

## MEDICAL POLICY

<b>POLICY TITLE</b>	<b>GENETIC TESTING FOR DEVELOPMENTAL DELAY/INTELLECTUAL DISABILITY, AUTISM SPECTRUM DISORDER, AND CONGENITAL ANOMALIES</b>
<b>POLICY NUMBER</b>	<b>MP 2.242</b>

<b>CLINICAL BENEFIT</b>	<input type="checkbox"/> MINIMIZE SAFETY RISK OR CONCERN. <input checked="" type="checkbox"/> MINIMIZE HARMFUL OR INEFFECTIVE INTERVENTIONS. <input type="checkbox"/> ASSURE APPROPRIATE LEVEL OF CARE. <input type="checkbox"/> ASSURE APPROPRIATE DURATION OF SERVICE FOR INTERVENTIONS. <input checked="" type="checkbox"/> ASSURE THAT RECOMMENDED MEDICAL PREREQUISITES HAVE BEEN MET. <input type="checkbox"/> ASSURE APPROPRIATE SITE OF TREATMENT OR SERVICE.
<b>Effective Date:</b>	<b>RETIRED 7/1/2026</b>

[POLICY RATIONALE](#)  
[CODING INFORMATION](#)

[PRODUCT VARIATIONS](#)  
[DEFINITIONS](#)  
[REFERENCES](#)

[DESCRIPTION/BACKGROUND](#)  
[DISCLAIMER](#)  
[POLICY HISTORY](#)

### I. POLICY

Chromosomal microarray analysis may be considered **medically necessary** in individuals with any of the following:

- Apparently nonsyndromic developmental delay/intellectual disability
- Autism spectrum disorder
- Multiple congenital anomalies not specific to a well-delineated genetic syndrome

Chromosomal microarray is considered **investigational** for the evaluation of all other conditions of delayed development, including but not limited to idiopathic growth or language delay. There is insufficient evidence to support a general conclusion concerning the health outcomes or benefits associated with this procedure for this indication.

Panel testing using next-generation sequencing is considered **investigational** in all cases of suspected genetic abnormality in children with developmental delay/intellectual disability, autism spectrum disorder, or congenital anomalies. There is insufficient evidence to support a general conclusion concerning the health outcomes or benefits associated with this procedure for this indication.

#### POLICY GUIDELINES

Use of chromosomal microarray (CMA) testing as outlined in this policy is not intended for use in the prenatal period. For trio testing and rapid WES/WGS please see **MP 2.324 Whole Exome and Whole Genome Sequencing for Diagnosis of Genetic Disorders**.

A guideline update from American College of Medical Genetics (ACMG, Schaefer et al [2013]) states that a stepwise (or tiered) approach to the clinical genetic diagnostic evaluation of autism spectrum disorder is recommended, with the recommendation being for first tier to include fragile X syndrome and CMA testing. CMA testing is recommended as first-tier evaluation in individuals who have the following:

- “Multiple anomalies not specific to a well-delineated genetic syndrome.”

## MEDICAL POLICY

<b>POLICY TITLE</b>	<b>GENETIC TESTING FOR DEVELOPMENTAL DELAY/INTELLECTUAL DISABILITY, AUTISM SPECTRUM DISORDER, AND CONGENITAL ANOMALIES</b>
<b>POLICY NUMBER</b>	<b>MP 2.242</b>

- “Apparently non-syndromic DD [developmental delay]/ID [intellectual disability].”
- “Autism spectrum disorders”

Recommendations from the American College of Medical Genetics (Manning and Hudgins [2010]) on array-based technologies and their clinical utilization for detecting chromosomal abnormalities include the following: “Appropriate follow-up is recommended in cases of chromosome imbalance identified by CMA, to include cytogenetic/FISH [fluorescent in situ hybridization] studies of the patient, parental evaluation, and clinical genetic evaluation and counseling.”

The International Standard Cytogenomic Array Consortium (ISCA, 2010) recommends offering CMA as a first-tier genetic test, in place of karyotype, for individuals with unexplained developmental delay/intellectual disability, autism spectrum disorders, or birth defects.

In some cases of CMA analysis, the laboratory performing the test confirms all reported copy number variants with an alternative technology such as fluorescent in situ hybridization analysis.

Per the American Academy of Pediatrics, chromosomal microarray and fragile X testing are recommended for all children with ASD to predict prognosis. Chromosomal microarray will reveal genetic abnormalities in up to 42% of children with ASD. Fragile X testing is positive in less than 1% of patients with ASD, but it is important for genetic counseling. Targeted testing for disorders such as tuberous sclerosis and Rett syndrome is useful only if presentation suggests these disorders. Whole exome sequencing shows an abnormality in up to one-fifth of patients with ASD and can be considered if other testing is negative.

In 2020, the Pediatric Exome Sequencing/Genome Sequencing Guideline Work Group (Peds ES/GS GWG) was convened to develop an evidence-based guideline for the clinical use of ES/GS in patients with CA/DD/ID. They strongly recommended exome sequencing and genome sequencing as a first-tier or second-tier test (guided by clinical judgment and often clinician-patient/ family shared decision making after CMA or focused testing) for patients with one or more CAs prior to one year of age or for patients with DD/ID with onset prior to 18 years of age. This recommendation is echoed by the ACMG.

ACMG (2012) stated “Before initiating GS/ES, counseling should be performed by a medical geneticist or an affiliated genetic counselor and should include written documentation of consent from the patient.”

### GENETICS NOMENCLATURE UPDATE

The Human Genome Variation Society nomenclature is used to report information on variants found in DNA and serves as an international standard in DNA diagnostics. The nomenclature is being implemented for genetic testing medical evidence review updates starting in 2017 (see Table PG1). The Society’s nomenclature is recommended by the Human Variome Project, the HUman Genome Organization, and by the Human Genome Variation Society itself.

The American College of Medical Genetics and Genomics and the Association for Molecular Pathology standards and guidelines for interpretation of sequence variants represent expert opinion from both organizations, in addition to the College of American Pathologists. These recommendations primarily apply to genetic tests used in clinical laboratories, including

## MEDICAL POLICY

<b>POLICY TITLE</b>	<b>GENETIC TESTING FOR DEVELOPMENTAL DELAY/INTELLECTUAL DISABILITY, AUTISM SPECTRUM DISORDER, AND CONGENITAL ANOMALIES</b>
<b>POLICY NUMBER</b>	<b>MP 2.242</b>

genotyping, single genes, panels, exomes, and genomes. Table PG2 shows the recommended standard terminology—“pathogenic,” “likely pathogenic,” “uncertain significance,” “likely benign,” and “benign”—to describe variants identified that cause Mendelian disorders.

**Table PG1. Nomenclature to Report on Variants Found in DNA**

<b>Previous</b>	<b>Updated</b>	<b>Definition</b>
<b>Mutation</b>	Disease-associated variant	Disease-associated change in the DNA sequence
	Variant	Change in the DNA sequence
	Familial variant	Disease-associated variant identified in a proband for use in subsequent targeted genetic testing in first-degree relatives

**Table PG2. ACMG-AMP Standards and Guidelines for Variant Classification**

<b>Variant Classification</b>	<b>Definition</b>
<b>Pathogenic</b>	Disease-causing change in the DNA sequence
<b>Likely pathogenic</b>	Likely disease-causing change in the DNA sequence
<b>Variant of uncertain significance</b>	Change in DNA sequence with uncertain effects on disease
<b>Likely benign</b>	Likely benign change in the DNA sequence
<b>Benign</b>	Benign change in the DNA sequence

ACMG: American College of Medical Genetics and Genomics; AMP: Association for Molecular Pathology

### GENETIC COUNSELING

Experts recommend formal genetic counseling for patients who are at risk for inherited disorders and who wish to undergo genetic testing. Interpreting the results of genetic tests and understanding risk factors can be difficult for some patients; genetic counseling helps individuals understand the impact of genetic testing, including the possible effects the test results could have on the individual or their family members. It should be noted that genetic counseling may alter the utilization of genetic testing substantially and may reduce inappropriate testing; further, genetic counseling should be performed by an individual with experience and expertise in genetic medicine and genetic testing methods. counseling should be performed by an individual with experience and expertise in genetic medicine and genetic testing methods.

**Cross-References:**

**MP 2.276 Genetic Testing for Pathogenic FMR1 Variants (Including Fragile X Syndrome)**

**MP 2.304 Autism Spectrum Disorders**

**MP 2.324 Whole Exome and Whole Genome Sequencing for Diagnosis of Genetic Disorders**

**MP 7.009 Preimplantation Genetic Testing**

## MEDICAL POLICY

<b>POLICY TITLE</b>	<b>GENETIC TESTING FOR DEVELOPMENTAL DELAY/INTELLECTUAL DISABILITY, AUTISM SPECTRUM DISORDER, AND CONGENITAL ANOMALIES</b>
<b>POLICY NUMBER</b>	<b>MP 2.242</b>

### II. PRODUCT VARIATIONS

[TOP](#)

This policy is only applicable to certain programs and products administered by Capital Blue Cross and subject to benefit variations as discussed in Section VI. Please see additional information below.

**FEP PPO** - Refer to FEP Medical Policy Manual. The FEP Medical Policy manual can be found at:

<https://www.fepblue.org/benefit-plans/medical-policies-and-utilization-management-guidelines/medical-policies>

### III. DESCRIPTION/BACKGROUND

[TOP](#)

#### Developmental Delay/Intellectual Disability

Developmental delay (DD), intellectual disability (ID), and congenital anomalies (CA) are among the most common indications for genetic referral in the pediatric population and comprise a heterogeneous group of conditions that can impact a child's physical, learning, or behavioral function. Identification of an underlying diagnosis for DD/ID/CA can lead to changes in management that will influence mortality, morbidity, and reduce the burden on patients and families searching for answers

Developmental delay is diagnosed in children 5 years or younger who show a significant delay in two or more developmental domains: gross or fine motor, speech/language, cognitive, social/personal, and activities of daily living. DD can precede the development of intellectual disability (ID) as the child ages.

ID is manifest by significant limitations in intellectual functioning and adaptive behavior. It is diagnosed at or after age 5 (when intelligence testing is considered valid and reliable) but prior to age 18 and is lifelong. The *Diagnostic and Statistical Manual of Mental Disorders: Fifth Edition (DSM-5)* defines ID as occurring during the developmental period and involving impairments of general mental abilities (e.g., IQ less than 70 or 75) that impact adaptive functioning in the conceptual, social, and practical domains. The causes of ID are extensive and include conditions that interfere with brain development and functioning. Among the known causes of ID, the majority are genetic abnormalities.

The national prevalence of DD and ID were estimated at 4.1% and 1.2%, respectively, in US children based on data from the 2009 to 2017 National Health Interview Survey. Both are influenced by genetic, environmental, infectious, and perinatal factors. Approximately 450 genes have been causally related to ID; most genes (~90%) are associated with syndromes. Inheritance of ID can be autosomal-dominant, recessive, or X-linked; and most nonsyndromic genes are located on the X chromosome. Prior to the advent of whole-exome and genome sequencing, Willemsen and Kleefstra (2014) concluded that 20% to 40% of ID cases could be attributed to a genetic variant. With the use of whole-genome sequencing, they estimated almost 60% of cases have an identifiable genetic etiology.

## MEDICAL POLICY

<b>POLICY TITLE</b>	<b>GENETIC TESTING FOR DEVELOPMENTAL DELAY/INTELLECTUAL DISABILITY, AUTISM SPECTRUM DISORDER, AND CONGENITAL ANOMALIES</b>
<b>POLICY NUMBER</b>	<b>MP 2.242</b>

Congenital anomalies are frequently present in children with DD and ID. In addition, a suspected etiology can often be established from history and physical examination (in skilled specialists as much as 20% to 40% of cases) without genetic testing. The recommended approach to evaluation in DD/ID begins with a 3-generation family history and physical (including neurologic) exam. Subsequent testing is used to confirm a suspected diagnosis (e.g., targeted fluorescent in situ hybridization [FISH] testing for DiGeorge or cri-du-chat syndromes). If no diagnosis is suspected, fragile X syndrome testing, metabolic testing for inborn errors of metabolism, and chromosomal microarray (CMA) testing (without karyotyping) are recommended-regardless of the presence or absence of dysmorphic features or congenital anomalies.

### Autism Spectrum Disorder

DSM-5 defines autism spectrum disorder (ASD) as the presence of

- Persistent deficits in social communication and social interaction across multiple contexts,
- Restricted, repetitive patterns of behavior, interests, or activities,
- Symptoms in the early developmental period (typically recognized in the first 2 years of life), and
- Symptoms cause clinically significant impairment in social, occupational, or other important areas of current functioning.

The estimated prevalence of ASD in US children based on data from the 2009 to 2017 National Health Interview Survey was 2.5%. ASD is 4 to 5 times more common in boys than girls, and white children are more often identified with ASD than black or Hispanic children. An accurate diagnosis can generally be made by age 2. The evaluation includes developmental screening and diagnostic evaluation (i.e., hearing, vision, neurologic, laboratory testing for metabolic disorders, and genetic testing).

A large body of evidence supports a genetic etiology in ASD. Twin studies estimate heritability between 60% and 90%. A family with an affected child has a 13% to 19% risk for recurrence in subsequent children. Based on Swedish genetic studies, Gaugler et al (2014) concluded that “the bulk of autism arises from genetic variation” (as opposed to environmental causes). Still, although genetic determinants can be heritable, most appear to arise de novo.

For these reasons, a child with ASD is often evaluated with genetic testing. Testing may be targeted when a child has a recognizable syndrome such as those shown in

Table 1. Alternatively, high-resolution cytogenetic analysis evaluating multiple genes-the focus of this evidence review-is used.

**Table 1. Examples of Specific Genes Associated With Disorders That Include Autistic Behaviors**

Gene (Syndrome)	Patient Selection	Yield, %	Reference
<i>FMR1</i> (fragile X)	Unselected autism	3-10	Schaefer and Mendelsohn

## MEDICAL POLICY

<b>POLICY TITLE</b>	<b>GENETIC TESTING FOR DEVELOPMENTAL DELAY/INTELLECTUAL DISABILITY, AUTISM SPECTRUM DISORDER, AND CONGENITAL ANOMALIES</b>
<b>POLICY NUMBER</b>	<b>MP 2.242</b>

<b>MECP2 (Rett)</b>	Females with nonsyndromic autism, intellectual disability, and cerebral palsy	3-13	(2008) <sup>24</sup>
<b>PTEN</b>	Autism with macrocephaly	≤17	Butler et al (2005) <sup>25</sup>

### Diagnostic Testing

#### Karyotyping and Fluorescent In Situ Hybridization

The goal of a cytogenetic evaluation is to identify chromosomal imbalances that cause a disorder. The most common imbalances are copy number variants (CNVs) or deletions and duplications of large segments of genomic material. CNVs are common in DD/ID and ASD but more often reflect the normal genetic variation. However, de novo CNVs are observed about four times more frequently in children with ASD than in normal individuals. Less frequently, other abnormalities such as balanced translocations (i.e., exchanges of equally sized DNA loci between chromosomes) may be pathogenic. For many well-described syndromes, the type and location of the associated chromosomal abnormality have been established by studying large patient samples. For others, few patients with similar abnormalities may have been evaluated to establish genotype-phenotype correlation. Finally, in some patients, cytogenetic analysis will discover chromosomal abnormalities that require study to determine their significance.

Prior to the advent of CMAs, the initial step in cytogenetic analysis was G-banded karyotyping, which evaluates all chromosomes. High-resolution G-banding can detect changes as small as 3 to 5 megabases in size, although standard G-banding evaluates more than ten megabases changes. In children with DD/ID, a review by Stankiewicz and Beaudet (2007) found G-banded karyotyping diagnostic in approximately 3% to 5% of cases. In ASD, high-resolution karyotyping appears to identify abnormalities in up to 5% of cases.

Chromosomal rearrangements in the gene-rich subtelomeric region are identified in approximately 4 to 6 percent of children with ID. Molecular screening using FISH of subtelomeric probes was used widely in the past to identify these abnormalities; however, CMA has replaced FISH as the test of choice, since the majority of diagnostic CMA arrays offer dense coverage of subtelomeric regions. FISH may still be substituted if array diagnosis is not available or if a specific telomeric disorder (e.g., DiGeorge syndrome, Cri-du-chat syndrome) is strongly suspected clinically.

In contrast, molecular cytogenetic techniques can detect small submicroscopic chromosomal alterations. Fluorescent in situ hybridization (FISH) a targeted approach, is used to identify specific chromosomal abnormalities associated with suspected diagnoses such as DiGeorge syndrome. Prior to CMAs, FISH was also used to screen the rearrangement-prone subtelomeric regions. Subtelomeric FISH was found to identify abnormalities in children with DD and ID, diagnostic in approximately 5% to 6% of those with negative karyotypes, but uncommonly in ASD.

#### Chromosomal Microarrays

## MEDICAL POLICY

<b>POLICY TITLE</b>	<b>GENETIC TESTING FOR DEVELOPMENTAL DELAY/INTELLECTUAL DISABILITY, AUTISM SPECTRUM DISORDER, AND CONGENITAL ANOMALIES</b>
<b>POLICY NUMBER</b>	<b>MP 2.242</b>

Two types of CMAs are considered here: array comparative genomic hybridization (aCGH) and single nucleotide variants (SNV) arrays. The aCGH approach uses DNA samples from a patient and a normal control. Each is labeled with distinct fluorescent dyes (red or green). The labeled samples are then mixed and hybridized to thousands of cloned or synthesized reference (normal) DNA fragments of known genomic locus immobilized on a glass slide (microarray) to conduct thousands of comparative reactions simultaneously. CNVs are determined by computer analysis of the array patterns and intensities of the hybridization signals. If the patient sequence is missing part of the normal sequence (a deletion) or has the normal sequence plus additional genomic material within that genomic location (e.g., a duplication), the sequence imbalance is detected as a difference in fluorescence intensity (Korf and Rehm [2013] offer an illustrative graphic). For this reason, aCGH cannot detect balanced chromosomal translocations (equal exchange of material between chromosomes) or sequence inversions (same sequence is present in reverse base pair order) because the fluorescence intensity would not change. A portion of the increased diagnostic yield from CMA over karyotyping comes from the discovery that chromosomal rearrangements that appear balanced (and therefore not pathogenic) by G-banded karyotype analysis are found to have small imbalances with greater resolution. It has been estimated that 40% of apparently balanced de novo or inherited translocations with abnormal phenotype are associated with cryptic deletion if analyzed by CMA testing.

Like aCGH, SNV arrays detect CNVs. In an SNV array, the two alleles for genes of interest are tagged with different fluorescent dyes. Comparative fluorescence intensity will be increased when there are duplications and diminished with deletions. The resolution provided by aCGH is higher than with SNV arrays. In addition, aCGH has better signal-to-background characteristics than SNV arrays. In contrast to aCGH, SNV arrays will also identify long stretches of DNA homozygosity, which may suggest uniparental disomy or consanguinity. Uniparental disomy occurs when a child inherits two copies of a chromosome from one parent and no copies from the other parent. Uniparental disomy can lead to syndromes such as Angelman and Prader-Willi.

Table 2 summarizes the cytogenetic tests used to evaluate children with DD/ID and autism. The table emphasizes the large difference in resolution between karyotyping and CMA.

**Table 2. Resolution and Analysis Comparison of FISH, Karyotyping, and CMA Analysis**

Test	Resolution in Kilobases <sup>a</sup>	Analysis
Karyotyping	3000-5000 kb	Genome-wide
CMA	≈50 kb	Genome-wide
FISH	≈500 to 1000 kb (depending on probe)	Targeted

CMA: chromosomal microarray; FISH: fluorescent in situ hybridization; kb: kilobases.

<sup>a</sup> One kb = 1000 bases, 1000 kb = 1 Mb.

Microarrays may be prepared by the laboratory using the technology or, more commonly, by commercial manufacturers, and sold to laboratories that must qualify and validate the product for use in their assay, in conjunction with computerized software for interpretation. The

## MEDICAL POLICY

<b>POLICY TITLE</b>	<b>GENETIC TESTING FOR DEVELOPMENTAL DELAY/INTELLECTUAL DISABILITY, AUTISM SPECTRUM DISORDER, AND CONGENITAL ANOMALIES</b>
<b>POLICY NUMBER</b>	<b>MP 2.242</b>

proliferation of laboratory-developed and commercially available platforms prompted the American College of Medical Genetics to publish guidelines for the design and performance expectations for clinical microarrays and associated software in the postnatal setting.

### Next-Generation Sequencing

Next-generation sequencing (NGS) has been proposed to detect single-gene causes of autism and possibly identify a syndrome that involves autism in patients with normal array-based testing. NGS involves the sequencing of millions of fragments of genetic material in a massively parallel fashion. NGS can be performed on segments of genetic material of various sizes—from the entire genome (whole-genome sequencing) to small subsets of genes (targeted sequencing). NGS allows the detection of SNVs, CNVs, insertions, and deletions. With higher resolution comes higher likelihood of detection of variants of uncertain significance.

### Exome and Genome Sequencing

Exome sequencing utilizes DNA-enrichment methods and massively parallel nucleotide sequencing to identify disease-associated variants throughout the human genome. Exome sequencing is limited to the DNA sequence of coding regions (exons) and flanking intronic regions of the genome, which is estimated to contain 85% of heritable disease-causing variants. Results of testing with ES include known pathogenic variants definitely associated with disease or a variant of uncertain significance. Due to the falling costs of sequencing and its high diagnostic yield, WES is rapidly becoming a clinical tool for the evaluation of ID, especially at specialty centers. Exome sequencing after standard testing increased the diagnostic yield at an additional cost compared to standard testing alone. However, using ES as a first- or second-tier test (e.g., after CMA or targeted testing) yielded more diagnoses at a lower cost than using ES only after extensive standard testing (e.g., large sequencing panels and/or multiple testing approaches) or using standard testing alone. With the anticipated further declines in cost, early use of genome-wide sequencing should continue to enable more timely diagnosis for patients with unexplained DD or multiple Cas.

Whole genome sequencing is costlier than more limited sequencing, because the whole genome is equivalent to approximately  $3.3 \times 10^9$  bases (3.3 gigabases [Gb]). Whole genome sequencing may become preferable to exome sequencing as cost decreases and more information about the role of non-coding DNA in human disease becomes available. Further, whole genome sequencing can be used to detect deletions and duplications typically detected only by array comparative genomic hybridization (aCGH) and thus is an opportunity to reduce the need for other supplemental testing. One of the most common medical indications for whole genome sequencing or whole exome sequencing is evaluation of severe intellectual disability or developmental delay believed to have a genetic etiology in a child with a negative initial evaluation.

The value of genetic counseling in exome and genome sequencing is well-established. Creating reasonable expectations, establishing an understanding of the value and limitations of testing, creating awareness of the potential harms, and allowing the family to make informed choices is a mainstay of informed consent for ES/GS. These visits should also be commensurate with the

## MEDICAL POLICY

<b>POLICY TITLE</b>	<b>GENETIC TESTING FOR DEVELOPMENTAL DELAY/INTELLECTUAL DISABILITY, AUTISM SPECTRUM DISORDER, AND CONGENITAL ANOMALIES</b>
<b>POLICY NUMBER</b>	<b>MP 2.242</b>

time spent as part of the clinical process including reimbursement for this type of counseling. Post-test counseling extends this benefit once the results are available regardless of the diagnostic yield. Elements of counseling should include a three-generation family pedigree; discussion of pathogenic/likely pathogenic results, benign results, and variants of uncertain significance; detection of misattributed paternity or consanguinity, and secondary findings unrelated to the reason for testing.

### Genetic Associations with DD/ID and ASD

For common phenotypes and syndromes, the pathogenicity of CNVs may be supported by considerable evidence; for uncommon phenotypes and uncommon CNVs determining pathogenicity requires a systematic evaluation that includes parental studies, examining databases for reported associations, and considering the molecular consequences of the identified variant. Parental studies (e.g., “trio” testing of affected child, father, and mother) can identify an inherited CNV from an unaffected parent and therefore considered benign. A variety of databases index the clinical implications of CNVs and their associations with a particular phenotype. CNVs are continuously cataloged and, with growth in CMA testing and improved resolution, databases have become increasingly extensive (e.g., DECIPHER, ClinVar). For uncommon CNVs, in addition to reports of CNV-phenotype associations, the location and size of the CNV can offer clues to pathogenicity; larger CNVs are more often pathogenic and the role of affected genes in brain circuitry and effect of CNV on gene expression can implicate pathogenicity. Although uncommon, an observed phenotype can result from unmasking a mutated recessive allele on the unaffected (non-CNV) chromosome. Other considerations when determining pathogenicity include CNV dosage, X linkage, number of reports in the literature of an association between CNV and phenotype, and findings in “normal” individuals.

The American College of Medical Genetics has published guidelines for evaluating, interpreting, and reporting pathogenicity reflecting these principles. The recommended categories of clinical significance for reporting are pathogenic, uncertain clinical significance (likely pathogenic, likely benign, or no subclassification), or benign. The International Standards for Cytogenomic Arrays Consortium more recently proposed “an evidence-based approach to guide the development of content on chromosomal microarrays and to support interpretation of clinically significant copy number variation.” The proposal defined levels of evidence describe how well or how poorly detected variants or CNVs correlate with phenotype.

### Regulatory Status

Clinical laboratories may develop and validate tests in-house and market them as a laboratory service; laboratory-developed tests must meet the general regulatory standards of the Clinical Laboratory Improvement Amendments. Lab tests for CMA testing and NGS are available under the auspices of Clinical Laboratory Improvement Amendments. Laboratories that offer laboratory-developed tests must be licensed by the Clinical Laboratory Improvement Amendments for high-complexity testing. To date, the U.S. Food and Drug Administration (FDA) has chosen not to require any regulatory review of this test.

## MEDICAL POLICY

<b>POLICY TITLE</b>	<b>GENETIC TESTING FOR DEVELOPMENTAL DELAY/INTELLECTUAL DISABILITY, AUTISM SPECTRUM DISORDER, AND CONGENITAL ANOMALIES</b>
<b>POLICY NUMBER</b>	<b>MP 2.242</b>

In 2010, FDA indicated that it would require microarray manufacturers to seek clearance to sell their products for use in clinical cytogenetics.

### CMA Testing

CMA testing is commercially available through many laboratories and includes targeted and whole- genome arrays, with or without SNV microarray analysis.

In January 2014, the Affymetrix CytoScan® Dx Assay (Thermo Fisher Scientific) was cleared by FDA through the de novo 510(k) process. FDA's review of the CytoScan® Dx Assay included an analytic evaluation of the test's ability to detect accurately numerous chromosomal variations of different types, sizes, and genome locations compared with several analytically validated test methods. The FDA found that the CytoScan® Dx Assay could detect CNVs across the genome and adequately detect CNVs in regions of the genome associated with ID/DD. Reproducibility decreased with the CNV gain or loss size, particularly when less than approximately four hundred kilobases (generally recommended as the lower reporting limit). As of July 2017, Affymetrix™ has reported 2.7 million markers for copy number, 750,000 SNVs and 1.9 million polymorphic probes (Affymetrix™ was acquired by Thermo Fisher Scientific in 2016). FDA product code: PFX.

FirstStep<sup>Dx</sup> PLUS® (Lineagen) uses Lineagen's custom-designed microarray platform manufactured by Affymetrix. As of July 2017, this microarray consists of a 2.8 million probe microarray for the detection of CNVs associated with neurodevelopmental disorders. The array includes probes that come standard on the Affymetrix CytoScan HD® microarray, with an additional 88,435 custom probes designed by Lineagen.

Ambry Genetics offers multiple tests (CMA and NGS) designed for diagnosing ASD and neurodevelopmental disorders. As of July 2017, the CMA offered by Ambry Genetics includes over 2.6 million probes for copy number and 750,000 SNV probes. The expanded NGS panel for neurodevelopmental disorders includes assesses 196 genes.

LabCorp offers the Reveal® SNP Microarray-Pediatric for individuals with nonsyndromic congenital anomalies, dysmorphic features, DD/ID, and/or ASD. The Reveal® microarray has 2695 million probes as of July 2017.

### Next-Generation Sequencing

A variety of commercial and academic laboratories offer NGS panels designed for the evaluation of ASD, DD/ID, and congenital anomalies, which vary in terms of the numbers of, and specific genes tested.

Emory Genetics Laboratory offers an NGS ASD panel of genes targeting genetic syndromes that include autism or autistic features. Greenwood Genetics Center offers an NGS panel for syndromic autism that includes eighty-three genes. Fulgent Genetics offers a next-generation sequencing ASD panel that includes 121 genes.

## MEDICAL POLICY

<b>POLICY TITLE</b>	<b>GENETIC TESTING FOR DEVELOPMENTAL DELAY/INTELLECTUAL DISABILITY, AUTISM SPECTRUM DISORDER, AND CONGENITAL ANOMALIES</b>
<b>POLICY NUMBER</b>	<b>MP 2.242</b>

### IV. RATIONALE

[TOP](#)

#### Summary of Evidence

For individuals who have DD/ID, ASD, or multiple congenital anomalies not specific to a well-delineated genetic syndrome who receive CMA testing, the evidence includes primarily case series. Relevant outcomes are test accuracy and validity, changes in reproductive decision making, morbid events, and resource utilization. The available evidence supports test validity. Although systematic studies of the impact of CMA on patient outcomes are lacking, the improvement in diagnostic yield over karyotyping has been well-demonstrated. Direct evidence of improved outcomes with CMA compared with karyotyping is lacking. However, for at least a subset of the disorders potentially diagnosed with CMA testing in this patient population, there are well-defined and accepted management steps associated with positive test results. Further, there is evidence of changes in reproductive decision making as a result of positive test results. The information derived from CMA testing can accomplish the following: it could end a long diagnostic odyssey; or reduce morbidity for certain conditions by initiating surveillance/management of associated comorbidities; or it could potentially impact future reproductive decision making for parents. The evidence is sufficient to determine that the technology results in a meaningful improvement in the net health outcome.

For exome and genome sequencing, the growing body of literature provides justification for a strong recommendation based desirable effects and limited harms. The evidence is sufficient to determine that the technology results in meaningful improvement in the net health outcome.

For individuals who have DD/ID, ASD, or multiple congenital anomalies not specific to a well-delineated genetic syndrome who receive next-generation sequencing panel testing, the evidence includes primarily case series. Relevant outcomes are test accuracy and validity, changes in reproductive decision making, morbid events, and resource utilization. The diagnostic yield associated with next-generation sequencing panel testing in this patient population is not well-characterized. The testing yield and likelihood of uncertain results are variable, based on gene panel, gene tested, and patient population; additionally, there are risks of uninterpretable and incidental results. The evidence is insufficient to determine the effects of the technology on health outcomes.

### V. DEFINITIONS

[TOP](#)

**ANALYTIC VALIDITY** of a genetic test defines its ability to accurately and reliably measure the genotype of interest.

**CHROMOSOME** is one of the threadlike “packages” of genes and other DNA in the nucleus of a cell.

**CLINICAL VALIDITY** of a genetic test defines its ability to detect or predict the associated disorder (phenotype).

## MEDICAL POLICY

<b>POLICY TITLE</b>	<b>GENETIC TESTING FOR DEVELOPMENTAL DELAY/INTELLECTUAL DISABILITY, AUTISM SPECTRUM DISORDER, AND CONGENITAL ANOMALIES</b>
<b>POLICY NUMBER</b>	<b>MP 2.242</b>

**COMPARATIVE GENOMIC HYBRIDIZATION (CGH)** also referred to as chromosomal microarray analysis (CMA), and array CGH (aCGH) is a technique which produces a map of DNA sequence copy number as a function of chromosomal location throughout the entire genetic genome, and allows the detection of genetic deletions, duplications, and amplifications.

**DNA** a large nucleic acid molecule, found principally in the chromosomes of the nucleus of a cell, that is the carrier of genetic information.

**FIRST-DEGREE RELATIVE** refers to a parent, sibling, or child.

**GENE** is the basic unit of heredity, made of DNA, the code for a specific protein.

**GENOTYPE** is the specific genetic makeup of an individual, usually in the form of DNA.

**KARYOTYPE** is the chromosomal complement of an individual, including the number of chromosomes and any abnormalities.

**MICROARRAY** is a tool for analyzing gene expression that consists of a small membrane or glass slide containing samples of many genes arranged in a regular pattern. Each spot on an array is associated with a particular gene. Each color in an array represents either healthy (control) or diseased (sample) tissue. Depending on the type of array used, the location and intensity of a color will indicate whether the gene, or mutation, is present in either the control and/or sample DNA. It will also provide an estimate of the expression level of the gene(s) in the sample and control DNA.

**MITOCHONDRIA** are intracellular organelles that are responsible for energy production and cellular respiration.

**MITOCHONDRIAL DISEASE** refers to one of hundreds of congenital illnesses that result from mutations in mitochondrial DNA. As a result, the mitochondria are unable to completely burn food and oxygen in order to generate energy.

**MUTATION** is a permanent structural alteration in DNA.

**PHENOTYPE** is the physical characteristics of an organism or the presence of a disease that may or may not be genetic.

**RNA** is a chemical similar to a single strand of DNA

### VI. **DISCLAIMER**

[TOP](#)

*Capital Blue Cross' medical policies are used to determine coverage for specific medical technologies, procedures, equipment, and services. These medical policies do not constitute medical advice and are subject to change as required by law or applicable clinical evidence from independent treatment guidelines. Treating providers are solely responsible for medical advice and treatment of members. These policies are not a guarantee of coverage or payment. Payment of claims is subject to a determination regarding the member's benefit program and eligibility on the date of service, and a determination that the services are medically necessary and appropriate. Final processing of a claim is based upon the terms of contract that applies to the members' benefit program, including benefit limitations and*

## MEDICAL POLICY

<b>POLICY TITLE</b>	<b>GENETIC TESTING FOR DEVELOPMENTAL DELAY/INTELLECTUAL DISABILITY, AUTISM SPECTRUM DISORDER, AND CONGENITAL ANOMALIES</b>
<b>POLICY NUMBER</b>	<b>MP 2.242</b>

*exclusions. If a provider or a member has a question concerning this medical policy, please contact Capital Blue Cross' Provider Services or Member Services.*

### VII. CODING INFORMATION

[TOP](#)

**Note:** This list of codes may not be all-inclusive, and codes are subject to change at any time. The identification of a code in this section does not denote coverage as coverage is determined by the terms of member benefit information. In addition, not all covered services are eligible for separate reimbursement.

#### Investigational; therefore, not covered.

Procedure Codes								
0156U	0170U							

#### Covered when medically necessary:

Procedure Codes								
S3870	0209U	0318U	81228	81229	81349	81470	81471	

ICD-10-CM Diagnosis Codes	Description
F70	Mild intellectual disabilities
F71	Moderate intellectual disabilities
F72	Severe intellectual disabilities
F73	Profound intellectual disabilities
F78	Other intellectual disabilities
F78.A1	SYNGAP1-related intellectual disability
F78.A9	Other genetic related intellectual disability
F80.0	Phonological disorder
F82	Specific developmental disorder of motor function
F84.0	Autistic disorder
F84.2	Rett's syndrome
F84.3	Other childhood disintegrative disorder
F84.5	Asperger's syndrome
F84.8	Other pervasive developmental disorders
F88	Other disorders of psychological development
Q89.7	Multiple congenital malformations, not elsewhere classified
QA0	Neurodevelopmental disorders related to specific genetic pathogenic variants

## MEDICAL POLICY

<b>POLICY TITLE</b>	<b>GENETIC TESTING FOR DEVELOPMENTAL DELAY/INTELLECTUAL DISABILITY, AUTISM SPECTRUM DISORDER, AND CONGENITAL ANOMALIES</b>
<b>POLICY NUMBER</b>	<b>MP 2.242</b>

<b>ICD-10-CM Diagnosis Codes</b>	<b>Description</b>
QA0.0	Neurodevelopmental disorders related to pathogenic variants in specific genes
QA0.01	Neurodevelopmental disorders related to pathogenic variants in certain specific genes
QA0.010	Neurodevelopmental disorders, related to pathogenic variants in ion channel genes
QA0.013	Neurodevelopmental disorders, related to pathogenic variants in other transporter and solute carrier genes
QA0.014	Neurodevelopmental disorders, related to pathogenic variants in synapse related genes
QA0.015	Neurodevelopmental disorders, related to genes associated with transcription and gene expression

### VIII. REFERENCES

[TOP](#)

1. Mikhail FM, Lose EJ, Robin NH, et al. Clinically relevant single gene or intragenic deletions encompassing critical neurodevelopmental genes in patients with developmental delay, mental retardation, and/or autism spectrum disorders. *Am J Med Genet A*. Oct 2011; 155A(10): 2386-96. PMID 22031302
2. Brandler WM, Sebat J. From de novo mutations to personalized therapeutic interventions in autism. *Annu Rev Med*. 2015; 66: 487-507. PMID 25587659
3. Stankiewicz P, Beaudet AL. Use of array CGH in the evaluation of dysmorphism, malformations, developmental delay, and idiopathic mental retardation. *Curr Opin Genet Dev*. Jun 2007; 17(3): 182-92. PMID 17467974
4. Stuart SW, King CH, Pai GS. Autism spectrum disorder, Klinefelter syndrome, and chromosome 3p21.31 duplication: a case report. *MedGenMed*. Dec 18, 2007; 9(4): 60. PMID 18311409
5. Moeschler JB. Medical genetics diagnostic evaluation of the child with global developmental delay or intellectual disability. *Curr Opin Neurol*. Apr 2008; 21(2): 117-22. PMID 18317267
6. Schaefer GB, Mendelsohn NJ. Clinical genetics evaluation in identifying the etiology of autism spectrum disorders. *Genet Med*. Apr 2008; 10(4): 301-5. PMID 18414214
7. Korf BR, Rehm HL. New approaches to molecular diagnosis. *JAMA*. Apr 10, 2013; 309(14): 1511-21. PMID 23571590
8. Kearney HM, South ST, Wolff DJ, et al. American College of Medical Genetics recommendations for the design and performance expectations for clinical genomic copy number microarrays intended for use in the postnatal setting for detection of constitutional abnormalities. *Genet Med*. Jul 2011; 13(7): 676-9. PMID 21681105
9. Rodriguez-Revenga L, Vallespin E, Madrigal I, et al. A parallel study of different array-CGH platforms in a set of Spanish patients with developmental delay and intellectual disability. *Gene*. May 25, 2013; 521(1): 82-6. PMID 23524024

## MEDICAL POLICY

<b>POLICY TITLE</b>	<b>GENETIC TESTING FOR DEVELOPMENTAL DELAY/INTELLECTUAL DISABILITY, AUTISM SPECTRUM DISORDER, AND CONGENITAL ANOMALIES</b>
<b>POLICY NUMBER</b>	<b>MP 2.242</b>

10. Kloosterman WP, Hochstenbach R. Deciphering the pathogenic consequences of chromosomal aberrations in human genetic disease. *Mol Cytogenet.* 2014; 7(1): 100. PMID 25606056
11. Kearney HM, Thorland EC, Brown KK, et al. American College of Medical Genetics standards and guidelines for interpretation and reporting of postnatal constitutional copy number variants. *Genet Med.* Jul 2011; 13(7): 680-5. PMID 21681106
12. Riggs ER, Church DM, Hanson K, et al. Towards an evidence-based process for the clinical interpretation of copy number variation. *Clin Genet.* May 2012; 81(5): 403- 12. PMID 22097934
13. Moeschler JB, Shevell M, Moeschler JB, et al. Comprehensive evaluation of the child with intellectual disability or global developmental delays. *Pediatrics.* Sep 2014; Reaffirmed October 2019. 134(3): e903-18. PMID 25157020
14. Shevell M, Ashwal S, Donley D, et al. Practice parameter: evaluation of the child with global developmental delay: report of the Quality Standards Subcommittee of the American Academy of Neurology and The Practice Committee of the Child Neurology Society. *Neurology.* Feb 11, 2003; 60(3): 367-80. PMID 12578916
15. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders, 5th Edition: DSM-5.* Washington (DC): American Psychiatric Association; 2013.
16. Zablotsky B, Black LI, Maenner MJ, et al. Prevalence and Trends of Developmental Disabilities among Children in the United States: 2009-2017. *Pediatrics.* Oct 2019; 144(4). PMID 31558576
17. Srouf M, Shevell M. Genetics and the investigation of developmental delay/intellectual disability. *Arch Dis Child.* Apr 2014; 99(4): 386-9. PMID 24344174
18. Willemsen MH, Kleefstra T. Making headway with genetic diagnostics of intellectual disabilities. *Clin Genet.* Feb 2014; 85(2): 101-10. PMID 23895455
19. Moeschler JB. Genetic evaluation of intellectual disabilities. *Semin Pediatr Neurol.* Mar 2008; 15(1): 2-9. PMID 18342255
20. Yuen RK, Thiruvahindrapuram B, Merico D, et al. Whole-genome sequencing of quartet families with autism spectrum disorder. *Nat Med.* Feb 2015; 21(2): 185-91. PMID 25621899
21. Gaugler T, Klei L, Sanders SJ, et al. Most genetic risk for autism resides with common variation. *Nat Genet.* Aug 2014; 46(8): 881-5. PMID 25038753
22. Schaefer GB, Mendelsohn NJ. Genetics evaluation for the etiologic diagnosis of autism spectrum disorders. *Genet Med.* Jan 2008; 10(1): 4-12. PMID 18197051
23. Butler MG, Dasouki MJ, Zhou XP, et al. Subset of individuals with autism spectrum disorders and extreme macrocephaly associated with germline PTEN tumour suppressor gene mutations. *J Med Genet.* Apr 2005; 42(4): 318-21. PMID 15805158
24. Wright CF, Fitzgerald TW, Jones WD, et al. Genetic diagnosis of developmental disorders in the DDD study: a scalable analysis of genome-wide research data. *Lancet.* Apr 04, 2015; 385(9975): 1305-14. PMID 25529582
25. Eriksson MA, Lieden A, Westerlund J, et al. Rare copy number variants are common in young children with autism spectrum disorder. *Acta Paediatr.* Jun 2015; 104(6): 610-8. PMID 25661985

## MEDICAL POLICY

<b>POLICY TITLE</b>	<b>GENETIC TESTING FOR DEVELOPMENTAL DELAY/INTELLECTUAL DISABILITY, AUTISM SPECTRUM DISORDER, AND CONGENITAL ANOMALIES</b>
<b>POLICY NUMBER</b>	<b>MP 2.242</b>

26. *Krepischi-Santos AC, Vianna-Morgante AM, Jehee FS, et al. Whole-genome array-CGH screening in undiagnosed syndromic patients: old syndromes revisited and new alterations. Cytogenet Genome Res. 2006; 115(3-4): 254-61. PMID 17124408*
27. *Bartnik M, Nowakowska B, Derwinska K, et al. Application of array comparative genomic hybridization in 256 patients with developmental delay or intellectual disability. J Appl Genet. Feb 2014; 55(1): 125-44. PMID 24297458*
28. *Bartnik M, Wisniowiecka-Kowalnik B, Nowakowska B, et al. The usefulness of array comparative genomic hybridization in clinical diagnostics of intellectual disability in children. Dev Period Med. Jul-Sep 2014; 18(3): 307-17. PMID 25182394*
29. *Chong WW, Lo IF, Lam ST, et al. Performance of chromosomal microarray for patients with intellectual disabilities/developmental delay, autism, and multiple congenital anomalies in a Chinese cohort. Mol Cytogenet. 2014; 7: 34. PMID 24926319*
30. *D'Amours G, Langlois M, Mathonnet G, et al. SNP arrays: comparing diagnostic yields for four platforms in children with developmental delay. BMC Med Genomics. Dec 24, 2014; 7: 70. PMID 25539807*
31. *Henderson LB, Applegate CD, Wohler E, et al. The impact of chromosomal microarray on clinical management: a retrospective analysis. Genet Med. Sep 2014; 16(9): 657-64. PMID 24625444*
32. *Nava C, Keren B, Mignot C, et al. Prospective diagnostic analysis of copy number variants using SNP microarrays in individuals with autism spectrum disorders. Eur J Hum Genet. Jan 2014; 22(1): 71-8. PMID 23632794*
33. *Nicholl J, Waters W, Mulley JC, et al. Cognitive deficit and autism spectrum disorders: prospective diagnosis by array CGH. Pathology. Jan 2014; 46(1): 41-5. PMID 24300712*
34. *Palmer E, Speirs H, Taylor PJ, et al. Changing interpretation of chromosomal microarray over time in a community cohort with intellectual disability. Am J Med Genet A. Feb 2014; 164A(2): 377-85. PMID 24311194*
35. *Preiksaitiene E, Molyte A, Kasnauskiene J, et al. Considering specific clinical features as evidence of pathogenic copy number variants. J Appl Genet. May 2014; 55(2): 189-96. PMID 24535828*
36. *Redin C, Gerard B, Lauer J, et al. Efficient strategy for the molecular diagnosis of intellectual disability using targeted high-throughput sequencing. J Med Genet. Nov 2014; 51(11): 724-36. PMID 25167861*
37. *Roberts JL, Hovanes K, Dasouki M, et al. Chromosomal microarray analysis of consecutive individuals with autism spectrum disorders or learning disability presenting for genetic services. Gene. Feb 01, 2014; 535(1): 70-8. PMID 24188901*
38. *Stobbe G, Liu Y, Wu R, et al. Diagnostic yield of array comparative genomic hybridization in adults with autism spectrum disorders. Genet Med. Jan 2014; 16(1): 70-7. PMID 23765050*
39. *Tao VQ, Chan KY, Chu YW, et al. The clinical impact of chromosomal microarray on paediatric care in Hong Kong. PLoS One. 2014; 9(10): e109629. PMID 25333781*
40. *Utine GE, Haliloglu G, Volkan-Salanci B, et al. Etiological yield of SNP microarrays in idiopathic intellectual disability. Eur J Paediatr Neurol. May 2014; 18(3): 327-37. PMID 24508361*

## MEDICAL POLICY

<b>POLICY TITLE</b>	<b>GENETIC TESTING FOR DEVELOPMENTAL DELAY/INTELLECTUAL DISABILITY, AUTISM SPECTRUM DISORDER, AND CONGENITAL ANOMALIES</b>
<b>POLICY NUMBER</b>	<b>MP 2.242</b>

41. Uwineza A, Caberg JH, Hitayezu J, et al. Array-CGH analysis in Rwandan patients presenting development delay/intellectual disability with multiple congenital anomalies. *BMC Med Genet.* Jul 12, 2014; 15: 79. PMID 25016475
42. Battaglia A, Doccini V, Bernardini L, et al. Confirmation of chromosomal microarray as a first-tier clinical diagnostic test for individuals with developmental delay, intellectual disability, autism spectrum disorders and dysmorphic features. *Eur J Paediatr Neurol.* Nov 2013; 17(6): 589-99. PMID 23711909
43. Lee CG, Park SJ, Yun JN, et al. Array-based comparative genomic hybridization in 190 Korean patients with developmental delay and/or intellectual disability: a single tertiary care university center study. *Yonsei Med J.* Nov 2013; 54(6): 1463-70. PMID 24142652
44. Shoukier M, Klein N, Auber B, et al. Array CGH in patients with developmental delay or intellectual disability: are there phenotypic clues to pathogenic copy number variants? *Clin Genet.* Jan 2013; 83(1): 53-65. PMID 22283495
45. Sorte HS, Gjevik E, Sponheim E, et al. Copy number variation findings among 50 children and adolescents with autism spectrum disorder. *Psychiatr Genet.* Apr 2013; 23(2): 61-9. PMID 23277134
46. Filges I, Suda L, Weber P, et al. High-resolution array in the clinical approach to chromosomal phenotypes. *Gene.* Mar 10, 2012; 495(2): 163-9. PMID 22240311
47. Iourov IY, Vorsanova SG, Kurinnaia OS, et al. Molecular karyotyping by array CGH in a Russian cohort of children with intellectual disability, autism, epilepsy and congenital anomalies. *Mol Cytogenet.* Dec 31, 2012; 5(1): 46. PMID 23272938
48. McGrew SG, Peters BR, Crittendon JA, et al. Diagnostic yield of chromosomal microarray analysis in an autism primary care practice: which guidelines to implement? *J Autism Dev Disord.* Aug 2012; 42(8): 1582-91. PMID 22089167
49. Tzetis M, Kitsiou-Tzeli S, Frysira H, et al. The clinical utility of molecular karyotyping using high-resolution array-comparative genomic hybridization. *Expert Rev Mol Diagn.* Jun 2012; 12(5): 449-57. PMID 22702362
50. Bremer A, Giacobini M, Eriksson M, et al. Copy number variation characteristics in subpopulations of patients with autism spectrum disorders. *Am J Med Genet B Neuropsychiatr Genet.* Mar 2011; 156(2): 115-24. PMID 21302340
51. Coulter ME, Miller DT, Harris DJ, et al. Chromosomal microarray testing influences medical management. *Genet Med.* Sep 2011; 13(9): 770-6. PMID 21716121
52. Wincent J, Anderlid BM, Lagerberg M, et al. High-resolution molecular karyotyping in patients with developmental delay and/or multiple congenital anomalies in a clinical setting. *Clin Genet.* Feb 2011; 79(2): 147-57. PMID 20486943
53. Manolakos E, Vetro A, Kefalas K, et al. The use of array-CGH in a cohort of Greek children with developmental delay. *Mol Cytogenet.* Nov 09, 2010; 3: 22. PMID 21062444
54. Rosenfeld JA, Ballif BC, Torchia BS, et al. Copy number variations associated with autism spectrum disorders contribute to a spectrum of neurodevelopmental disorders. *Genet Med.* Nov 2010; 12(11): 694-702. PMID 20808228
55. Schaefer GB, Starr L, Pickering D, et al. Array comparative genomic hybridization findings in a cohort referred for an autism evaluation. *J Child Neurol.* Dec 2010; 25(12): 1498-503. PMID 20729506

## MEDICAL POLICY

<b>POLICY TITLE</b>	<b>GENETIC TESTING FOR DEVELOPMENTAL DELAY/INTELLECTUAL DISABILITY, AUTISM SPECTRUM DISORDER, AND CONGENITAL ANOMALIES</b>
<b>POLICY NUMBER</b>	<b>MP 2.242</b>

56. Shen Y, Dies KA, Holm IA, et al. Clinical genetic testing for patients with autism spectrum disorders. *Pediatrics*. Apr 2010; 125(4): e727-35. PMID 20231187
57. Bruno DL, Ganesamoorthy D, Schoumans J, et al. Detection of cryptic pathogenic copy number variations and constitutional loss of heterozygosity using high resolution SNP microarray analysis in 117 patients referred for cytogenetic analysis and impact on clinical practice. *J Med Genet*. Feb 2009; 46(2): 123-31. PMID 19015223
58. Friedman J, Adam S, Arbour L, et al. Detection of pathogenic copy number variants in children with idiopathic intellectual disability using 500 K SNP array genomic hybridization. *BMC Genomics*. Nov 16, 2009; 10: 526. PMID 19917086
59. Baldwin EL, Lee JY, Blake DM, et al. Enhanced detection of clinically relevant genomic imbalances using a targeted plus whole genome oligonucleotide microarray. *Genet Med*. Jun 2008; 10(6): 415-29. PMID 18496225
60. Christian SL, Brune CW, Sudi J, et al. Novel submicroscopic chromosomal abnormalities detected in autism spectrum disorder. *Biol Psychiatry*. Jun 15, 2008; 63(12): 1111-7. PMID 18374305
61. Marshall CR, Noor A, Vincent JB, et al. Structural variation of chromosomes in autism spectrum disorder. *Am J Hum Genet*. Feb 2008; 82(2): 477-88. PMID 18252227
62. Pickering DL, Eudy JD, Olney AH, et al. Array-based comparative genomic hybridization analysis of 1176 consecutive clinical genetics investigations. *Genet Med*. Apr 2008; 10(4): 262-6. PMID 18414209
63. Saam J, Gudgeon J, Aston E, et al. How physicians use array comparative genomic hybridization results to guide patient management in children with developmental delay. *Genet Med*. Mar 2008; 10(3): 181-6. PMID 18344707
64. Shevell MI, Bejjani BA, Srour M, et al. Array comparative genomic hybridization in global developmental delay. *Am J Med Genet B Neuropsychiatr Genet*. Oct 05, 2008; 147B(7): 1101-8. PMID 18361433
65. Aradhya S, Manning MA, Splendore A, et al. Whole-genome array-CGH identifies novel contiguous gene deletions and duplications associated with developmental delay, mental retardation, and dysmorphic features. *Am J Med Genet A*. Jul 01, 2007; 143A(13): 1431-41. PMID 17568414
66. Ballif BC, Sulpizio SG, Lloyd RM, et al. The clinical utility of enhanced subtelomeric coverage in array CGH. *Am J Med Genet A*. Aug 15, 2007; 143A(16): 1850-7. PMID 17632771
67. Froyen G, Van Esch H, Bauters M, et al. Detection of genomic copy number changes in patients with idiopathic mental retardation by high-resolution X-array-CGH: important role for increased gene dosage of XLMR genes. *Hum Mutat*. Oct 2007; 28(10): 1034-42. PMID 17546640
68. Hoyer J, Dreweke A, Becker C, et al. Molecular karyotyping in patients with mental retardation using 100K single-nucleotide polymorphism arrays. *J Med Genet*. Oct 2007; 44(10): 629-36. PMID 17601928
69. Lu X, Shaw CA, Patel A, et al. Clinical implementation of chromosomal microarray analysis: summary of 2513 postnatal cases. *PLoS One*. Mar 28, 2007; 2(3): e327. PMID 17389918

## MEDICAL POLICY

<b>POLICY TITLE</b>	<b>GENETIC TESTING FOR DEVELOPMENTAL DELAY/INTELLECTUAL DISABILITY, AUTISM SPECTRUM DISORDER, AND CONGENITAL ANOMALIES</b>
<b>POLICY NUMBER</b>	<b>MP 2.242</b>

70. Madrigal I, Rodriguez-Reventa L, Armengol L, et al. X-chromosome tiling path array detection of copy number variants in patients with chromosome X-linked mental retardation. *BMC Genomics*. Nov 29, 2007; 8: 443. PMID 18047645
71. Sebat J, Lakshmi B, Malhotra D, et al. Strong association of de novo copy number mutations with autism. *Science*. Apr 20, 2007; 316(5823): 445-9. PMID 17363630
72. Shen Y, Irons M, Miller DT, et al. Development of a focused oligonucleotide-array comparative genomic hybridization chip for clinical diagnosis of genomic imbalance. *Clin Chem*. Dec 2007; 53(12): 2051-9. PMID 17901113
73. Thureson AC, Bondeson ML, Edeby C, et al. Whole-genome array-CGH for detection of submicroscopic chromosomal imbalances in children with mental retardation. *Cytogenet Genome Res*. 2007; 118(1): 1-7. PMID 17901693
74. Wagenstaller J, Spranger S, Lorenz-Depiereux B, et al. Copy-number variations measured by single-nucleotide-polymorphism oligonucleotide arrays in patients with mental retardation. *Am J Hum Genet*. Oct 2007; 81(4): 768-79. PMID 17847001
75. Ballif BC, Rorem EA, Sundin K, et al. Detection of low-level mosaicism by array CGH in routine diagnostic specimens. *Am J Med Genet A*. Dec 15, 2006; 140(24): 2757- 67. PMID 17103431
76. Friedman JM, Baross A, Delaney AD, et al. Oligonucleotide microarray analysis of genomic imbalance in children with mental retardation. *Am J Hum Genet*. Sep 2006; 79(3): 500-13. PMID 16909388
77. Jacquemont ML, Sanlaville D, Redon R, et al. Array-based comparative genomic hybridisation identifies high frequency of cryptic chromosomal rearrangements in patients with syndromic autism spectrum disorders. *J Med Genet*. Nov 2006; 43(11): 843-9. PMID 16840569
78. Lugtenberg D, de Brouwer AP, Kleefstra T, et al. Chromosomal copy number changes in patients with non-syndromic X linked mental retardation detected by array CGH. *J Med Genet*. Apr 2006; 43(4): 362-70. PMID 16169931
79. Menten B, Maas N, Thienpont B, et al. Emerging patterns of cryptic chromosomal imbalance in patients with idiopathic mental retardation and multiple congenital anomalies: a new series of 140 patients and review of published reports. *J Med Genet*. Aug 2006; 43(8): 625-33. PMID 16490798
80. Miyake N, Shimokawa O, Harada N, et al. BAC array CGH reveals genomic aberrations in idiopathic mental retardation. *Am J Med Genet A*. Feb 01, 2006; 140(3): 205- 11. PMID 16419101
81. Rosenberg C, Knijnenburg J, Bakker E, et al. Array-CGH detection of micro rearrangements in mentally retarded individuals: clinical significance of imbalances present both in affected children and normal parents. *J Med Genet*. Feb 2006; 43(2): 180-6. PMID 15980116
82. Shaffer LG, Kashork CD, Saleki R, et al. Targeted genomic microarray analysis for identification of chromosome abnormalities in 1500 consecutive clinical cases. *J Pediatr*. Jul 2006; 149(1): 98-102. PMID 16860135
83. Sharp AJ, Hansen S, Selzer RR, et al. Discovery of previously unidentified genomic disorders from the duplication architecture of the human genome. *Nat Genet*. Sep 2006; 38(9): 1038-42. PMID 16906162

## MEDICAL POLICY

<b>POLICY TITLE</b>	<b>GENETIC TESTING FOR DEVELOPMENTAL DELAY/INTELLECTUAL DISABILITY, AUTISM SPECTRUM DISORDER, AND CONGENITAL ANOMALIES</b>
<b>POLICY NUMBER</b>	<b>MP 2.242</b>

84. de Vries BB, Pfundt R, Leisink M, et al. Diagnostic genome profiling in mental retardation. *Am J Hum Genet.* Oct 2005; 77(4): 606-16. PMID 16175506
85. Schoumans J, Ruivenkamp C, Holmberg E, et al. Detection of chromosomal imbalances in children with idiopathic mental retardation by array based comparative genomic hybridisation (array-CGH). *J Med Genet.* Sep 2005; 42(9): 699-705. PMID 16141005
86. Tyson C, Harvard C, Locker R, et al. Submicroscopic deletions and duplications in individuals with intellectual disability detected by array-CGH. *Am J Med Genet A.* Dec 15, 2005; 139(3): 173-85. PMID 16283669
87. Harada N, Hatchwell E, Okamoto N, et al. Subtelomere specific microarray based comparative genomic hybridisation: a rapid detection system for cryptic rearrangements in idiopathic mental retardation. *J Med Genet.* Feb 2004; 41(2): 130-6. PMID 14757861
88. Shaw-Smith C, Redon R, Rickman L, et al. Microarray based comparative genomic hybridisation (array-CGH) detects submicroscopic chromosomal deletions and duplications in patients with learning disability/mental retardation and dysmorphic features. *J Med Genet.* Apr 2004; 41(4): 241-8. PMID 15060094
89. Vissers LE, de Vries BB, Osoegawa K, et al. Array-based comparative genomic hybridization for the genomewide detection of submicroscopic chromosomal abnormalities. *Am J Hum Genet.* Dec 2003; 73(6): 1261-70. PMID 14628292
90. Chaves TF, Baretto N, Oliveira LF, et al. Copy Number Variations in a Cohort of 420 Individuals with Neurodevelopmental Disorders From the South of Brazil. *Sci Rep.* Nov 28, 2019; 9(1): 17776. PMID 31780800
91. Hu T, Zhang Z, Wang J, et al. Chromosomal Aberrations in Pediatric Patients with Developmental Delay/Intellectual Disability: A Single-Center Clinical Investigation. *Biomed Res Int.* 2019; 2019: 9352581. PMID 31781653
92. Xu M, Ji Y, Zhang T, et al. Clinical Application of Chromosome Microarray Analysis in Han Chinese Children with Neurodevelopmental Disorders. *Neurosci Bull.* Dec 2018; 34(6): 981-991. PMID 29948840
93. Sansovic I, Ivankov AM, Bobinec A, et al. Chromosomal microarray in clinical diagnosis: a study of 337 patients with congenital anomalies and developmental delays or intellectual disability. *Croat Med J.* Jun 14, 2017; 58(3): 231-238. PMID 28613040
94. Ho KS, Twede H, Vanzo R, et al. Clinical Performance of an Ultrahigh Resolution Chromosomal Microarray Optimized for Neurodevelopmental Disorders. *Biomed Res Int.* 2016; 2016: 3284534. PMID 27975050
95. Ho KS, Wassman ER, Baxter AL, et al. Chromosomal Microarray Analysis of Consecutive Individuals with Autism Spectrum Disorders Using an Ultra-High Resolution Chromosomal Microarray Optimized for Neurodevelopmental Disorders. *Int J Mol Sci.* Dec 09, 2016; 17(12). PMID 27941670
96. Hu G, Fan Y, Wang L, et al. Copy number variations in 119 Chinese children with idiopathic short stature identified by the custom genome-wide microarray. *Mol Cytogenet.* 2016; 9: 16. PMID 26884814
97. Lu XY, Phung MT, Shaw CA, et al. Genomic imbalances in neonates with birth defects: high detection rates by using chromosomal microarray analysis. *Pediatrics.* Dec 2008; 122(6): 1310-8. PMID 19047251

## MEDICAL POLICY

<b>POLICY TITLE</b>	<b>GENETIC TESTING FOR DEVELOPMENTAL DELAY/INTELLECTUAL DISABILITY, AUTISM SPECTRUM DISORDER, AND CONGENITAL ANOMALIES</b>
<b>POLICY NUMBER</b>	<b>MP 2.242</b>

98. Gogarty B. *Parents as partners. A report and guidelines on the investigation of children with developmental delay; by parents, for professionals* Cambridge: Cambridge Genetics Knowledge Park; 2006.
99. Mroch AR, Flanagan JD, Stein QP. Solving the puzzle: case examples of array comparative genomic hybridization as a tool to end the diagnostic odyssey. *Curr Probl Pediatr Adolesc Health Care*. Mar 2012; 42(3): 74-8. PMID 22325475
100. Turner G, Boyle J, Partington MW, et al. Restoring reproductive confidence in families with X-linked mental retardation by finding the causal mutation. *Clin Genet*. Feb 2008; 73(2): 188-90. PMID 18070138
101. Lingen M, Albers L, Borchers M, et al. Obtaining a genetic diagnosis in a child with disability: impact on parental quality of life. *Clin Genet*. Feb 2016; 89(2): 258-66. PMID 26084449
102. Hayeems RZ, Hoang N, Chenier S, et al. Capturing the clinical utility of genomic testing: medical recommendations following pediatric microarray. *Eur J Hum Genet*. Sep 2015; 23(9): 1135-41. PMID 25491637
103. Ellison JW, Ravnán JB, Rosenfeld JA, et al. Clinical utility of chromosomal microarray analysis. *Pediatrics*. Nov 2012; 130(5): e1085-95. PMID 23071206
104. Hoffmann TJ, Windham GC, Anderson M, et al. Evidence of reproductive stoppage in families with autism spectrum disorder: a large, population-based cohort study. *JAMA Psychiatry*. Aug 2014; 71(8): 943-51. PMID 24942798
105. Grozeva D, Carss K, Spasic-Boskovic O, et al. Targeted Next-Generation Sequencing Analysis of 1,000 Individuals with Intellectual Disability. *Hum Mutat*. Dec 2015; 36(12): 1197-204. PMID 26350204
106. Kalsner L, Twachtman-Bassett J, Tokarski K, et al. Genetic testing including targeted gene panel in a diverse clinical population of children with autism spectrum disorder: Findings and implications. *Mol Genet Genomic Med*. Mar 2018; 6(2): 171-185. PMID 29271092
107. Volkmar F, Siegel M, Woodbury-Smith M, et al. Practice parameter for the assessment and treatment of children and adolescents with autism spectrum disorder. *J Am Acad Child Adolesc Psychiatry*. Feb 2014; 53(2): 237-57. PMID 24472258
108. Michelson DJ, Shevell MI, Sherr EH, et al. Evidence report: Genetic and metabolic testing on children with global developmental delay: report of the Quality Standards Subcommittee of the American Academy of Neurology and the Practice Committee of the Child Neurology Society. *Neurology*. Oct 25, 2011; 77(17): 1629-35. PMID 21956720
109. Manning M, Hudgins L. Array-based technology and recommendations for utilization in medical genetics practice for detection of chromosomal abnormalities. *Genet Med*. Nov 2010; 12(11): 742-5. PMID 20962661
110. Manning M, Hudgins L. Addendum: Array-based technology and recommendations for utilization in medical genetics practice for detection of chromosomal abnormalities. *Genet Med*. Jun 08, 2020. PMID 32514088
111. South ST, Lee C, Lamb AN, et al. ACMG Standards and Guidelines for constitutional cytogenomic microarray analysis, including postnatal and prenatal applications: revision 2013. *Genet Med*. Nov 2013; 15(11): 901-9. PMID 24071793

## MEDICAL POLICY

<b>POLICY TITLE</b>	<b>GENETIC TESTING FOR DEVELOPMENTAL DELAY/INTELLECTUAL DISABILITY, AUTISM SPECTRUM DISORDER, AND CONGENITAL ANOMALIES</b>
<b>POLICY NUMBER</b>	<b>MP 2.242</b>

112. Schaefer GB, Mendelsohn NJ. *Clinical genetics evaluation in identifying the etiology of autism spectrum disorders: 2013 guideline revisions. Genet Med. May 2013; 15(5): 399-407. PMID 23519317*
113. Manickam K, McClain MR, Demmer LA, et al. *Exome and genome sequencing for pediatric patients with congenital anomalies or intellectual disability: an evidence-based clinical guideline of the American College of Medical Genetics and Genomics (ACMG). Genet Med. 2021;23(11):2029-2037. doi:10.1038/s41436-021-01242-6*
114. Vasudevan P, Suri M. *A clinical approach to developmental delay and intellectual disability. Clin Med (Lond). 2017;17(6):558-561. doi:10.7861/clinmedicine.17-6-558*
115. Savatt JM, Myers SM. *Genetic Testing in Neurodevelopmental Disorders. Front Pediatr. 2021;9:526779. Published 2021 Feb 19. doi:10.3389/fped.2021.526779*
116. Ewans LJ, Schofield D, Shrestha R, et al. *Whole-exome sequencing reanalysis at 12 months boosts diagnosis and is cost-effective when applied early in Mendelian disorders. Genet Med. 2018;20(12):1564-1574. doi:10.1038/gim.2018.39 PMID 29595814*
117. Hulick, Peter J MD, *Next-generation sequencing (NGS): Principles and clinical applications. In: Up To Date Online. Waltham, MA: Last updated May 02, 2025. Literature review through June 2025*
118. Pivalizza, P MD and Lalani, S MD, *Intellectual disability (ID) in children: Clinical features, evaluation, and diagnosis. In: UpToDate Online. Waltham, MA: Last updated Feb. 14, 2024. Literature review June 2025.*
119. Sanchack, K; *Autism Spectrum Disorder: Updated Guidelines from the American Academy of Pediatrics. American Family Physician, 2020;102(9):629-631*
120. Aldosari AN, Aldosari TS. *Comprehensive evaluation of the child with global developmental delays or intellectual disability. Clin Exp Pediatr. 2024;67(9):435-446. doi:10.3345/cep.2023.01697. PMID: 38810986*

### IX. POLICY HISTORY

[TOP](#)

<b>MP 2.242</b>	<b>01/01/2020 Administrative Update.</b> New code 0156U added.
	<b>02/26/2020 Consensus Review.</b> No changes to policy statement. Coding reviewed. New code 0170U added as investigational, effective 04/01/2020.
	<b>10/01/2020 Administrative Update.</b> New code 0209U added, effective 10/01/2020.
	<b>09/07/2021 Administrative Update.</b> New codes F78.A1 and F78.A0 added. Effective 10/01/2021
	<b>10/04/2021 Consensus Review.</b> No change to policy statement. Description/background and references updated. Coding section revision: 0156U moved from medically necessary to investigational.
	<b>03/11/2022 Administrative Update.</b> New code 0318U added as conditionally covered; effective 04/01/2022.
	<b>08/30/2022 Minor Review.</b> Exome and genome sequencing now MN. Literature review and update. Societal recommendations added to policy

## MEDICAL POLICY

<b>POLICY TITLE</b>	<b>GENETIC TESTING FOR DEVELOPMENTAL DELAY/INTELLECTUAL DISABILITY, AUTISM SPECTRUM DISORDER, AND CONGENITAL ANOMALIES</b>
<b>POLICY NUMBER</b>	<b>MP 2.242</b>

	guidelines. Supporting information added to background. Added 2.324 to cross ref. Ref update. 81349 added, 0209U moved from INV to MN.
	<b>07/26/2023 Minor Review.</b> Added criteria for standard whole exome and genome sequencing to including genetic counseling and clarify appropriate timing of testing. Updates to background, references. CPT codes added to include WES/WGS codes: 0212U, 0214U, 0265U, 0267U, 81415, 81417, 81425, 81427.
	<b>01/19/2024 Administrative Update.</b> Clinical benefit added.
	<b>10/07/2024 Consensus Review.</b> No change to policy stance. Updated references.
	<b>07/07/2025 Minor review.</b> Removed criteria regarding whole exome and whole genome sequencing as this is addressed in <b>MP 2.324</b> . Codes 0267U, 0212U, 0214U, 0265U, 81415, 81417, 81425, and 81427 removed from policy; remain on <b>MP 2.324</b> .
	<b>09/02/2025 Administrative Update.</b> New ICD10 codes added effective 10/1/25
	<b>10/09/2025 Administrative Update.</b> Removed Benefit Variations Section and updated Disclaimer.
	<b>03/05/2026 Retirement Review.</b> EviCore delegation.

[Top](#)

*Health care benefit programs issued or administered by Capital Blue Cross and/or its subsidiaries, Capital Advantage Insurance Company®, Capital Advantage Assurance Company® and Keystone Health Plan® Central. Independent licensees of the Blue Cross BlueShield Association. Communications issued by Capital Blue Cross in its capacity as administrator of programs and provider relations for all companies.*