

MEDICAL POLICY

POLICY TITLE	GENETIC TESTING OF CADASIL SYNDROME
POLICY NUMBER	MP 2.247

CLINICAL BENEFIT	☐ MINIMIZE SAFETY RISK OR CONCERN. ☐ MINIMIZE HARMFUL OR INEFFECTIVE INTERVENTIONS. ☐ ASSURE APPROPRIATE LEVEL OF CARE. ☐ ASSURE APPROPRIATE DURATION OF SERVICE FOR INTERVENTIONS. ☒ ASSURE THAT RECOMMENDED MEDICAL PREREQUISITES HAVE BEEN MET. ☐ ASSURE APPROPRIATE SITE OF TREATMENT OR SERVICE.
Effective Date:	RETIRED 7/1/2026

POLICY

Genetic testing of *NOTCH3* to confirm the diagnosis of cerebral autosomal dominant arteriopathy with subcortical infarcts and leukoencephalopathy (CADASIL) syndrome may be considered **medically necessary** under the following conditions:

- Clinical signs, symptoms, and imaging results are consistent with CADASIL, indicating that the pre-test probability of CADASIL is at least in the moderate to high range (see Policy Guidelines); **and**
- The diagnosis of CADASIL is inconclusive following alternate methods of testing, including magnetic resonance imaging.

For individuals who are asymptomatic with a family member with a diagnosis of CADASIL syndrome:

- If there is a family member (first- and second-degree relative) with a known variant, targeted genetic testing of the known *NOTCH3* familial variant may be considered **medically necessary**
- If the family member's genetic status is unknown, genetic testing of *NOTCH3* (see Policy Guidelines) may be considered **medically necessary**.

Genetic testing of *NOTCH3* to confirm the diagnosis of CADASIL syndrome in all other situations is considered **investigational**. There is insufficient evidence to support a general conclusion concerning the health outcomes or benefits associated with this listed procedure.

Policy Guidelines

Genetic testing for *NOTCH3* comprises targeted sequencing of specific exons (e.g., exon 4 only, exons 2-6), general sequencing of *NOTCH3* exons (e.g., exons 2-24 or all 33 exons), or targeted testing for known *NOTCH3* pathogenic variants. Skin biopsy should be reserved for patients where *NOTCH3* genetic testing is inconclusive (e.g. variants of uncertain significance).

The probability that CADASIL is present in an individualized assessment depends on numerous factors such as family history, symptoms, imaging results, and other specialized testing such as skin biopsy.

MEDICAL POLICY

POLICY TITLE	GENETIC TESTING OF CADASIL SYNDROME
POLICY NUMBER	MP 2.247

Pescini et al (2012) attempted to identify clinical factors that increase the likelihood of a pathogenic variant being present. Table PG1 summarizes the pooled frequency of clinical and radiologic features, and the points assigned for each finding. The authors recommended that a total score of 14 be used to select patients for testing, because this score resulted in a high sensitivity (96.7%) and a moderately high specificity (74.2%).

Table PG1. Pooled Frequency of Clinical and Radiologic Features

Features	No. With <i>NOTCH3</i> Variant	Percent With <i>NOTCH3</i> Variant	Points
Clinical			
Migraine	239/463	52%	1
Migraine with aura	65/85	76%	3
Transient ischemic attack/stroke	380/526	72%	1 (2 if <50 y)
Psychiatric disturbance	106/380	28%	1
Cognitive decline	188/434	43%	3
Radiologic			
LE	277Pescini/277	100%	3
LE extended to temporal pole	174/235	74%	1
LE extended to external capsule	228/303	75%	5
Subcortical infarcts	210/254	83%	2

Genetics Nomenclature Update

Human Genome Variation Society nomenclature is used to report information on variants found in DNA and serves as an international standard in DNA diagnostics. It was implemented for genetic testing medical evidence review updates 2017 (see Table PG2). The Society's nomenclature is recommended by the Human Variome Project, the Human Genome Organization, and by the Human Genome Variation Society itself.

The American College of Medical Genetics and Genomics and Association for Molecular Pathology standards and guidelines for interpretation of sequence variants represent expert opinion from both organizations, in addition to the College of American Pathologists. These recommendations primarily apply to genetic tests used in clinical laboratories, including genotyping, single genes, panels, exomes, and genomes. Table PG3 shows the recommended standard terminology—"pathogenic," "likely pathogenic," "uncertain significance," "likely benign," and "benign"—to describe variants identified that cause Mendelian disorders.

Table PG2. Nomenclature to Report on Variants Found in DNA

Previous	Updated	Definition
Mutation	Disease-associated variant	Disease-associated change in the DNA sequence

MEDICAL POLICY

POLICY TITLE	GENETIC TESTING OF CADASIL SYNDROME
POLICY NUMBER	MP 2.247

	Variant	Change in the DNA sequence
	Familial variant	Disease-associated variant identified in a proband for use in subsequent targeted genetic testing in first-degree relatives

Table PG3. ACMG-AMP Standards and Guidelines for Variant Classification

Variant Classification	Definition
Pathogenic	Disease-causing change in the DNA sequence
Likely pathogenic	Likely disease-causing change in the DNA sequence
Variant of uncertain significance	Change in DNA sequence with uncertain effects on disease
Likely benign	Likely benign change in the DNA sequence
Benign	Benign change in the DNA sequence

ACMG: American College of Medical Genetics and Genomics; AMP: Association for Molecular Pathology.

Genetic Counseling

Experts recommend formal genetic counseling for individuals who are at risk for inherited disorders and who wish to undergo genetic testing. Interpreting the results of genetic tests and understanding risk factors can be difficult for some individuals; genetic counseling helps individuals understand the impact of genetic testing, including the possible effects the test results could have on the individual or their family members. It should be noted that genetic counseling may alter the utilization of genetic testing substantially and may reduce inappropriate testing; further, genetic counseling should be performed by an individual with experience and expertise in genetic medicine and genetic testing methods.

Cross-Reference:

MP 7.009 Preimplantation Genetic Testing

PRODUCT VARIATIONS

This policy is only applicable to certain programs and products administered by Capital BlueCross and subject to benefit variations. Please see additional information below.

FEP PPO: Refer to FEP Medical Policy Manual. The FEP Medical Policy manual can be found at: <https://www.fepblue.org/benefit-plans/medical-policies-and-utilization-management-guidelines/medical-policies>.

DESCRIPTION/BACKGROUND

Variants in the *NOTCH3* gene have been causally associated with cerebral autosomal dominant arteriopathy with subcortical infarcts and leukoencephalopathy (CADASIL). Genetic testing is available to determine if pathogenic variants exist in the *NOTCH3* gene for individuals with suspected CADASIL and their family members.

MEDICAL POLICY

POLICY TITLE	GENETIC TESTING OF CADASIL SYNDROME
POLICY NUMBER	MP 2.247

CADASIL

Cerebral autosomal dominant arteriopathy with subcortical infarcts and leukoencephalopathy (CADASIL) is an uncommon, autosomal dominant disease, though it is the most common cause of hereditary stroke and hereditary vascular dementia in adults. CADASIL syndrome is an adult-onset, disabling systemic condition, characterized by a migraine with aura, recurrent lacunar strokes, progressive cognitive impairment, and psychiatric disorders. The overall prevalence of the disease is unknown in the general population.

Diagnosis

The differential diagnosis of CADASIL includes the following conditions (see Table 1).

Table 1. Differential Diagnosis of CADASIL

Acquired Disorders	Inherited Disorders
- Sporadic SVD with or without hypertension as the main risk factor - Multiple sclerosis - Primary angiitis of the central nervous system	- Fabry disease - Cerebral autosomal recessive arteriopathy with subcortical infarcts and leukoencephalopathy - Familial SVD caused by heterozygous variants in the <i>HTRA1</i> gene - Some forms of leukodystrophy

SVD: small vessel disease.

Since the clinical presentation of CADASIL varies, the condition may be confused with multiple sclerosis, Alzheimer dementia, and Binswanger disease. The specific clinical signs and symptoms, along with family history and brain magnetic resonance imaging findings, are extremely important in diagnosing CADASIL. The clinical features and mode of inheritance (autosomal dominant vs autosomal recessive) help to distinguish CADASIL from other inherited disorders in a differential diagnosis.

When the differential diagnosis includes CADASIL, various diagnostic tests are available:

- Genetic testing, by direct sequencing of select exons or of exons 2 through 24 of the *NOTCH3* gene (see the Rationale section). Identification of a *NOTCH3* pathogenic variant definitively establishes a diagnosis of CADASIL without the need for additional diagnostic testing (e.g., skin biopsy)
- Genetic testing can facilitate the differentiation of *NOTCH3*-associated cerebral small vessel disease (CSVD) from other CADASIL-like disorders (e.g., *HTRA1*-associated CSVD and cerebral autosomal recessive arteriopathy with subcortical infarcts and leukoencephalopathy).
- Immunohistochemistry assay of a skin biopsy sample, using a monoclonal antibody with reactivity against the extracellular domain of the *NOTCH3* receptor. Positive immunostaining reveals the accumulation of the *NOTCH3* protein in the walls of small blood vessels. Lesnick Oberstein et al (2003) estimated the sensitivity and specificity at 85% to 90% and 95% to 100%, respectively, for 2 observers of the test results in a

MEDICAL POLICY

POLICY TITLE	GENETIC TESTING OF CADASIL SYNDROME
POLICY NUMBER	MP 2.247

population of patients and controls correlated with clinical, genetic, and magnetic resonance imaging parameters

- Detection of granular osmiophilic material in the same skin biopsy sample by electron microscopy. The major component of granular osmiophilic material is the ectodomain of the *NOTCH3* gene product. Granular osmiophilic material accumulates directly in vascular smooth muscle cells and, when present, is considered a hallmark of the disease. However, granular osmiophilic material may not be present in all biopsy samples. Sensitivity has been reported as low as 45% and 57% but specificity is generally near or at 100%.
- Examination of brain tissue for the presence of granular osmiophilic material was originally described as limited to brain blood vessels. Examination of brain biopsy or autopsy after death was an early criterion standard for diagnosis. In some cases, peripheral staining for granular osmiophilic material has been absent even though positive results were seen in brain blood vessels.

NOTCH3 Variants

Variants in *NOTCH3* have been identified as the underlying cause of CADASIL. In almost all cases, the pathogenic variants lead to loss or gain of a cysteine residue that can lead to increased reactivity of the *NOTCH3* protein, resulting in ligand-binding and toxic effects.

The *NOTCH3* gene is found on chromosome 19p13.12 and encodes the third discovered human homologue of the *Drosophila melanogaster* type I membrane protein *NOTCH*. The *NOTCH3* protein consists of 2321 amino acids, primarily expressed in vascular smooth muscle cells, and plays an important role in the control of vascular transduction. It has an extracellular ligand-binding domain of 34 epidermal growth factor (EGF)-like repeats, traverses the membrane once, and has an intracellular domain required for signal transduction.

Variants in the *NOTCH3* gene have been differentiated into those causative of the CADASIL syndrome (pathogenic variants) and those of uncertain significance. Pathogenic variants affect conserved cysteine residues within 34 EGF-like repeat domains in the extracellular portion of the *NOTCH3* protein. More than 150 pathogenic variants have been reported in at least 500 pedigrees. *NOTCH3* has 33 exons but all CADASIL variants reported to date have occurred in exons 2 to 24, which encode the 34 EGF-like repeats, with strong clustering in exons 3 and 4, which encode EGF receptors 2 to 5 (>40% of variants in >70% of families occur in these exons). A study by Hack et al (2023) identified 3 clinically distinct risk categories (high, medium, and low) of *NOTCH3* EGF-like repeats using data from CADASIL and population cohorts (N=4221). Some studies have indicated that the clinical variability in CADASIL presentation, particularly about the development of white matter hyperintensities on magnetic resonance imaging, may be related to genetic modifiers outside the *NOTCH3* locus but the specific role of these modifiers is not well-delineated. Dupé et al (2023) investigated the phenotypic variability in individuals with CADASIL and found the mutation location in the *NOTCH3* gene (n=436) to be strongly related to clinical severity, and found male sex, arterial hypertension, and smoking to be associated with increased disease severity.

MEDICAL POLICY

POLICY TITLE	GENETIC TESTING OF CADASIL SYNDROME
POLICY NUMBER	MP 2.247

The probability that CADASIL is present in an individualized assessment depends on numerous factors such as family history; symptoms, imaging results, and other specialized testing (e.g., skin biopsy). Pescini et al (2012) attempted to identify clinical factors that increase the likelihood of a pathogenic variant being present, with increasing likelihood with the presence of one or several factors, including a migraine, migraine with aura, transient ischemic attack/stroke, psychiatric disturbance, cognitive decline, leukoencephalopathy (with greater risk for leukoencephalopathy extending to the temporal pole or external capsule), and subcortical infarcts.

Regulatory Status

Clinical laboratories may develop and validate tests in-house and market them as a laboratory service; laboratory-developed tests must meet the general regulatory standards of the Clinical Laboratory Improvement Amendments (CLIA). Genetic testing of *NOTCH3* is available under the auspices of the CLIA. Laboratories that offer laboratory-developed tests must be licensed by the CLIA for high-complexity testing. To date, the U.S. Food and Drug Administration has chosen not to require any regulatory review of this test.

RATIONALE

Summary of Evidence

For individuals with suspected CADASIL syndrome who receive *NOTCH3* genetic testing, the evidence includes case reports, case series, and genotype-phenotype correlation studies evaluating the clinical validity and genetic testing yield for *NOTCH3*. The relevant outcomes are overall survival, test accuracy and validity, changes in reproductive decision making, change in disease status, and morbid events. The clinical validity studies have demonstrated that a *NOTCH3* pathogenic variant is found in a high percentage of individuals with a clinical diagnosis of CADASIL, with studies reporting a clinical sensitivity of 90% to 100%. Limited data on specificity derives from testing small numbers of healthy controls, and no false-positive *NOTCH3* pathogenic variants have been reported in these populations. The diagnostic yield studies have reported a variable yield, ranging from 10% to 54%. These lower numbers likely reflect testing in heterogeneous populations that include individuals with other disorders. No direct evidence was identified demonstrating outcome improvements associated with genetic testing for CADASIL. However, a chain of evidence can be constructed to demonstrate that identification of a *NOTCH3* pathogenic variant establishes the diagnosis of CADASIL without the need for a skin biopsy and reduces the need for other diagnostic tests used to exclude other conditions in a differential diagnosis. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who are asymptomatic with family members who have CADASIL syndrome who receive targeted genetic testing for a known *NOTCH3* familial variant, the evidence is limited. The relevant outcomes are overall survival, test accuracy and validity, changes in reproductive decision making, change in disease status, and morbid events. For asymptomatic family members of an individual with known CADASIL, knowledge of the presence of a familial variant may lead to changes in lifestyle decisions for the affected individual (e.g., reproduction,

MEDICAL POLICY

POLICY TITLE	GENETIC TESTING OF CADASIL SYNDROME
POLICY NUMBER	MP 2.247

employment). However, the impact of these lifestyle decisions on health outcomes is uncertain, and there are no interventions for asymptomatic individuals that are known to delay or prevent disease onset. A chain of evidence can be constructed to demonstrate that identification of a *NOTCH3* familial variant predicts future development of CADASIL in an asymptomatic individual, eliminates the need for additional diagnostic testing, allows for earlier monitoring for development of systems, aids in reproductive planning, and helps determine the likelihood of an affected offspring. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who are asymptomatic with family members who have CADASIL syndrome whose genetic status is unknown who receive *NOTCH3* genetic testing, the evidence is limited. The relevant outcomes are overall survival, test accuracy and validity, changes in reproductive decision making, change in disease status, and morbid events. For asymptomatic family members of an individual with known CADASIL whose genetic status is unknown, knowledge of the presence of a *NOTCH3* pathogenic variant may lead to changes in lifestyle decisions for the affected individual (e.g., reproduction, employment). However, the impact of these lifestyle decisions on health outcomes is uncertain, and there are no interventions for asymptomatic individuals that are known to delay or prevent disease onset. A chain of evidence can be constructed to demonstrate that identification of a *NOTCH3* pathogenic variant predicts future development of CADASIL in an asymptomatic individual, eliminates the need for additional diagnostic testing, allows for earlier monitoring for development of systems, aids in reproductive planning, and helps determine the likelihood of an affected offspring. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

DEFINITIONS

AUTOSOMAL DOMINANT is a gene on one of the non-sex chromosomes that is always expressed, even if only one copy is present. The chance of passing the gene to offspring is 50% for each pregnancy.

GENOTYPE is the specific genetic makeup of an individual, usually in the form of DNA.

LEUKOENCEPHALOPATHY refers to any of a group of diseases affecting the white substance of the brain.

MUTATION is a permanent structural alteration in DNA.

DISCLAIMER

Capital Blue Cross' medical policies are used to determine coverage for specific medical technologies, procedures, equipment, and services. These medical policies do not constitute medical advice and are subject to change as required by law or applicable clinical evidence from independent treatment guidelines. Treating providers are solely responsible for medical advice and treatment of members. These policies are not a guarantee of coverage or

MEDICAL POLICY

POLICY TITLE	GENETIC TESTING OF CADASIL SYNDROME
POLICY NUMBER	MP 2.247

payment. Payment of claims is subject to a determination regarding the member's benefit program and eligibility on the date of service, and a determination that the services are medically necessary and appropriate. Final processing of a claim is based upon the terms of contract that applies to the members' benefit program, including benefit limitations and exclusions. If a provider or a member has a question concerning this medical policy, please contact Capital Blue Cross' Provider Services or Member Services.

CODING INFORMATION

Note: This list of codes may not be all-inclusive, and codes are subject to change at any time. The identification of a code in this section does not denote coverage as coverage is determined by the terms of member benefit information. In addition, not all covered services are eligible for separate reimbursement.

Covered when Medically Necessary:

Procedure Codes								
81406								

ICD-10-CM Diagnosis Codes	Description
A81.2	Progressive multifocal leukoencephalopathy
E75.25	Metachromatic leukodystrophy
E75.27	Pelizaeus-Merzbacher disease
F01.50	Vascular dementia without behavioral disturbance
F01.51	Vascular dementia with behavioral disturbance
F06.31	Mood disorder due to known physiological condition with depressive features
F06.32	Mood disorder due to known physiological condition with major depressive-like epis
F06.4	Anxiety disorder due to known physiological condition
F06.8	Other specified mental disorders due to known physiological condition
G43.001	Migraine without aura, not intractable, with status migrainosus
G43.009	Migraine without aura, not intractable, without status migrainosus
G43.011	Migraine without aura, intractable, with status migrainosus
G43.019	Migraine without aura, intractable, without status migrainosus
G43.101	Migraine with aura, not intractable, with status migrainosus
G43.109	Migraine with aura, not intractable, without status migrainosus
G43.111	Migraine with aura, intractable, with status migrainosus
G43.119	Migraine with aura, intractable, without status migrainosus
G43.401	Hemiplegic migraine, not intractable, with status migrainosus
G43.409	Hemiplegic migraine, not intractable, without status migrainosus
G43.411	Hemiplegic migraine, intractable, with status migrainosus
G43.419	Hemiplegic migraine, intractable, without status migrainosus

MEDICAL POLICY

POLICY TITLE	GENETIC TESTING OF CADASIL SYNDROME
POLICY NUMBER	MP 2.247

ICD-10-CM Diagnosis Codes	Description
G43.501	Persistent migraine aura without cerebral infarction, not intractable, with status migrainosus
G43.509	Persistent migraine aura without cerebral infarction, not intractable, without status migrainosus
G43.511	Persistent migraine aura without cerebral infarction, intractable, with status migrainosus
G43.519	Persistent migraine aura without cerebral infarction, intractable, without status migrainosus
G43.601	Persistent migraine aura with cerebral infarction, not intractable, with status migrainosus
G43.609	Persistent migraine aura with cerebral infarction, not intractable, without status migrainosus
G43.611	Persistent migraine aura with cerebral infarction, intractable, with status migrainosus
G43.619	Persistent migraine aura with cerebral infarction, intractable, without status migrainosus
G43.801	Other migraine, not intractable, with status migrainosus
G43.809	Other migraine, not intractable, without status migrainosus
G43.811	Other migraine, intractable, with status migrainosus
G43.819	Other migraine, intractable, without status migrainosus
G43.E01	Chronic migraine with aura, not intractable, with status migrainosus
G43.E09	Chronic migraine with aura, not intractable, without status migrainosus
G43.E11	Chronic migraine with aura, intractable, with status migrainosus
G43.E19	Chronic migraine with aura, intractable, without status migrainosus
G45.8	Other transient cerebral ischemic attacks and related syndromes
G45.9	Transient cerebral ischemic attack, unspecified
I63.311	Cerebral infarction due to thrombosis of right middle cerebral artery
I63.312	Cerebral infarction due to thrombosis of left middle cerebral artery
I63.313	Cerebral infarction due to thrombosis of bilateral middle cerebral arteries
I63.321	Cerebral infarction due to thrombosis of right anterior cerebral artery
I63.322	Cerebral infarction due to thrombosis of left anterior cerebral artery
I63.323	Cerebral infarction due to thrombosis of bilateral anterior arteries
I63.331	Cerebral infarction due to thrombosis of right posterior cerebral artery
I63.332	Cerebral infarction due to thrombosis of left posterior cerebral artery
I63.333	Cerebral infarction due to thrombosis of bilateral posterior cerebral arteries
I63.341	Cerebral infarction due to thrombosis of right cerebellar artery
I63.342	Cerebral infarction due to thrombosis of left cerebellar artery
I63.343	Cerebral infarction due to thrombosis of bilateral cerebellar arteries
I63.39	Cerebral infarction due to thrombosis of other cerebral artery

MEDICAL POLICY

POLICY TITLE	GENETIC TESTING OF CADASIL SYNDROME
POLICY NUMBER	MP 2.247

ICD-10-CM Diagnosis Codes	Description
I63.411	Cerebral infarction due to embolism of right middle cerebral artery
I63.412	Cerebral infarction due to embolism of left middle cerebral artery
I63.413	Cerebral infarction due to embolism of bilateral middle cerebral arteries
I63.421	Cerebral infarction due to embolism of right anterior cerebral artery
I63.422	Cerebral infarction due to embolism of left anterior cerebral artery
I63.423	Cerebral infarction due to embolism of bilateral anterior cerebral arteries
I63.431	Cerebral infarction due to embolism of right posterior cerebral artery
I63.432	Cerebral infarction due to embolism of left posterior cerebral artery
I63.433	Cerebral infarction due to embolism of bilateral posterior cerebral arteries
I63.441	Cerebral infarction due to embolism of right cerebellar artery
I63.442	Cerebral infarction due to embolism of left cerebellar artery
I63.443	Cerebral infarction due to embolism of bilateral cerebellar arteries
I63.49	Cerebral infarction due to embolism of other cerebral artery
I63.511	Cerebral infarction due to unspecified occlusion or stenosis of right middle cerebral artery
I63.512	Cerebral infarction due to unspecified occlusion or stenosis of left middle cerebral artery
I63.513	Cerebral infarction due to unspecified occlusion or stenosis of bilateral middle arteries
I63.521	Cerebral infarction due to unspecified occlusion or stenosis of right anterior cerebral artery
I63.522	Cerebral infarction due to unspecified occlusion or stenosis of left anterior cerebral artery
I63.523	Cerebral infarction due to unspecified occlusion or stenosis of bilateral anterior arteries
I63.531	Cerebral infarction due to unspecified occlusion or stenosis of right posterior cerebral artery
I63.532	Cerebral infarction due to unspecified occlusion or stenosis of left posterior cerebral artery
I63.533	Cerebral infarction due to unspecified occlusion or stenosis of bilateral posterior arteries
I63.541	Cerebral infarction due to unspecified occlusion or stenosis of right cerebellar artery
I63.542	Cerebral infarction due to unspecified occlusion or stenosis of left cerebellar artery
I63.543	Cerebral infarction due to unspecified occlusion or stenosis of bilateral cerebellar arteries
I63.59	Cerebral infarction due to unspecified occlusion or stenosis of other cerebral artery
I67.3	Progressive vascular leukoencephalopathy
I63.81	Other cerebral infarction due to occlusion or stenosis of small artery
I63.89	Other cerebral infarction
I67.850	Cerebral autosomal dominant arteriopathy with subcortical infarcts and leukoencephalopathy

MEDICAL POLICY

POLICY TITLE	GENETIC TESTING OF CADASIL SYNDROME
POLICY NUMBER	MP 2.247

ICD-10-CM Diagnosis Codes	Description
R41.81	Age-related cognitive decline
R41.82	Altered mental status, unspecified
Z82.49	Family history of ischemic heart disease and other diseases of the circulatory system

REFERENCES

1. Yuan L, Chen X, Jankovic J, et al. CADASIL: A NOTCH3-associated cerebral small vessel disease. *J Adv Res.* Dec 2024; 66: 223-235. PMID 38176524
2. Joutel A, Favrole P, Labauge P, et al. Skin biopsy immunostaining with a Notch3 monoclonal antibody for CADASIL diagnosis. *Lancet.* Dec 15 2001; 358(9298): 2049-51. PMID 11755616
3. Lesnik Oberstein SA, van Duinen SG, van den Boom R, et al. Evaluation of diagnostic NOTCH3 immunostaining in CADASIL. *Acta Neuropathol.* Aug 2003; 106(2): 107-11. PMID 12756589
4. Muqtadar H, Testai FD. Single gene disorders associated with stroke: a review and update on treatment options. *Curr Treat Options Cardiovasc Med.* Jun 2012; 14(3): 288-97. PMID 22528196
5. del Río-Espínola A, Mendióroz M, Domingues-Montanari S, et al. CADASIL management or what to do when there is little one can do. *Expert Rev Neurother.* Feb 2009; 9(2): 197-210. PMID 19210195
6. Malandrini A, Gaudio C, Gambelli S, et al. Diagnostic value of ultrastructural skin biopsy studies in CADASIL. *Neurology.* Apr 24 2007; 68(17): 1430-2. PMID 17452591
7. Brulin P, Godfraind C, Leteurtre E, et al. Morphometric analysis of ultrastructural vascular changes in CADASIL: analysis of 50 skin biopsy specimens and pathogenic implications. *Acta Neuropathol.* Sep 2002; 104(3): 241-8. PMID 12172909
8. Markus HS, Martin RJ, Simpson MA, et al. Diagnostic strategies in CADASIL. *Neurology.* Oct 22 2002; 59(8): 1134-8. PMID 12395806
9. Choi JC, Lee KH, Song SK, et al. Screening for NOTCH3 gene mutations among 151 consecutive Korean patients with acute ischemic stroke. *J Stroke Cerebrovasc Dis.* Jul 2013; 22(5): 608-14. PMID 22133740
10. Mosca L, Marazzi R, Ciccone A, et al. NOTCH3 gene mutations in subjects clinically suspected of CADASIL. *J Neurol Sci.* Aug 15 2011; 307(1-2): 144-8. PMID 21616505
11. Hack RJ, Rutten J, Lesnik Oberstein SAJ. In: Adam MP, Feldman J, Mirzaa GM, et al., eds. *GeneReviews*, Seattle, WA: University of Washington; 2019.
12. Donahue CP, Kosik KS. Distribution pattern of Notch3 mutations suggests a gain-of-function mechanism for CADASIL. *Genomics.* Jan 2004; 83(1): 59-65. PMID 14667809
13. Chabriat H, Joutel A, Dichgans M, et al. Cadasil. *Lancet Neurol.* Jul 2009; 8(7): 643-53. PMID 19539236
14. Hack RJ, Gravesteijn G, Cerfontaine MN, et al. Three-tiered EGFr domain risk stratification for individualized NOTCH3-small vessel disease prediction. *Brain.* Jul 03 2023; 146(7): 2913-2927. PMID 36535904

MEDICAL POLICY

POLICY TITLE	GENETIC TESTING OF CADASIL SYNDROME
POLICY NUMBER	MP 2.247

15. Opherck C, Gonik M, Duering M, et al. Genome-wide genotyping demonstrates a polygenic risk score associated with white matter hyperintensity volume in CADASIL. *Stroke*. Apr 2014; 45(4): 968-72. PMID 24578207
16. Dupé C, Guey S, Biard L, et al. Phenotypic variability in 446 CADASIL patients: Impact of NOTCH3 gene mutation location in addition to the effects of age, sex and vascular risk factors. *J Cereb Blood Flow Metab*. Jan 2023; 43(1): 153-166. PMID 36254369
17. Pescini F, Nannucci S, Bertaccini B, et al. The Cerebral Autosomal-Dominant Arteriopathy With Subcortical Infarcts and Leukoencephalopathy (CADASIL) Scale: a screening tool to select patients for NOTCH3 gene analysis. *Stroke*. Nov 2012; 43(11): 2871-6. PMID 22996955
18. Lee YC, Liu CS, Chang MH, et al. Population-specific spectrum of NOTCH3 mutations, MRI features and founder effect of CADASIL in Chinese. *J Neurol*. Feb 2009; 256(2): 249-55. PMID 19242647
19. Yin X, Wu D, Wan J, et al. Cerebral autosomal dominant arteriopathy with subcortical infarcts and leukoencephalopathy: Phenotypic and mutational spectrum in patients from mainland China. *Int J Neurosci*. 2015; 125(8): 585-92. PMID 25105908
20. Joutel A, Vahedi K, Corpechot C, et al. Strong clustering and stereotyped nature of Notch3 mutations in CADASIL patients. *Lancet*. Nov 22 1997; 350(9090): 1511-5. PMID 9388399
21. Abramychewa N, Stepanova M, Kalashnikova L, et al. New mutations in the Notch3 gene in patients with cerebral autosomal dominant arteriopathy with subcortical infarcts and leukoencephalopathy (CADASIL). *J Neurol Sci*. Feb 15 2015; 349(1-2): 196-201. PMID 25623805
22. Maksemous N, Smith RA, Haupt LM, et al. Targeted next generation sequencing identifies novel NOTCH3 gene mutations in CADASIL diagnostics patients. *Hum Genomics*. Nov 24 2016; 10(1): 38. PMID 27881154
23. Gorukmez O, Gorukmez O, Topak A, et al. NOTCH3 Variants in Patients with Suspected CADASIL. *Ann Indian Acad Neurol*. 2023; 26(4): 484-490. PMID 37970308
24. Peters N, Opherck C, Bergmann T, et al. Spectrum of mutations in biopsy-proven CADASIL: implications for diagnostic strategies. *Arch Neurol*. Jul 2005; 62(7): 1091-4. PMID 16009764
25. Tikka S, Mykkänen K, Ruchoux MM, et al. Congruence between NOTCH3 mutations and GOM in 131 CADASIL patients. *Brain*. Apr 2009; 132(Pt 4): 933-9. PMID 19174371
26. Dotti MT, Federico A, Mazzei R, et al. The spectrum of Notch3 mutations in 28 Italian CADASIL families. *J Neurol Neurosurg Psychiatry*. May 2005; 76(5): 736-8. PMID 15834039
27. Dichgans M, Markus HS, Salloway S, et al. Donepezil in patients with subcortical vascular cognitive impairment: a randomised double-blind trial in CADASIL. *Lancet Neurol*. Apr 2008; 7(4): 310-8. PMID 18296124
28. De Maria R, Campolo J, Frontali M, et al. Effects of sapropterin on endothelium-dependent vasodilation in patients with CADASIL: a randomized controlled trial. *Stroke*. Oct 2014; 45(10): 2959-66. PMID 25184356
29. Huang L, Yang Q, Zhang L, et al. Acetazolamide improves cerebral hemodynamics in CADASIL. *J Neurol Sci*. May 15 2010; 292(1-2): 77-80. PMID 20227091

MEDICAL POLICY

POLICY TITLE	GENETIC TESTING OF CADASIL SYNDROME
POLICY NUMBER	MP 2.247

30. Peters N, Freilinger T, Opherck C, et al. Effects of short term atorvastatin treatment on cerebral hemodynamics in CADASIL. J Neurol Sci. Sep 15 2007; 260(1-2): 100-5. PMID 17531269
31. Meschia JF, Worrall BB, Elahi FM, et al. Management of Inherited CNS Small Vessel Diseases: The CADASIL Example: A Scientific Statement From the American Heart Association. Stroke. Oct 2023; 54(10): e452-e464. PMID 37602377

POLICY HISTORY

MP 2.247	04/10/2020 Consensus Review. No change to policy statement. References checked and updated. Coding checked with no changes.
	05/19/2021 Minor Review. To align with BCBSA's policy stance, skin biopsy was taken out as an MN requirement in the 1 st and 2 nd bullet point. Updated Policy Guidelines and References. No changes to coding.
	05/19/2022 Consensus Review. References updated. Coding reviewed.
	10/01/2022 Administrative Update. New ICD10 codes added to policy from new code review.
	08/31/2023 Consensus Review. No change to policy statement. Product Variation language and Background updated.
	10/01/2023 Administrative Update. New ICD10 codes added to policy from new code review.
	10/08/2024 Consensus Review. No change to policy statement. Updated references. Coding reviewed and no changes.
	09/25/2025 Consensus Review. No change to policy statement. Updated background, ICD-10-CM diagnosis code table, and references. Coding reviewed.
	03/09/2026 Retirement Review. Services managed by EviCore.

Health care benefit programs issued or administered by Capital Blue Cross and/or its subsidiaries, Capital Advantage Insurance Company®, Capital Advantage Assurance Company®, and Keystone Health Plan® Central. Independent licensees of the Blue Cross BlueShield Association. Communications issued by Capital Blue Cross in its capacity as administrator of programs and provider relations for all companies.