

POLICY TITLE	ABLATION PROCEDURES FOR PERIPHERAL NEUROMAS
POLICY NUMBER	MP-2.084

Original Issue Date (Created):	12/1/2011
Most Recent Review Date (Revised):	4/15/2020
Effective Date:	7/1/2020

- | | | |
|----------------------------------|--|---|
| POLICY RATIONALE | PRODUCT VARIATIONS DEFINITIONS | DESCRIPTION/BACKGROUND BENEFIT VARIATIONS |
| DISCLAIMER | CODING INFORMATION | REFERENCES |
| POLICY HISTORY | | |

I. POLICY

Minimally invasive ablation procedures, radiofrequency ablation (RFA), and cryoablation are considered **investigational** for treatment of peripheral neuromas. There is insufficient evidence to support a conclusion concerning the health outcomes or benefits associated with this procedure.

Cross-references:

- MP-2.034** Extracorporeal Shock Wave Treatment for Plantar Fasciitis and Other Musculoskeletal Conditions.
- MP-2.018** Foot Care Services

II. PRODUCT VARIATIONS

[Top](#)

This policy is only applicable to certain programs and products administered by Capital BlueCross and subject to benefit variations as discussed in Section VI. Please see additional information below.

*Note** - The Federal Employee Program (FEP) Service Benefit Plan does not have a medical policy related to these services.

III. DESCRIPTION/BACKGROUND

[Top](#)

Neuroma

A neuroma is pathology of peripheral nerve that develops as part of a normal reparative process. Neuromas may develop after injury to a nerve or as a result of chronic irritation, pressure, stretch, poor repair of nerve lesions or previous neuromas, laceration, crush injury, or blunt trauma. Neuromas typically appear about 6 to 10 weeks after trauma, with most presenting within 1 to 12 months after injury or surgery. They may gradually enlarge over a

POLICY TITLE	ABLATION PROCEDURES FOR PERIPHERAL NEUROMAS
POLICY NUMBER	MP-2.084

period of 2 to 3 years and may or may not be painful. Pain from a neuroma may be secondary to traction on the nerve by scar tissue, compression of the sensitive nerve endings by adjacent soft tissues, ischemia of the nervous tissue, or ectopic foci of ion channels that elicit neuropathic pain. Patients may describe the pain as a low intensity dull pain, or intense paroxysmal burning pain, often triggered by external stimuli such as touch or temperature. Neuroma formation has been implicated as a contributor of neuropathic pain in residual limb pain, postthoracotomy, postmastectomy, and postherniorrhaphy pain syndromes. They may coexist with phantom pain or can predispose to it.

Morton Neuroma

Morton intermetatarsal neuroma is a common and painful compression neuropathy of the common digital nerve of the foot that may be referred to by other names, including interdigital neuroma, interdigital neuritis, and interdigital or Morton metatarsalgia. It is histologically characterized by perineural fibrosis, endoneurial edema, axonal degeneration, and local vascular proliferation. Thus, some investigators do not consider Morton neuroma to be a true neuroma; instead they consider it to be an entrapment neuropathy that occurs secondary to compression of the common digital nerve under the overlying transverse metatarsal ligament. The incidence and prevalence of Morton neuroma are not clear, but it appears 10-fold more often in women than in men with an average age at presentation of around 50 years.

Diagnosis of Morton Neuroma

Although a host of imaging methods are used to diagnosis Morton neuroma, including plain radiographs, magnetic resonance imaging, and ultrasonography, objective findings are unique to this condition and are primarily used to establish a clinical diagnosis. Thus, a patient’s toes often show splaying or divergence. Patients may describe the feeling of a “lump” on the foot bottom or a feeling of walking on a rolled-up or wrinkled sock. Clinical examination with medial and lateral compression may reproduce the painful symptoms with a palpable “click” on interspace compression (Mulder sign).

Treatment of Morton Neuroma

Management of patients diagnosed with Morton neuroma typically starts with conservative approaches, such as the use of metatarsal pads in shoes and orthotic devices that alter supination and pronation of the affected foot. These approaches try to reduce pressure and irritation of the affected nerve. They may provide relief, but do not alter the underlying pathology. There is scant evidence to support the effectiveness or comparative effectiveness of these practices. In one case series (1995), investigators evaluated a 3-stage protocol of “stepped care” through which private practice patients (N=115) advanced from stage I (education plus footwear modifications, and a metatarsal pad) to stage II (steroid injections with local anesthetic or local anesthetic alone), and into stage III (surgical resection) if stages I and II were not relieved within 3 months. Overall, 97 (85%) of 115 patients believed that pain had been reduced with the treatment program. However, 24 (21%) patients eventually required surgical excision of the nerve, and 23 (96%) of them had satisfactory results.

POLICY TITLE	ABLATION PROCEDURES FOR PERIPHERAL NEUROMAS
POLICY NUMBER	MP-2.084

Ablation Techniques

Several minimally invasive procedures to treat refractory Morton neuroma are aimed at in situ destruction of the pathology: radiofrequency ablation (RFA) and cryoablation (also known as cryoneurolysis, cryolysis, cryoanalgesia). RFA uses heat generated by an electrode that conducts electromagnetic energy into a tissue or lesion to denature proteins and destroy cells. RFA is used to ablate a wide range of tissues or lesions, including osteoid osteoma; cardiovascular system pathologies; cervical pain syndromes; liver, lung, and other cancers; and varicosities. Cryoablation uses a coolant to chill a cryoprobe to temperatures below -75°C, which when inserted into a lesion, freezes and kills the tissue. It has been used to treat Morton neuroma, other chronic nerve pain syndromes, and conditions for which RFA has been used.

This review primarily focuses on evidence for the use of RFA and cryoablation on painful neuromas, with emphasis on Morton neuroma and the comparative effectiveness of these less invasive therapies with open surgical resection of the nerve pathology.

Regulatory Status

Although radiofrequency ablation probes and generators and cryoablation equipment have been cleared for marketing by the U.S. Food and Drug Administration through the 510(k) process, none appear to be specifically indicated for treatment of Morton neuroma or any other specific peripheral neuroma.

IV. RATIONALE

[Top](#)

Summary of Evidence

For individuals who have Morton neuroma who receive radiofrequency ablation (RFA), the evidence includes case series. Relevant outcomes are symptoms, functional outcomes, and treatment-related morbidity. Three case series identified reported outcomes for RFA to treat Morton neuroma. The body of evidence is highly heterogeneous regarding RFA protocols, prior conservative management, patient characteristics, follow-up durations, outcome measures, and reporting of outcomes. Variable proportions of patients require surgery after RFA, making the benefit of RFA for avoiding more invasive treatment uncertain. The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals who have Morton neuroma who receive cryoablation, the evidence includes case series. Relevant outcomes are symptoms, functional outcomes, and treatment-related morbidity. Only two retrospective case series on the use of cryoablation to treat peripheral nerve pain were identified in our literature review. The case series were heterogeneous regarding cryoablation protocols and length of follow-up. Outcome measures did not provide information on functional end points. The evidence is insufficient to determine the effects of the technology on health outcomes.

POLICY TITLE	ABLATION PROCEDURES FOR PERIPHERAL NEUROMAS
POLICY NUMBER	MP-2.084

For individuals who have peripheral neuroma(s) other than Morton neuroma who receive ablation, the evidence is very limited: no published literature was identified. Relevant outcomes are symptoms, functional outcomes, and treatment-related morbidity. The evidence is insufficient to determine the effects of the technology on health outcomes.

V. DEFINITIONS

[Top](#)

NEUROMA is the formal term for any type of tumor comprised of nerve cells. Classification is made with respect to the specific portion of the nerve involved. For example, ganglionated neuroma is a neuroma composed of true nerve cells.

VI. BENEFIT VARIATIONS

[Top](#)

The existence of this medical policy does not mean that this service is a covered benefit under the member's health benefit plan. Benefit determinations should be based in all cases on the applicable health benefit plan language. Medical policies do not constitute a description of benefits. A member's health benefit plan governs which services are covered, which are excluded, which are subject to benefit limits and which require preauthorization. There are different benefit plan designs in each product administered by Capital BlueCross. Members and providers should consult the member's health benefit plan for information or contact Capital BlueCross for benefit information.

VII. DISCLAIMER

[Top](#)

Capital BlueCross's medical policies are developed to assist in administering a member's benefits, do not constitute medical advice and are subject to change. Treating providers are solely responsible for medical advice and treatment of members. Members should discuss any medical policy related to their coverage or condition with their provider and consult their benefit information to determine if the service is covered. If there is a discrepancy between this medical policy and a member's benefit information, the benefit information will govern. If a provider or a member has a question concerning the application of this medical policy to a specific member's plan of benefits, please contact Capital BlueCross' Provider Services or Member Services. Capital BlueCross considers the information contained in this medical policy to be proprietary and it may only be disseminated as permitted by law.

VIII. CODING INFORMATION

[Top](#)

Note: This list of codes may not be all-inclusive, and codes are subject to change at any time. The identification of a code in this section does not denote coverage as coverage is

MEDICAL POLICY

POLICY TITLE	ABLATION PROCEDURES FOR PERIPHERAL NEUROMAS
POLICY NUMBER	MP-2.084

determined by the terms of member benefit information. In addition, not all covered services are eligible for separate reimbursement.

Minimally invasive ablation procedures, radiofrequency ablation (RFA), and cryoablation, are considered investigational for treatment of peripheral neuromas; therefore, not covered:

CPT Codes ®							
64624	64632	64640	0441T				

Current Procedural Terminology (CPT) copyrighted by American Medical Association. All rights reserved.

ICD-10-CM Diagnosis Codes	Description
D36.10	Benign neoplasm of peripheral nerves and autonomic nervous system, unspecified
D36.13	Benign neoplasm of peripheral nerves and autonomic nervous system of lower limb, including hip
G57.80	Other specified mononeuropathies of unspecified lower limb
G57.81	Other specified mononeuropathies of right lower limb
G57.82	Other specified mononeuropathies of left lower limb
G57.83	Other specified mononeuropathies of bilateral lower limbs
G57.90	Unspecified mononeuropathy of unspecified lower limb
G57.91	Unspecified mononeuropathy of right lower limb
G57.92	Unspecified mononeuropathy of left lower limb
G57.93	Unspecified mononeuropathy of bilateral lower limbs

POLICY TITLE	ABLATION PROCEDURES FOR PERIPHERAL NEUROMAS
POLICY NUMBER	MP-2.084

IX. REFERENCES

[Top](#)

1. Rajput K, Reddy S, Shankar H. Painful neuromas. *Clin J Pain.* Sep 2012;28(7):639-645. PMID 22699131
2. Jain S, Mannan K. The diagnosis and management of Morton's neuroma: a literature review. *Foot Ankle Spec.* Aug 2013;6(4):307-317. PMID 23811947
3. Clinical Practice Guideline Forefoot Disorders P, Thomas JL, Blitch ELt, et al. Diagnosis and treatment of forefoot disorders. Section 3. Morton's intermetatarsal neuroma. *J Foot Ankle Surg.* Mar-Apr 2009;48(2):251-256. PMID 19232980
4. Wu KK. Morton's interdigital neuroma: a clinical review of its etiology, treatment, and results. *J Foot Ankle Surg.* Mar-Apr 1996;35(2):112-119; discussion 187-118. PMID 8722878
5. Mulder JD. The causative mechanism in morton's metatarsalgia. *J Bone Joint Surg Br.* Feb 1951;33-B(1):94-95. PMID 14814167
6. Adams WR, 2nd. Morton's neuroma. *Clin Podiatr Med Surg.* Oct 2010;27(4):535-545. PMID 20934103
7. Thomson CE, Gibson JN, Martin D. Interventions for the treatment of Morton's neuroma. *Cochrane Database Syst Rev.* 2004(3):CD003118. PMID 15266472
8. Bennett GL, Graham CE, Mauldin DM. Morton's interdigital neuroma: a comprehensive treatment protocol. *Foot Ankle Int.* Dec 1995;16(12):760-763. PMID 8749346
9. Nashi M, Venkatachalam AK, Muddu BN. Surgery of Morton's neuroma: dorsal or plantar approach? *J R Coll Surg Edinb.* Feb 1997;42(1):36-37. PMID 9046143
10. Pace A, Scammell B, Dhar S. The outcome of Morton's neurectomy in the treatment of metatarsalgia. *Int Orthop.* Apr 2010;34(4):511-515. PMID 19484237
11. Kasparek M, Schneider W. Surgical treatment of Morton's neuroma: clinical results after open excision. *Int Orthop.* Sep 2013;37(9):1857-1861. PMID 23851648
12. Dierselhuis EF, van den Eerden PJ, Hoekstra HJ, et al. Radiofrequency ablation in the treatment of cartilaginous lesions in the long bones: results of a pilot study. *Bone Joint J.* Nov 2014;96-B(11):1540-1545. PMID 25371471
13. Boersma D, van Eekeren RR, Kelder HJ, et al. Mechanochemical endovenous ablation versus radiofrequency ablation in the treatment of primary small saphenous vein insufficiency (MESSI trial): study protocol for a randomized controlled trial. *Trials.* Oct 29 2014;15(1):421. PMID 25354769
14. Di Costanzo GG, Tortora R, D'Adamo G, et al. Radiofrequency ablation versus laser ablation for the treatment of small hepatocellular carcinoma in cirrhosis: a randomized trial. *J Gastroenterol Hepatol.* Mar 2015;30(3):559-565. PMID 25251043

POLICY TITLE	ABLATION PROCEDURES FOR PERIPHERAL NEUROMAS
POLICY NUMBER	MP-2.084

15. Anchala PR, Irving WD, Hillen TJ, et al. Treatment of metastatic spinal lesions with a navigational bipolar radiofrequency ablation device: a multicenter retrospective study. *Pain Physician*. Jul-Aug 2014;17(4):317-327. PMID 25054391
16. Hillen TJ, Anchala P, Friedman MV, et al. Treatment of metastatic posterior vertebral body osseous tumors by using a targeted bipolar radiofrequency ablation device: technical note. *Radiology*. Oct 2014;273(1):261-267. PMID 24927327
17. Wang X, Wang X, Song Y, et al. Efficiency of radiofrequency ablation for surgical treatment of chronic atrial fibrillation in rheumatic valvular disease. *Int J Cardiol*. Jul 01 2014;174(3):497-502. PMID 24820759
18. Huang WZ, Wu YM, Ye HY, et al. Comparison of the outcomes of monopolar and bipolar radiofrequency ablation in surgical treatment of atrial fibrillation. *Chin Med Sci J*. Mar 2014;29(1):28-32. PMID 24698675
19. Avery J, Kumar K, Thakur V, et al. Radiofrequency ablation as first-line treatment of varicose veins. *Am Surg*. Mar 2014;80(3):231-235. PMID 24666862
20. Hiraki T, Gobara H, Iguchi T, et al. Radiofrequency ablation as treatment for pulmonary metastasis of colorectal cancer. *World J Gastroenterol*. Jan 28 2014;20(4):988-996. PMID 24574771
21. Morillo CA, Verma A, Connolly SJ, et al. Radiofrequency ablation vs antiarrhythmic drugs as first-line treatment of paroxysmal atrial fibrillation (RAAFT-2): a randomized trial. *JAMA*. Feb 19 2014;311(7):692-700. PMID 24549549
22. Fuller CW, Nguyen SA, Lohia S, et al. Radiofrequency ablation for treatment of benign thyroid nodules: systematic review. *Laryngoscope*. Jan 2014;124(1):346-353. PMID 24122763
23. Huang XM, Hu JQ, Li ZF, et al. Symptomatic sinus tachycardia with perpetuating slow pathway: successful treatment with radiofrequency ablation. *Pacing Clin Electrophysiol*. Oct 2014;37(10):e1-4. PMID 21077914
24. Prologo JD, Passalacqua M, Patel I, et al. Image-guided cryoablation for the treatment of painful musculoskeletal metastatic disease: a single-center experience. *Skeletal Radiol*. Nov 2014;43(11):1551-1559. PMID 24972918
25. Kim EH, Tanagho YS, Saad NE, et al. Comparison of laparoscopic and percutaneous cryoablation for treatment of renal masses. *Urology*. May 2014;83(5):1081-1087. PMID 24560975
26. Durand M, Barret E, Galiano M, et al. Focal cryoablation: a treatment option for unilateral low-risk prostate cancer. *BJU Int*. Jan 2014;113(1):56-64. PMID 24053685
27. Duarte R, Pereira T, Pinto P, et al. [Percutaneous image-guided cryoablation for localized bone plasmacytoma treatment]. *Radiologia*. Sep-Oct 2014;56(5):e1-4. PMID 22621822
28. Rodriguez-Entem FJ, Exposito V, Gonzalez-Enriquez S, et al. Cryoablation versus radiofrequency ablation for the treatment of atrioventricular nodal reentrant tachycardia:

POLICY TITLE	ABLATION PROCEDURES FOR PERIPHERAL NEUROMAS
POLICY NUMBER	MP-2.084

- results of a prospective randomized study. J Interv Card Electrophysiol. Jan 2013;36(1):41-45; discussion 45. PMID 23080326*
29. Yamauchi Y, Izumi Y, Hashimoto K, et al. Percutaneous cryoablation for the treatment of medically inoperable stage I non-small cell lung cancer. *PLoS One.* 2012;7(3):e33223. PMID 22413004
 30. Collins KK, Schaffer MS. Use of cryoablation for treatment of tachyarrhythmias in 2010: survey of current practices of pediatric electrophysiologists. *Pacing Clin Electrophysiol.* Mar 2011;34(3):304-308. PMID 21077912
 31. Kaufman CS, Bachman B, Littrup PJ, et al. Cryoablation treatment of benign breast lesions with 12-month follow-up. *Am J Surg.* Oct 2004;188(4):340-348. PMID 15474424
 32. Genon MP, Chin TY, Bedi HS, et al. Radio-frequency ablation for the treatment of Morton's neuroma. *ANZ J Surg.* Sep 2010;80(9):583-585. PMID 20857612
 33. Moore JL, Rosen R, Cohen J, et al. Radiofrequency thermoneurolysis for the treatment of Morton's neuroma. *J Foot Ankle Surg.* Jan-Feb 2012;51(1):20-22. PMID 22055491
 34. Chuter GS, Chua YP, Connell DA, et al. Ultrasound-guided radiofrequency ablation in the management of interdigital (Morton's) neuroma. *Skeletal Radiol.* Jan 2013;42(1):107-111. PMID 23073898
 35. Friedman T, Richman D, Adler R. Sonographically guided cryoneurolysis: preliminary experience and clinical outcomes. *J Ultrasound Med.* Dec 2012;31(12):2025-2034. PMID 23197557
 36. Cazzato RL, Garnon J, Ramamurthy N, et al. Percutaneous MR-guided cryoablation of Morton's neuroma: rationale and technical details after the first 20 patients. *Cardiovasc Intervent Radiol.* Oct 2016;39(10):1491-1498. PMID 27189181
 37. Association of Extremity Nerve Surgeons (AENS). *Clinical Practice Guidelines, Edition 1. 2014*; <https://www.aens.us/images/aens/AENSGuidelinesFinal-12082014.pdf>. Accessed April 13, 2020.
 38. Blue Cross Blue Shield Association Medical Policy Reference Manual. 7.01.147, Ablation Procedures for Peripheral Neuromas July 2019.

X. POLICY HISTORY

[Top](#)

MP-2.084	CAC 7/26/11 New policy. No previous policy statements regarding this treatment. Standard FEP variation in place.
	7/24/13 Admin coding review complete
	CAC 9/24/13 Consensus review. No change to policy statements. References reviewed and updated.
	CAC 9/30/14 Consensus review. References updated. Rationale added. No changes to the policy statements.

MEDICAL POLICY



POLICY TITLE	ABLATION PROCEDURES FOR PERIPHERAL NEUROMAS
POLICY NUMBER	MP-2.084

	<p>CAC 9/29/15 Minor review. BCBSA adopted. Changed “Stereotactic radiofrequency thermal lesioning, or radiofrequency nerve ablation” to “Minimally invasive ablation procedures, radiofrequency ablation (RFA), and cryoablation”. Changed “for treatment of foot conditions” to “treatment of peripheral neuromas”. Changed title to Ablation Procedures for Peripheral Neuromas. Formerly Radiofrequency Thermal Lesioning for Treatment of Foot Conditions. Revised rationale and references. Coding reviewed.</p>
	<p>CAC 7/26/16 Consensus review. No change to policy statements. Background, rationale and references reviewed. Coding updated.</p>
	<p>10/24/16 Administrative Update Variation reformatting</p>
	<p>CAC 9/26/17 Consensus review. Policy statement unchanged. Medicare variation added d/t difference in denial reason. Description/Background, Rationale and Reference sections updated. Coding Reviewed.</p>
	<p>1/1/18 Admin Update: Medicare variations removed from Commercial Policies.</p>
	<p>6/7/18 Consensus review. No change to policy statements. References reviewed. Condensed rationale.</p>
	<p>4/8/19 Consensus review. No change to policy statements. References, background and summary of evidence reviewed.</p>
	<p>1/1/2020 Admin Coding Update: Added new code 64624.</p>
	<p>4/15/2020 Consensus review. No change to policy statement. Surgical treatment section removed to match BCBSA policy. References and coding reviewed. Added diagnosis codes.</p>

[Top](#)

Health care benefit programs issued or administered by Capital BlueCross and/or its subsidiaries, Capital Advantage Insurance Company®, Capital Advantage Assurance Company® and Keystone Health Plan® Central. Independent licensees of the BlueCross BlueShield Association. Communications issued by Capital BlueCross in its capacity as administrator of programs and provider relations for all companies.