

MEDICAL POLICY

POLICY TITLE	GENETIC TESTING FOR DIAGNOSIS AND MANAGEMENT OF MENTAL HEALTH CONDITIONS
POLICY NUMBER	MP 2.264

CLINICAL BENEFIT	☐ MINIMIZE SAFETY RISK OR CONCERN. ☒ MINIMIZE HARMFUL OR INEFFECTIVE INTERVENTIONS. ☐ ASSURE APPROPRIATE LEVEL OF CARE. ☐ ASSURE APPROPRIATE DURATION OF SERVICE FOR INTERVENTIONS. ☒ ASSURE THAT RECOMMENDED MEDICAL PREREQUISITES HAVE BEEN MET. ☐ ASSURE APPROPRIATE SITE OF TREATMENT OR SERVICE.
Effective Date:	RETIRED 7/1/2026

POLICY

Genetic testing for diagnosis and management of mental health disorders is considered **investigational** in all situations, including but not limited to the following:

- To confirm a diagnosis of a mental health disorder in an individual with symptoms.
- To predict future risk of a mental health disorder in an asymptomatic individual.
- To inform the selection or dose of medications used to treat mental health disorders, including but not limited to the following medications:
 - selective serotonin reuptake inhibitors;
 - selective norepinephrine reuptake inhibitors and serotonin-norepinephrine reuptake inhibitors;
 - tricyclic antidepressants;
 - antipsychotic drugs.

Genetic testing panels for mental health disorders, including but not limited to the Genecept Assay, STA²R test, the GeneSight Psychotropic panel, the Proove Opioid Risk assay, and the Mental Health DNA Insight panel, are considered **investigational** for all indications. There is insufficient evidence to support a general conclusion concerning the health outcomes or benefits associated with this procedure.

Cross-References:

MP 2.234 Cytochrome p450 Genotype Guided Treatment Strategy

MP 2.253 Genetic Testing for Inherited Thrombophilia

MP 2.323 General Approach to Evaluating the Utility of Genetic Panels

PRODUCT VARIATIONS

This policy is only applicable to certain programs and products administered by Capital Blue Cross and subject to benefit variations. Please see additional information below.

FEP PPO - Refer to FEP Medical Policy Manual. The FEP Medical Policy manual can be found at: <https://www.fepblue.org/benefit-plans/medical-policies-and-utilization-management-guidelines/medical-policies>

MEDICAL POLICY

POLICY TITLE	GENETIC TESTING FOR DIAGNOSIS AND MANAGEMENT OF MENTAL HEALTH CONDITIONS
POLICY NUMBER	MP 2.264

DESCRIPTION/BACKGROUND

Individual genes have been shown to be associated with the risk of psychiatric disorders and specific aspects of psychiatric drug treatment such as drug metabolism, treatment response, and risk of adverse events. Commercially available testing panels include several of these genes and are intended to aid in the diagnosis and management of mental health disorders.

This evidence review assesses whether genetic testing for the diagnosis and management of mental health conditions is clinically useful. To make a clinical management decision that improves the net health outcome; the balance of benefits and harms must be better when the test is used to manage the condition than when another test or no test is used. The net health outcome can be improved if individuals receive correct therapy, or more effective therapy, or avoid unnecessary therapy, or avoid unnecessary testing.

The primary goal of pharmacogenomic testing and personalized medicine is to achieve better clinical outcomes compared to managing the condition with the standard of care. Drug response varies greatly between individuals, and genetic factors are known to play a role. However, in most cases, the genetic variation only explains a modest portion of the variance in the individual response because clinical outcomes are also affected by a wide variety of factors including alternate pathways of metabolism and patient- and disease-related factors that may affect absorption, distribution, and elimination of the drug.

Therefore, assessment of clinical utility of a pharmacogenetic test cannot be made by a chain of evidence from clinical validity data alone. In such cases, evidence evaluation requires studies that directly demonstrate that the use of the pharmacogenomic test to make management decisions alters clinical outcomes; it is not sufficient to demonstrate that the test predicts a disorder or a phenotype. Direct evidence of clinical utility is provided by studies that compare health outcomes for patients managed with or without the test. Because these are intervention studies, the preferred evidence is from randomized controlled trials.

REGULATORY STATUS

Clinical laboratories may develop and validate tests in-house and market them as a laboratory service; laboratory-developed tests must meet the general regulatory standards of the Clinical Laboratory Improvement Amendments. The tests discussed in this section are available under the auspices of the Clinical Laboratory Improvement Amendments. Laboratories that offer laboratory-developed tests must be licensed by the Clinical Laboratory Improvement Amendments for high-complexity testing. To date, the U.S. Food and Drug Administration has chosen not to require any regulatory review of this test.

Examples of commercially available panels include the following:

- Genecept™ Assay (Genomind);
- STA²R test (SureGene Test for Antipsychotic and Antidepressant Response; Clinical Reference Laboratory). Specific variants included in the panel were not easily identified from the manufacturer's website.

MEDICAL POLICY

POLICY TITLE	GENETIC TESTING FOR DIAGNOSIS AND MANAGEMENT OF MENTAL HEALTH CONDITIONS
POLICY NUMBER	MP 2.264

- GeneSight® Psychotropic panel (Assurex Health);
- Mental Health DNA Insight™ panel (Pathway Genomics);
- IDgenetix-branded tests (AltheaDx).

Also, many labs offer genetic testing for individual genes, including *MTFHR* (GeneSight Rx and other laboratories), cytochrome P450 variants, and *SULT4A1*.

AltheaDx offers a number of IDgenetix-branded tests, which include several panels focusing on variants that affect medication pharmacokinetics for a variety of disorders, including psychiatric disorders.

RATIONALE

SUMMARY OF EVIDENCE

For individuals who are evaluated for diagnosis or risk of a mental illness who receive genetic testing for risk of that disorder, the evidence includes various observational studies (cohort, case-control, genome-wide association study). Relevant outcomes are changes in disease status, morbid events, functional outcomes, health status measures, quality of life, and treatment-related morbidity. Most studies evaluated the association between genotype and mental health disorders or gene-drug interactions among individuals at risk for mental health conditions. No studies were identified that evaluated whether testing for variants changed clinical management or affected health outcomes. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For adult individuals with major depressive disorder (MDD) who receive GeneSight testing guided drug treatment, the evidence includes 4 randomized controlled trials (RCTs). Relevant outcomes are symptoms, change in disease status, morbid events, functional outcomes, health status measures, quality of life, and treatment-related morbidity. The RCTs compared response ($\geq 50\%$ decrease in Hamilton Depression Rating Scale-17 [HAM-D17] or Patient Health Questionnaire-9 [PHQ-9]), remission (HAM-D17 ≤ 7 or PHQ-9 ≤ 5), and symptom improvement (mean % change in HAM-D17 or PHQ-9) with antidepressant therapy informed by GeneSight test results to antidepressant therapy selected without GeneSight test results (i.e., standard of care [SOC]). The PRecision Medicine In Mental Health Care (PRIME Care) trial did not find a statistically significant difference between GeneSight guided treatment and SOC in the primary outcome of remission at 24 weeks follow-up, but significant differences in the secondary outcome of symptom score improvement and treatment response were observed, favoring the GeneSight group. However, this trial had a high loss to follow-up (21%) and had inadequate participant recruitment based on a priori sample size estimation and power analysis. The GUIDED trial reported statistically significant improvements in response and remission in the GeneSight arm compared to SOC at 8 weeks among individuals with MDD. However, depending on the population (intention to treat [ITT] or per protocol), up to one-third of GUIDED randomized participants were missing from the reported results; the extent of missing data following randomization precludes conclusions on outcomes at 8 weeks. The GAPP-MDD trial,

MEDICAL POLICY

POLICY TITLE	GENETIC TESTING FOR DIAGNOSIS AND MANAGEMENT OF MENTAL HEALTH CONDITIONS
POLICY NUMBER	MP 2.264

also comparing GeneSight guided treatment with SOC, found no statistically significant differences between groups in response, remission or symptom improvement at 8 weeks follow-up, although like the GUIDED trial, a high proportion (up to 69%) of randomized participants were excluded from outcome analysis and the study was not adequately powered to detect between-group differences. In the third trial, a small, single-center pilot study by Winner et al (2013), depression outcomes did not differ significantly between GeneSight-guided care and SOC groups at the 10-week follow-up, though the study was underpowered to detect significant differences in outcomes between study arms. All of these trials have major limitations in design and conduct and in consistency and precision, thus none provided adequate evidence. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For adult individuals with MDD who receive NeuroIDgenetix testing guided drug treatment, the evidence includes 2 RCTs. Relevant outcomes are symptoms, changes in disease status, morbid events, functional outcomes, health status measures, quality of life, and treatment-related morbidity. Bradley et al (2018) conducted a double-blind RCT among patients with MDD and reported statistically significant improvement in response ($\geq 50\%$ decrease in HAM-D17) in the NeuroIDgenetix arm (64% of 140) compared to SOC (46% of 121) at 12 weeks ($p=.01$) and significant improvement in remission ($\text{HAM-D17} \leq 7$) in the NeuroIDgenetix arm (35% of 40) compared to SOC (13% of 53) at 12 weeks ($p=.02$). There was evidence of reporting bias and it was unclear if the analysis was based on ITT population; there was also high loss to follow-up (15%). In the RCT conducted by Olson et al (2017), among patients with neuropsychiatric disorders, those receiving SOC reported significantly more adverse events (53%) than those receiving NeuroIDgenetix-guided care (28%), however, the study did not report the number of patients included in this analysis. The study did not describe the randomization procedure, and in clinicalTrials.gov, neurocognitive measures were listed as co-primary outcomes, which were not reported, suggesting possible selective reporting. None of these trials provided adequate evidence. The Olson et al (2017) study had major relevance limitations and both studies have major limitations in design and conduct and in consistency and precision. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For adult individuals with MDD who receive Neuropharmagen testing guided drug treatment, the evidence includes 2 RCTs. Relevant outcomes are symptoms, changes in disease status, morbid events, functional outcomes, health status measures, quality of life, and treatment-related morbidity. The 2 RCTs compared response ($\geq 50\%$ decrease in HAM-D17) and remission ($\text{HAM-D17} \leq 7$) with antidepressant therapy informed by Neuropharmagen test results to antidepressant therapy selected without Neuropharmagen test results (i.e., SOC). The single-blinded RCT by Han et al (2018) reported statistically significant improvement in response (72% of 52 vs. 44% of 48; $p=.01$) but no statistically significant improvement in remission (46% of 52 vs. 26% of 48; $p=.07$) in the Neuropharmagen arm compared to SOC at 8 weeks among patients with MDD. The study reported an early dropout of 25% in guided-care and 38% in the standard care arm and used the last observation carried forward (LOCF) approach in the ITT analysis of effectiveness. Use of LOCF assumes data are missing completely at random, which

MEDICAL POLICY

POLICY TITLE	GENETIC TESTING FOR DIAGNOSIS AND MANAGEMENT OF MENTAL HEALTH CONDITIONS
POLICY NUMBER	MP 2.264

is unlikely to hold in this analysis. Also, the study did not report registration in any clinical trial database. The single-blinded RCT by Perez et al (2017) reported non-statistically significant improvement in response (45% of 141 vs. 40% of 139; $p=.39$) and remission (34% of 141 vs. 33% of 139; $p=.87$) in the Neuropharmagen arm compared to SOC at 12 weeks among individuals with MDD. Response and remission data were missing for 9% of individuals in the guided care group and 14% in the SOC group. None of these trials provided adequate evidence. Both studies have major limitations in design and conduct and in consistency and precision. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals with a mental illness other than depression who are undergoing drug treatment who receive genetic testing for genes associated with medication pharmacokinetics and pharmacodynamics, the evidence includes a systematic review and meta-analysis and RCTs evaluating associations between specific genes and outcomes of drug treatment. Relevant outcomes are symptoms, changes in disease status, morbid events, functional outcomes, health status measures, quality of life, and treatment-related morbidity. The systematic review and meta-analysis by Hartwell et al (2020) included 7 RCTs and reported no significant moderating effect of rs1799971, a single nucleotide polymorphism (SNP) that encodes a non-synonymous substitution (Asn40Asp) in the mu-opioid receptor gene, OPRM1 on response to naltrexone treatment of alcohol use disorder. Bradley et al (2018) conducted a double-blind RCT among individuals with anxiety disorders and reported statistically significant improvement in response ($\geq 50\%$ decrease in Hamilton Rating Scale for Anxiety [HAM-A]) in the NeuroIDgenetix arm (63% of 82) compared to SOC (50% of 95) at 12 weeks among a moderate and severe group of patients ($p=.04$). There was evidence of reporting bias and, it was unclear if the analysis was based on the ITT population. Furthermore, among the randomized moderate and severe anxiety patients with only anxiety, 25% in the experimental arm and 17% in the SOC arm were lost to follow-up over the 12-week period. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

DEFINITIONS

N/A

DISCLAIMER

Capital Blue Cross' medical policies are used to determine coverage for specific medical technologies, procedures, equipment, and services. These medical policies do not constitute medical advice and are subject to change as required by law or applicable clinical evidence from independent treatment guidelines. Treating providers are solely responsible for medical advice and treatment of members. These policies are not a guarantee of coverage or payment. Payment of claims is subject to a determination regarding the member's benefit program and eligibility on the date of service, and a determination that the services are medically necessary and appropriate. Final processing of a claim is based upon the terms of contract that applies to the members' benefit program, including benefit limitations and

MEDICAL POLICY

POLICY TITLE	GENETIC TESTING FOR DIAGNOSIS AND MANAGEMENT OF MENTAL HEALTH CONDITIONS
POLICY NUMBER	MP 2.264

exclusions. If a provider or a member has a question concerning this medical policy, please contact Capital Blue Cross' Provider Services or Member Services.

CODING INFORMATION

Note: This list of codes may not be all-inclusive, and codes are subject to change at any time. The identification of a code in this section does not denote coverage as coverage is determined by the terms of member benefit information. In addition, not all covered services are eligible for separate reimbursement.

Genetic testing for mutations associated with mental health disorders are Investigational; therefore, not covered:

Procedure Codes								
0173U	0175U	0291U	0292U	0293U	0345U	0347U	0348U	0349U
0392U	0411U	0419U	0423U	0434U	0437U	0438U	0460U	0461U
81225	81226	81230	81231	81401	81418	81479	81599	0622U
0627U								

REFERENCES

1. Koyama E, Zai CC, Bryushkova L, et al. Predicting risk of suicidal ideation in youth using a multigene panel for impulsive aggression. *Psychiatry Res.* Mar 2020; 285: 112726. PMID 31870620
2. Ghafouri-Fard S, Taheri M, Omrani MD, et al. Application of Single-Nucleotide Polymorphisms in the Diagnosis of Autism Spectrum Disorders: A Preliminary Study with Artificial Neural Networks. *J Mol Neurosci.* Aug 2019; 68(4): 515-521. PMID 30937628
3. Ran L, Ai M, Wang W, et al. Rare variants in SLC6A4 cause susceptibility to major depressive disorder with suicidal ideation in Han Chinese adolescents and young adults. *Gene.* Feb 05 2020; 726: 144147. PMID 31629822
4. Wan L, Zhang G, Liu M, et al. Sex-specific effects of methylenetetrahydrofolate reductase polymorphisms on schizophrenia with methylation changes. *Compr Psychiatry.* Oct 2019; 94: 152121. PMID 31476590
5. Zhu D, Yin J, Liang C, et al. CACNA1C (rs1006737) may be a susceptibility gene for schizophrenia: An updated meta-analysis. *Brain Behav.* Jun 2019; 9(6): e01292. PMID 31033230
6. Schröter K, Brum M, Brunkhorst-Kanaan N, et al. Longitudinal multi-level biomarker analysis of BDNF in major depression and bipolar disorder. *Eur Arch Psychiatry Clin Neurosci.* Mar 2020; 270(2): 169-181. PMID 30929061
7. Chen X, Wang M, Zhang Q, et al. Stress response genes associated with attention deficit hyperactivity disorder: A case-control study in Chinese children. *Behav Brain Res.* May 02 2019; 363: 126-134. PMID 30707907

MEDICAL POLICY

POLICY TITLE	GENETIC TESTING FOR DIAGNOSIS AND MANAGEMENT OF MENTAL HEALTH CONDITIONS
POLICY NUMBER	MP 2.264

8. Zhang L, Hu XZ, Benedek DM, et al. Genetic predictor of current suicidal ideation in US service members deployed to Iraq and Afghanistan. *J Psychiatr Res.* Jun 2019; 113: 65-71. PMID 30904785
9. Bonin L. Pediatric unipolar depression: Epidemiology, clinical features, assessment, and diagnosis
10. Gaynes BN, Warden D, Trivedi MH, et al. What did STAR*D teach us? Results from a large-scale, practical, clinical trial for patients with depression. *Psychiatr Serv.* Nov 2009; 60(11): 1439-45. PMID 19880458
11. Browin VK. DNA test for antidepressants raises questions from FDA. *Bloomberg.* August 14 2019
12. Rohan KJ, Rough JN, Evans M, et al. A protocol for the Hamilton Rating Scale for Depression: Item scoring rules, Rater training, and outcome accuracy with data on its application in a clinical trial. *J Affect Disord.* Aug 2016; 200: 111-8. PMID 27130960
13. CADTH Common Drug Reviews. Aripiprazole (Abilify): Depression, Major Depressive Disorder (MDD). Ottawa (ON): Canadian Agency for Drugs and Technologies in Health, Copyright (c) CADTH 2016.; 2016.
14. Kroenke K, Spitzer RL, Williams JB. The PHQ-9: validity of a brief depression severity measure. *J Gen Intern Med.* Sep 2001; 16(9): 606-13. PMID 11556941
15. Costantini L, Pasquarella C, Odone A, et al. Screening for depression in primary care with Patient Health Questionnaire-9 (PHQ-9): A systematic review. *J Affect Disord.* Jan 15 2021; 279: 473-483. PMID 33126078
16. Spielmans GI, McFall JP. A comparative meta-analysis of Clinical Global Impressions change in antidepressant trials. *J Nerv Ment Dis.* Nov 2006; 194(11): 845-52. PMID 17102709
17. Sheehan DV, Harnett-Sheehan K, Raj BA. The measurement of disability. *Int Clin Psychopharmacol.* Jun 1996; 11 Suppl 3: 89-95. PMID 8923116
18. Leon AC, Olfson M, Portera L, et al. Assessing psychiatric impairment in primary care with the Sheehan Disability Scale. *Int J Psychiatry Med.* 1997; 27(2): 93-105. PMID 9565717
19. Brown L, Vranjkovic O, Li J, et al. The clinical utility of combinatorial pharmacogenomic testing for patients with depression: a meta-analysis. *Pharmacogenomics.* Jun 2020; 21(8): 559-569. PMID 32301649
20. Oslin DW, Lynch KG, Shih MC, et al. Effect of Pharmacogenomic Testing for Drug-Gene Interactions on Medication Selection and Remission of Symptoms in Major Depressive Disorder: The PRIME Care Randomized Clinical Trial. *JAMA.* Jul 12 2022; 328(2): 151-161. PMID 35819423
21. Greden JF, Parikh SV, Rothschild AJ, et al. Impact of pharmacogenomics on clinical outcomes in major depressive disorder in the GUIDED trial: A large, patient- and rater-blinded, randomized, controlled study. *J Psychiatr Res.* Apr 2019; 111: 59-67. PMID 30677646
22. Tiwari AK, Zai CC, Altar CA, et al. Clinical utility of combinatorial pharmacogenomic testing in depression: A Canadian patient- and rater-blinded, randomized, controlled trial. *Transl Psychiatry.* Mar 14 2022; 12(1): 101. PMID 35288545
23. Winner JG, Carhart JM, Altar CA, et al. A prospective, randomized, double-blind study assessing the clinical impact of integrated pharmacogenomic testing for major depressive disorder. *Discov Med.* Nov 2013; 16(89): 219-27. PMID 24229738

MEDICAL POLICY

POLICY TITLE	GENETIC TESTING FOR DIAGNOSIS AND MANAGEMENT OF MENTAL HEALTH CONDITIONS
POLICY NUMBER	MP 2.264

24. Noordam R, Avery CL, Visser LE, et al. Identifying genetic loci affecting antidepressant drug response in depression using drug-gene interaction models. *Pharmacogenomics*. Jun 2016; 17(9): 1029-40. PMID 27248517
25. International Conference on Harmonization. *Statistical principles for clinical trials*: E9. 1998
26. Newell DJ. Intention-to-treat analysis: implications for quantitative and qualitative research. *Int J Epidemiol*. Oct 1992; 21(5): 837-41. PMID 1468842
27. Bradley P, Shiekh M, Mehra V, et al. Improved efficacy with targeted pharmacogenetic-guided treatment of patients with depression and anxiety: A randomized clinical trial demonstrating clinical utility. *J Psychiatr Res*. Jan 2018; 96: 100-107. PMID 28992526
28. Olson MC, Maciel A, Garipey JF, et al. Clinical Impact of Pharmacogenetic-Guided Treatment for Patients Exhibiting Neuropsychiatric Disorders: A Randomized Controlled Trial. *Prim Care Companion CNS Disord*. Mar 16 2017; 19(2). PMID 28314093
29. Vilches S, Tuson M, Vieta E, et al. Effectiveness of a Pharmacogenetic Tool at Improving Treatment Efficacy in Major Depressive Disorder: A Meta-Analysis of Three Clinical Studies. *Pharmaceutics*. Sep 02 2019; 11(9). PMID 31480800
30. Han C, Wang SM, Bahk WM, et al. A Pharmacogenomic-based Antidepressant Treatment for Patients with Major Depressive Disorder: Results from an 8-week, Randomized, Single-blinded Clinical Trial. *Clin Psychopharmacol Neurosci*. Nov 30 2018; 16(4): 469-480. PMID 30466219
31. Pérez V, Salavert A, Espadaler J, et al. Efficacy of prospective pharmacogenetic testing in the treatment of major depressive disorder: results of a randomized, double-blind clinical trial. *BMC Psychiatry*. Jul 14 2017; 17(1): 250. PMID 28705252
32. Espadaler J, Tuson M, Lopez-Ibor JM, et al. Pharmacogenetic testing for the guidance of psychiatric treatment: a multicenter retrospective analysis. *CNS Spectr*. Aug 2017; 22(4): 315-324. PMID 27098095
33. Lachin JM. Fallacies of last observation carried forward analyses. *Clin Trials*. Apr 2016; 13(2): 161-8. PMID 26400875
34. Hartwell EE, Feinn R, Morris PE, et al. Systematic review and meta-analysis of the moderating effect of rs1799971 in OPRM1, the mu-opioid receptor gene, on response to naltrexone treatment of alcohol use disorder. *Addiction*. Aug 2020; 115(8): 1426-1437. PMID 31961981
35. Kampangkaew JP, Spellicy CJ, Nielsen EM, et al. Pharmacogenetic role of dopamine transporter (SLC6A3) variation on response to disulfiram treatment for cocaine addiction. *Am J Addict*. Jul 2019; 28(4): 311-317. PMID 31087723
36. Naumova D, Grizenko N, Sengupta SM, et al. DRD4 exon 3 genotype and ADHD: Randomised pharmacodynamic investigation of treatment response to methylphenidate. *World J Biol Psychiatry*. Jul 2019; 20(6): 486-495. PMID 29182037
37. Jukic MM, Smith RL, Haslemo T, et al. Effect of CYP2D6 genotype on exposure and efficacy of risperidone and aripiprazole: a retrospective, cohort study. *Lancet Psychiatry*. May 2019; 6(5): 418-426. PMID 31000417
38. Caudle KE, Klein TE, Hoffman JM, et al. Incorporation of pharmacogenomics into routine clinical practice: the Clinical Pharmacogenetics Implementation Consortium (CPIC) guideline development process. *Curr Drug Metab*. Feb 2014; 15(2): 209-17. PMID 24479687
39. Bousman CA, Stevenson JM, Ramsey LB, et al. Clinical Pharmacogenetics Implementation Consortium (CPIC) Guideline for CYP2D6, CYP2C19, CYP2B6, SLC6A4, and HTR2A

MEDICAL POLICY

POLICY TITLE	GENETIC TESTING FOR DIAGNOSIS AND MANAGEMENT OF MENTAL HEALTH CONDITIONS
POLICY NUMBER	MP 2.264

Genotypes and Serotonin Reuptake Inhibitor Antidepressants. Clin Pharmacol Ther. Jul 2023; 114(1): 51-68. PMID 37032427

40. *International Society of Psychiatric Genetics. Genetic Testing and Psychiatric Disorders: A Statement from the International Society of Psychiatric Genetics*

41. *Bousman CA, Bengesser SA, Aitchison KJ, et al. Review and Consensus on Pharmacogenomic Testing in Psychiatry. Pharmacopsychiatry. Jan 2021; 54(1): 5-17. PMID 33147643*

POLICY HISTORY

MP 2.264	04/21/2020 Consensus Review. Policy statement unchanged. Policy guideline, Background, Rationale, and References updated. Coding reviewed.
	05/20/2020 Administrative Update. New code 0175U added to the policy. Product Variation, Benefit Variation, and Disclaimer updated.
	02/25/2021 Major Review. Policy statement updated to include LifeKit Prescript as MN. Rationale, Background, coding, and references updated.
	12/01/2021 Administrative Update. Added 0290U-0293U
	04/07/2022 Minor Review. MindX blood tests for mood disorders added to policy as INV. Policy guidelines, Product Variations, and Summary of Evidence updated. References added.
	09/12/2022 Administrative Update. New Codes 0345U, 0347U, 0348U & 0349U added
	03/16/2023 Administrative Update. New Code 0380U added.
	06/13/2023 Administrative Update. New Code 0392U added.
	08/11/2023 Consensus Review. Regulatory status updated. References reviewed and updated. Coding reviewed.
	09/07/2023 Administrative Update. New Codes 0411U, 0419U added.
	12/13/2023 Administrative Update. New codes added: 0423U, 0437U, 0434U, & 0438U
	01/19/2024 Administrative Update. Clinical benefit added.
	06/10/2024 Administrative Update. Added 0460U and 0461U. Effective 07/01/2024.
	10/15/2024 Consensus Review. No change to policy statement. References updated.
	12/10/2024 Administrative Update. Removed code 0380U, effective 01/01/2025
	04/10/2025 Minor Review. All indications for genetic testing for mental health now investigational. Policy Guidelines removed. Background, Rationale and References updated. Removed 0032U, 0290U and 81291. Added 0173U and 81418.
	07/10/2025 Administrative Update. Removed Benefit Variations Section and updated Disclaimer.

MEDICAL POLICY

POLICY TITLE	GENETIC TESTING FOR DIAGNOSIS AND MANAGEMENT OF MENTAL HEALTH CONDITIONS
POLICY NUMBER	MP 2.264

	12/04/2025 Administrative Update. Removed code 0033U, effective 01/01/2025.
	03/05/2026 Retirement Review. Service is to be managed by the vendor Evicore.
	03/12/2026 Administrative Update. Added 0622U and 0627U as part of New Code Process. Eff date 4/1/2026

Health care benefit programs issued or administered by Capital Blue Cross and/or its subsidiaries, Capital Advantage Insurance Company[®], Capital Advantage Assurance Company[®] and Keystone Health Plan[®] Central. Independent licensees of the Blue Cross BlueShield Association. Communications issued by Capital Blue Cross in its capacity as administrator of programs and provider relations for all companies.