

MEDICAL POLICY

POLICY TITLE	SPHENOPALATINE GANGLION BLOCK
POLICY NUMBER	MP-4.046

Original Issue Date (Created):	2/1/2018
Most Recent Review Date (Revised):	7/5/2018
Effective Date:	9/1/2018

[POLICY RATIONALE](#)
[DISCLAIMER](#)
[POLICY HISTORY](#)

[PRODUCT VARIATIONS](#)
[DEFINITIONS](#)
[CODING INFORMATION](#)
[APPENDIX](#)

[DESCRIPTION/BACKGROUND](#)
[BENEFIT VARIATIONS](#)
[REFERENCES](#)

I. POLICY

Sphenopalatine ganglion blocks are considered **investigational** for all indications, including but not limited to the treatment of migraines and non-migraine headaches as there is insufficient evidence to support a conclusion concerning the health outcomes or benefits associated with this procedure.

Cross-reference:

- MP-6.020** Transcutaneous Electrical Nerve Stimulation
- MP-2.064** Biofeedback and Neurofeedback Therapy
- MP-2.006** Botulinum Toxin

II. PRODUCT VARIATIONS

[TOP](#)

This policy is applicable to all programs and products administered by Capital BlueCross unless otherwise indicated below.

FEP PPO: The FEP program dictates that all drugs, devices or biological products approved by the U.S. Food and Drug Administration (FDA) may not be considered investigational. Therefore, FDA-approved drugs, devices or biological products may be assessed on the basis of medical necessity.

III. DESCRIPTION/BACKGROUND

[TOP](#)

HEADACHES AND HEADACHE TREATMENTS

Headaches are common neurologic disorders and are among the top reasons that patients seek medical care. Headaches affect approximately 50% of the general population in a given year and over 90% of people have a lifetime history of headache.¹ The 2 most common types

MEDICAL POLICY

POLICY TITLE	SPHENOPALATINE GANGLION BLOCK
POLICY NUMBER	MP-4.046

of headache are tension-type headaches and migraines. Tension headaches have a prevalence of approximately 40%.² Diagnostic criteria include the presence of at least 2 of the following characteristics: bilateral headache location, nonpulsating pain, mild-to-moderate intensity, and headache not aggravated by physical activity.³ Migraines are the second-most common headache disorder, with a 1-year migraine prevalence of approximately 12% in the United States.² They are characterized by severe pain on 1 or both sides of the head, nausea, and, at times, disturbed vision. Migraines can be categorized by headache frequency, and by the presence or absence of aura. Chronic migraine is defined as attacks on at least 15 days per month for more than 3 months, with features of migraine on at least 8 days per month.³

Cluster headaches are less common than tension or migraine headaches, with an estimated prevalence of 0.1% of the population.² They are characterized by severe unilateral orbital, supraorbital, and/or temporal pain that also includes other symptoms in the eye and/or nose on the same side (eg, rhinorrhea, eyelid edema or drooping).

Treatment

A variety of medications are used to treat acute migraine episodes. They include medications taken at the onset of an attack to abort the attack (triptans, ergotamines) and medications to treat the pain and other symptoms of migraines once they are established (nonsteroidal anti-inflammatory drugs, antiemetics). Prophylactic medication therapy may be appropriate for people with migraines that occur more than 2 days per week. In addition to medication, behavioral treatments (eg, relaxation, cognitive therapy) are used to manage migraine headache. Botulinum toxin type A injections are a U.S. Food and Drug Administration–approved treatment for chronic migraine.

Severe acute cluster headaches may be treated with abortive therapy including breathing 100% oxygen, and triptan medications. Other medications used to treat cluster headaches include steroids, calcium channel blockers, and nerve pain medications. Due to the severity of pain associated with cluster headaches, patients may seek emergency treatment. Tension-type headaches are generally treated with over the counter pain medication.

Sphenopalatine Ganglion Block

Sphenopalatine ganglion (SPG) blocks are a proposed treatment option for chronic migraines and some severe non-migraine headaches. The SPG is a group of nerve cells located behind the bony structures of the nose. The nerve bundle is linked to the trigeminal nerve, the primary nerve involved in headache disorders. The SPG has both autonomic nerves, which in this case are associated with functions such as tearing and nasal congestion, and sensory nerves, associated with pain perception. SPG blocks involve topical application of local anesthetic to mucosa overlying the SPG. The rationale for using SPG blocks to treat headaches is that local anesthetics in low concentrations could block the sensory fibers and thereby reduce pain while maintaining autonomic function.

The proposed procedure for SPG blockade is to insert intranasally a catheter that is attached to a syringe carrying local anesthetic (eg, lidocaine, bupivacaine). Once the catheter is in place, the local anesthetic is applied to the posterior wall of the nasal cavity and reaches the

MEDICAL POLICY

POLICY TITLE	SPHENOPALATINE GANGLION BLOCK
POLICY NUMBER	MP-4.046

SPG. Some form of SPG blocking procedure has been used for many years. Originally, SPG blocks were done by inserting a cotton-tipped applicator dabbed with local anesthetic into the nose; this technique may be less accurate and effective than the currently proposed procedure. Another variation is to insert a needle into the cheek and inject local anesthetic but this no longer appears to be used because it is more invasive and can be painful. Neurostimulation of the SPG and SPG blockade with radiofrequency lesioning have been used outside of the United States,⁴ but these treatments are not cleared or approved by FDA.

Three catheter devices are commercially available in the United States for performing SPG blocks. The catheters have somewhat different designs but all are attached to syringes that contain local anesthetic. The catheters are inserted intranasally and, once in place, the local anesthetic is applied through the catheter. With 2 of the 3 commercially available catheters (the SpenoCath®, Allevio™), patients are positioned on their back with their nose pointed vertically and their head turned to the side. With the Tx360® device, patients remain seated.⁵

The company marketing the Tx360® device proposes its use in the context of the MiRx™ protocol.⁶ This 2-part protocol includes a medical component for immediate pain relief and a physical component to reduce headache recurrences. The medical component involves clinical evaluation and, if the patient is considered eligible, an SPG block procedure. The physical component can include any of a number of approaches such as physical therapy, ergonomic modifications, massage, and dietary recommendations.

The optimal number and frequency of SPG treatments is unclear. Information from the American Migraine Foundation suggests that the procedure can be repeated as often as needed to control pain.⁵ A randomized controlled trial (RCT) has described a course of treatment for migraines consisting of SPG blocks twice a week for 6 weeks (total, 12 treatments).

SGB blocks are proposed for both short- and long-term treatment of headaches and migraines. When used in the emergency setting in patients with severe acute headaches, the goal of treatment is to abort the current headache while the patient is in the emergency department. In the RCT that provided a 6-week course of treatment with SPG blocks for chronic migraine (mentioned above), short-term outcomes were assessed up to 24 hours after each treatment, and the duration and frequency of chronic migraines were assessed at 1 and 6 months after the course of treatment.

REGULATORY STATUS

The Tx360® Nasal Applicator (Tian Medical), the Allevio™ SPG Nerve Block Catheter (JET Medical), and the SpenoCath® (Dolor Technologies) are considered class I devices by the U.S. Food and Drug Administration (FDA) and are exempt from 510(k) requirements. This classification does not require submission of clinical data on efficacy but only notification of FDA prior to marketing. All 3 devices are used to apply numbing medication intranasally.

MEDICAL POLICY

POLICY TITLE	SPHENOPALATINE GANGLION BLOCK
POLICY NUMBER	MP-4.046

IV. RATIONALE

[TOP](#)

Summary of Evidence

For individuals who have chronic migraine who receive sphenopalatine ganglion (SPG) block(s), the evidence includes a randomized controlled trial (RCT) and a case report. Relevant outcomes are symptoms, functional outcomes, quality of life, and treatment-related morbidity. The randomized trial evaluated a regimen of 12 SPG blocks over 6 weeks and was double-blind and placebo-controlled. The trial found significantly greater short-term (up to 24 hours) benefits from active treatment than from placebo. There were no significant longer term effects (ie, 1 and 6 months after 12 treatments), although the trial was underpowered to detect longer term efficacy. Given that SPG blocks are being proposed as a preventive therapy for chronic migraines, evidence demonstrating reduced migraine frequency, severity, or other objective outcomes from robust trials is still needed. The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals who have severe acute headache treated in the emergency setting who receive SPG block(s), the evidence includes 1 RCT. Relevant outcomes are symptoms, functional outcomes, quality of life, and treatment-related morbidity. The randomized, double-blind, placebo-controlled trial was evaluated a single SPG block for severe acute headache of mixed etiologies. There was no statistically significant difference between active treatment and placebo for the primary outcome (pain reduction 15 minutes postintervention). The trialists did not collect pain again data until 24 hours posttreatment, at which time significantly more patients were headache-free in the active treatment arm than in the placebo arm. Additional studies, preferably RCTs, are needed to determine whether SPG blocks are an effective treatment in the emergency setting. The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals who have cluster headache who receive SPG block(s), the evidence includes case series. Relevant outcomes are symptoms, functional outcomes, quality of life, and treatment-related morbidity. Two small case series, both of which evaluate an approach for intranasal SPG blocks that differs from the intervention currently available in the United States, were identified. In these series, 40% to 50% of patients experienced complete symptom relief for a variable length of time and about 20% had treatment related complications. However, it is not clear from these series the degree to which the procedures evaluated differ in safety and efficacy from an intranasal SPG block using a device cleared by the Food and Drug Administration. Additional studies, preferably RCTs, are needed to evaluate SPG blocks for treating cluster headaches. The evidence is insufficient to determine the effects of the technology on health outcomes.

V. DEFINITIONS

[TOP](#)

N/A

MEDICAL POLICY

POLICY TITLE	SPHENOPALATINE GANGLION BLOCK
POLICY NUMBER	MP-4.046

VI. BENEFIT VARIATIONS

[TOP](#)

The existence of this medical policy does not mean that this service is a covered benefit under the member's contract. Benefit determinations should be based in all cases on the applicable contract language. Medical policies do not constitute a description of benefits. A member's individual or group customer benefits govern which services are covered, which are excluded, and which are subject to benefit limits and which require preauthorization. Members and providers should consult the member's benefit information or contact Capital BlueCross for benefit information.

VII. DISCLAIMER

[TOP](#)

Capital BlueCross medical policies are developed to assist in administering a member's benefits, do not constitute medical advice and are subject to change. Treating providers are solely responsible for medical advice and treatment of members. Members should discuss any medical policy related to their coverage or condition with their provider and consult their benefit information to determine if the service is covered. If there is a discrepancy between this medical policy and a member's benefit information, the benefit information will govern. Capital BlueCross considers the information contained in this medical policy to be proprietary and it may only be disseminated as permitted by law.

VIII. CODING INFORMATION

[TOP](#)

Note: This list of codes may not be all-inclusive, and codes are subject to change at any time. The identification of a code in this section does not denote coverage as coverage is determined by the terms of member benefit information. In addition, not all covered services are eligible for separate reimbursement.

Investigational; therefore not covered:

CPT Codes®							
64999	64505						

Current Procedural Terminology (CPT) copyrighted by American Medical Association. All Rights Reserved.

IX. REFERENCES

[TOP](#)

1. International Association for the Study of Pain (IASP). *Epidemiology of Headache*. 2011; <https://s3.amazonaws.com/rdcms-iasp/files/production/public/Content/ContentFolders/GlobalYearAgainstPain2/HeadacheFactSheets/1-Epidemiology.pdf>. Accessed July 5, 2018.
2. Singh A, Soares WE. Management strategies for acute headache in the emergency department. *Emerg Med Pract*. Jun 2012;14(6):1-23; quiz 23-24. PMID 22830180

MEDICAL POLICY

POLICY TITLE	SPHENOPALATINE GANGLION BLOCK
POLICY NUMBER	MP-4.046

3. Headache Classification Committee of the International Headache S. *The International Classification of Headache Disorders, 3rd edition (beta version)*. Cephalalgia. Jul 2013;33(9):629-808. PMID 23771276
4. Sanders M, Zuurmond WW. Efficacy of sphenopalatine ganglion blockade in 66 patients suffering from cluster headache: a 12- to 70-month follow-up evaluation. *J Neurosurg*. Dec 1997;87(6):876-880. PMID 9384398
5. American Migraine Foundation. *Sphenopalatine Ganglion Blocks in Headache Disorders*. 2016; <https://americanmigrainefoundation.org/understanding-migraine/sphenopalatine-ganglion-blocks-in-headachedisorders/>. Accessed July 5, 2018.
6. The MiRx™ Protocol. 2013; <http://www.mirxprotocol.com/>. Accessed July 5, 2018.
7. Cady R, Saper J, Dexter K, et al. A double-blind, placebo-controlled study of repetitive transnasal sphenopalatine ganglion blockade with tx360 ((R)) as acute treatment for chronic migraine. *Headache*. Jan 2015;55(1):101-116. PMID 25338927
8. Cady RK, Saper J, Dexter K, et al. Long-term efficacy of a double-blind, placebo-controlled, randomized study for repetitive sphenopalatine blockade with bupivacaine vs. saline with the Tx360 device for treatment of chronic migraine. *Headache*. Apr 2015;55(4):529-542. PMID 25828648
9. Candido KD, Massey ST, Sauer R, et al. A novel revision to the classical transnasal topical sphenopalatine ganglion block for the treatment of headache and facial pain. *Pain Physician*. Nov-Dec 2013;16(6):E769-778. PMID 24284858
10. Headache Classification Subcommittee of the International Headache S. *The International Classification of Headache Disorders: 2nd edition*. Cephalalgia. 2004;24 Suppl 1:9-160. PMID 14979299
11. Schaffer JT, Hunter BR, Ball KM, et al. Noninvasive sphenopalatine ganglion block for acute headache in the emergency department: a randomized placebo-controlled trial. *Ann Emerg Med*. May 2015;65(5):503-510. PMID 25577713
12. Felisati G, Arnone F, Lozza P, et al. Sphenopalatine endoscopic ganglion block: a revision of a traditional technique for cluster headache. *Laryngoscope*. Aug 2006;116(8):1447-1450. PMID 16885751
13. Pipolo C, Bussone G, Leone M, et al. Sphenopalatine endoscopic ganglion block in cluster headache: a reevaluation of the procedure after 5 years. *Neurol Sci*. Jun 2010;31 Suppl 1:S197-199. PMID 20464621
14. Blue Cross Blue Shield Association Medical Policy Reference Manual. 7.01.159, Sphenopalatine Ganglion Block for Headache. May 2017.

X. POLICY HISTORY

[TOP](#)

MP 4.046	CAC 9/26/17 New policy. BCBSA adopted. Sphenopalatine ganglion blocks are considered investigational for all indications, including but not
-----------------	--

MEDICAL POLICY

POLICY TITLE	SPHENOPALATINE GANGLION BLOCK
POLICY NUMBER	MP-4.046

	limited to the treatment of migraines and non-migraine headaches. FEP variation added. Coding reviewed.
	7/05/18 Consensus review. Policy statement unchanged. Description/Background, Rationale and Reference sections updated.

[Top](#)

Health care benefit programs issued or administered by Capital BlueCross and/or its subsidiaries, Capital Advantage Insurance Company®, Capital Advantage Assurance Company® and Keystone Health Plan® Central. Independent licensees of the BlueCross BlueShield Association. Communications issued by Capital BlueCross in its capacity as administrator of programs and provider relations for all companies.