

MEDICAL POLICY

POLICY TITLE	ENDOVASCULAR GRAFTS FOR ABDOMINAL AORTIC ANEURYSMS
POLICY NUMBER	MP 1.090

CLINICAL BENEFIT	<input type="checkbox"/> Minimize safety risk or concern. <input type="checkbox"/> Minimize harmful or ineffective interventions. <input type="checkbox"/> Assure appropriate level of care. <input type="checkbox"/> Assure appropriate duration of service for interventions. <input checked="" type="checkbox"/> Assure that recommended medical prerequisites have been met. <input type="checkbox"/> Assure appropriate site of treatment or service.
Effective Date:	9/1/2024

[POLICY RATIONALE](#)
[DISCLAIMER](#)
[POLICY HISTORY](#)

[PRODUCT VARIATIONS](#)
[DEFINITIONS](#)
[CODING INFORMATION](#)

[DESCRIPTION/BACKGROUND](#)
[BENEFIT VARIATIONS](#)
[REFERENCES](#)

I. POLICY

Abdominal Aortic Aneurysm Repair

The use of endoprostheses approved by the U.S. Food and Drug Administration (FDA) as a treatment of abdominal aortic aneurysms (AAA) may be considered **medically necessary** in any of the following clinical situations:

- An aneurysmal diameter greater than 5.0 cm **OR**
- An aneurysmal diameter of 4.0 - 5.0 cm that has increased in size by 0.5 cm in the last 6 months; **OR**
- An aneurysmal diameter that measures twice the size of the normal infrarenal aorta; **OR**
- A ruptured abdominal aortic aneurysm; **OR**
- Treatment of aneurysms that meet the recommended threshold for surgery in patients who are ineligible for open repair due to physical limitations or other factors.

The use of endoprostheses approved by the FDA is considered **investigational** for treatment of smaller abdominal aortic aneurysms that do not meet the current recommended threshold for surgery, as there is insufficient evidence to support a general conclusion concerning the health outcomes or benefits associated with this procedure for the above indications.

Cross-reference:

MP 1.132 Endovascular Stent Grafts for Disorders of the Thoracic Aorta

POLICY GUIDELINES

MEDICAL POLICY

POLICY TITLE	ENDOVASCULAR GRAFTS FOR ABDOMINAL AORTIC ANEURYSMS
POLICY NUMBER	MP 1.090

For treatment of ruptured abdominal aortic aneurysms with endoprotheses, several factors must be considered including the following:

- The member must be sufficiently stable to undergo detailed computed tomography examination for anatomic measurements,
- The aneurysm should be anatomically appropriate for endovascular repair, and
- Specialized personnel should be available.

To monitor for leaking of the graft after implantation, patients will typically undergo routine imaging with computed tomography or ultrasonography every 6 to 12 months, or more frequently if perivascular leaks or aneurysm enlargement are detected.

II. PRODUCT VARIATIONS

[TOP](#)

This policy is only applicable to certain programs and products administered by Capital Blue Cross and subject to benefit variations as discussed in Section VI. Please see additional information below.

FEP PPO - Refer to FEP Medical Policy Manual. The FEP Medical Policy manual can be found at: <https://www.fepblue.org/benefit-plans/medical-policies-and-utilization-management-guidelines/medical-policies>.

III. DESCRIPTION/BACKGROUND

[TOP](#)

Conventional management of a clinically significant abdominal aortic aneurysm (AAA) consists of surgical excision with the placement of a sutured woven graft. Surgical excision is associated with a perioperative mortality rate between 1% and 5%. Perioperative morbidity and mortality are highest in older female patients with cardiac, pulmonary, or kidney disease; the most common cause of death is multisystem organ failure.

Due to the high mortality rate, endovascular prostheses have been developed as a less risky and minimally invasive, catheter-based alternative to open surgical excision of AAAs. These devices are deployed across the aneurysm such that the aneurysm is effectively “excluded” from the circulation, with subsequent restoration of normal blood flow.

The main potential advantage of endovascular grafts for an AAA is that they offer a less invasive and less risky approach to the repair of abdominal aneurysms. While the use of an endovascular approach has the potential to reduce the relatively high perioperative morbidity and mortality associated with open AAA repair, use of endovascular grafts also has potential disadvantages. There are concerns about the durability of the anchoring system, aneurysm expansion, and other late complications related to the prosthetic graft. Aneurysm expansion may result from perivascular leaks, also known as endoleaks, which are a unique complication of endoprotheses. Perivascular leaks may result from an incompetent seal at one of the graft attachment sites, blood flow in aneurysm tributaries (these tributaries are ligated during open surgery), or perforation of graft fabric.

MEDICAL POLICY

POLICY TITLE	ENDOVASCULAR GRAFTS FOR ABDOMINAL AORTIC ANEURYSMS
POLICY NUMBER	MP 1.090

Several types of grafts are currently in use: straight grafts, in which both ends are anchored to the infrarenal aorta, and bifurcated grafts, in which the proximal end is anchored to the infrarenal aorta, and the distal ends are anchored to the iliac arteries. Fenestrated grafts have also been investigated. These grafts are designed with openings in the wall that can be placed across the renal or celiac arteries while still protecting vessel patency through these critical arteries. Also, extensions can be placed from inside the main endograft body into the visceral arteries to create a hemostatic seal.

In about 10% of AAA, there is an associated iliac artery aneurysm (IAA). Isolated iliac artery aneurysms are uncommon, comprising less than 2% of all abdominal aneurysmal disease. IAA is defined as a dilatation of the iliac artery of 1.5-fold the normal diameter. Despite their rarity, several reports have suggested that isolated iliac artery aneurysms have a high risk of rupture, with an associated high mortality rate of up to 80%. Most of these aneurysms remain asymptomatic. Consequently, diagnosis of iliac artery aneurysms is usually made incidentally during ultrasound, CT scanning, or angiography.

Based on results of retrospective reviews of ruptured IAA and expert consensus opinion, an IAA diameter of 3 cm is the traditional threshold above which asymptomatic iliac aneurysm repair is recommended.

Utilizing endovascular technique, has significantly decreased in-operative morbidity and mortality as well as, fewer complications and a shorter length of hospital stay. Open surgical techniques for repair of IAA, although usually straightforward, can be complicated by difficult exposures, blood loss, morbidity, and mortality.

In December 2022, the American College of Cardiology and the American Heart Association published the “Guideline for the Diagnosis and Management of Aortic Disease: A Report of the American Heart Association/American College of Cardiology Joint Committee on Clinical Practice Guidelines.” Regarding Open Versus Endovascular Repair of AAA, a 2a B-NR recommendation states “In patients with nonruptured AAA and a high perioperative risk, EVAR is reasonable to reduce the risk of 30-day morbidity, mortality, or both.” The document goes on to explain that “EVAR-2 (UK Endovascular Aneurysm Repair 2) was an RCT that evaluated outcomes of EVAR in high-risk patients. Patients were enrolled if they were determined to be unfit for open surgery, with fitness assessed using cardiac, respiratory, and renal criteria. In these patients, the trial initially showed that EVAR did not improve survival compared with the control of no intervention; however, more than a decade later, those treated with EVAR had significantly lower aneurysm-related mortality (hazard ratio, 0.55; 95% CI, 0.34–0.91). Contemporary analyses of outcomes in high-risk patients show that perioperative death after EVAR has markedly decreased (e.g., 9% in EVAR-2 versus 1.9% in the ACS national registry). Furthermore, in evaluating a propensity-matched Medicare population, postoperative complications that are more likely to affect high-risk patients, such as myocardial infarction, pneumonia, acute renal failure, and need for dialysis, were all significantly less likely to occur after infrarenal EVAR compared with open repair.

MEDICAL POLICY

POLICY TITLE	ENDOVASCULAR GRAFTS FOR ABDOMINAL AORTIC ANEURYSMS
POLICY NUMBER	MP 1.090

REGULATORY STATUS

A large number of endovascular grafts have been approved by the U.S. Food and Drug Administration (FDA) through the premarket approval (PMA) process for treatment of AAAs (see Table 1). The original PMA dates are shown. Most stents have undergone device modification, name changes, and have approved supplements to the original PMA. FDA product code MIH.

Table 1. Abdominal Aortic Stent Grafts Approved by FDA

Stent Name	PMA Applicant	Approved	PMA No.
AneuRx® Prosthesis System (AneuRx AAA Advantage Stent Graft)	Medtronic Vascular	1999	P990020
Ancure® Aortoiliac System	Guidant Endovascular Technologies	2002	P990017
Gore® Excluder®	W.L. Gore & Associates	2002	P020004
Zenith® AAA Endovascular Graft	Cook	2003	P020018
Endologix Powerlink® (Afx Endovascular AAA system)	Endologix	2004	P040002
Talent® Abdominal Stent Graft System	Medtronic	2008	P070027
Endurant® II AAA Stent Graft System	Medtronic	2010	P100021
Valiant Thoracic Stent Graft System	Medtronic	2011	P100040
Relay Thoracic Stent-Graft with Plus Delivery System	Bolton Medical	2012	P110038
Ovation™ Abdominal Stent Graft System	TriVascular	2012	P120006
Aorfix™ AAA Flexible Stent Graft System	Lombard Medical	2013	P110032
Incraft® AAA Stent Graft System	Cordis	2018	P150002
TREO® Abdominal Stent-Graft System	Bolton Medical, Inc.	2020	P190015
Alto™ Abdominal Stent Graft System	Endologix, Inc.	2020	P120006/S031

FDA: Food and Drug Administration; PMA: premarket approval.

MEDICAL POLICY

POLICY TITLE	ENDOVASCULAR GRAFTS FOR ABDOMINAL AORTIC ANEURYSMS
POLICY NUMBER	MP 1.090

IV. RATIONALE

[TOP](#)

SUMMARY OF EVIDENCE

For individuals who have AAAs eligible for open repair who receive endovascular stent grafts, the evidence includes randomized controlled trials (RCTs), systematic reviews of RCTs and cohort studies, and nonrandomized comparative studies. Relevant outcomes are overall survival, morbid events, and treatment-related mortality and morbidity. Evidence from a patient-level meta-analysis of 4 RCTs comparing endovascular aneurysm repair (EVAR) with open repair for elective treatment of AAAs has indicated that neither approach is clearly superior to the other. While EVAR is associated with an early reduction in mortality, outcomes at 5 years or longer have shown greater reintervention rates and endovascular mortality and comparable overall survival rates for EVAR and open repair. Thus, the early advantage of EVAR is offset by a higher rate of late complications over the long term. Based on these data, EVAR may be considered as an alternative to open surgery in patients who are candidates for both procedures. The evidence is sufficient to determine that the technology results in a meaningful improvement in the net health outcome.

For individuals who have ruptured AAAs who receive endovascular stent grafts, the evidence includes RCTs, systematic reviews of RCTs and nonrandomized comparative studies. Relevant outcomes are overall survival, morbid events, and treatment-related mortality and morbidity. For patients with ruptured AAAs, evidence from 4 RCTs and a patient-level meta-analysis has indicated that short- and intermediate-term survival following EVAR is comparable with open repair. Evidence from RCTs and nonrandomized matched comparisons has shown that EVAR is associated with lower perioperative morbidity. The evidence is sufficient to determine that the technology results in a meaningful improvement in the net health outcome.

For individuals who have AAAs ineligible for open repair who receive endovascular stent grafts, the evidence includes RCTs. Relevant outcomes are overall survival, morbid events, and treatment-related mortality and morbidity. At least 2 RCTs have compared EVAR with no surgical intervention for patient's ineligible for open repair, either because of aneurysm size or prohibitive surgical risk. These trials did not report superior outcomes with EVAR and thus do not support the use of EVAR in this population.

V. DEFINITIONS

[TOP](#)

AORTA is the largest artery in the body, originating from the left ventricle of the heart and extending down to the abdomen, where it branches off into two smaller arteries (the common iliacs). The aorta distributes oxygenated blood to all parts of the body through the systemic circulation. It is usually divided into five segments/sections:

- Ascending aorta—the section between the heart and the arch of aorta
- Arch of aorta—the peak part that looks somewhat like an inverted "U"

MEDICAL POLICY

POLICY TITLE	ENDOVASCULAR GRAFTS FOR ABDOMINAL AORTIC ANEURYSMS
POLICY NUMBER	MP 1.090

- Descending aorta—the section from the arch of aorta to the point where it divides into the common iliac arteries
- Thoracic aorta—the half of the descending aorta above the diaphragm o Abdominal aorta—the half of the descending aorta below the diaphragm

STENT refers to any material or device used to hold tissue in place, to maintain open blood vessels, or to provide support for a graft or anastomoses while healing is taking place.

THORACIC refers to the chest or thorax.

VI. BENEFIT VARIATIONS

[TOP](#)

The existence of this medical policy does not mean that this service is a covered benefit under the member's health benefit plan. Benefit determinations should be based in all cases on the applicable health benefit plan language. Medical policies do not constitute a description of benefits. A member's health benefit plan governs which services are covered, which are excluded, which are subject to benefit limits, and which require preauthorization. There are different benefit plan designs in each product administered by Capital Blue Cross. Members and providers should consult the member's health benefit plan for information or contact Capital Blue Cross for benefit information.

VII. DISCLAIMER

[TOP](#)

Capital Blue Cross' medical policies are developed to assist in administering a member's benefits, do not constitute medical advice and are subject to change. Treating providers are solely responsible for medical advice and treatment of members. Members should discuss any medical policy related to their coverage or condition with their provider and consult their benefit information to determine if the service is covered. If there is a discrepancy between this medical policy and a member's benefit information, the benefit information will govern. If a provider or a member has a question concerning the application of this medical policy to a specific member's plan of benefits, please contact Capital Blue Cross' Provider Services or Member Services. Capital Blue Cross considers the information contained in this medical policy to be proprietary and it may only be disseminated as permitted by law.

VIII. CODING INFORMATION

[TOP](#)

Note: This list of codes may not be all-inclusive, and codes are subject to change at any time. The identification of a code in this section does not denote coverage as coverage is determined by the terms of member benefit information. In addition, not all covered services are eligible for separate reimbursement.

MEDICAL POLICY

POLICY TITLE	ENDOVASCULAR GRAFTS FOR ABDOMINAL AORTIC ANEURYSMS
POLICY NUMBER	MP 1.090

Covered when medically necessary:

Procedure Codes							
34701	34702	34703	34704	34705	34706	34707	34708
34709	34711	34712	34713	34714	34715	34716	34717
34718	34808	34812	34813	34820	34833	34834	34839
34841	34842	34843	34844	34845	34846	34847	34848
36200	36245						

ICD-10-CM Diagnosis Codes	Description
I71.00	Dissection of unspecified site of aorta
I71.02	Dissection of abdominal aorta
I71.03	Dissection of thoracoabdominal aorta
I71.30	Abdominal aortic aneurysm, ruptured, unspecified
I71.31	Pararenal abdominal aortic aneurysm, ruptured
I71.32	Juxtarenal abdominal aortic aneurysm, ruptured
I71.33	Infrarenal abdominal aortic aneurysm, ruptured
I71.40	Abdominal aortic aneurysm, without rupture, unspecified
I71.41	Pararenal abdominal aortic aneurysm, without rupture
I71.42	Juxtarenal abdominal aortic aneurysm, without rupture
I71.43	Infrarenal abdominal aortic aneurysm, without rupture
I71.50	Thoroacoabdominal aortic aneurysm, ruptured, unspecified
I71.51	Supraceliac aneurysm of the abdominal aorta
I71.52	Paravisceral aneurysm of the abdominal aorta, ruptured
I71.60	Thoracoabdominal aortic aneurysm, without rupture, unspecified
I71.61	Supraceliac aneurysm of the abdominal aorta, without rupture
I71.62	Paravisceral aneurysm of the abdominal aorta, without rupture
I71.9	Aortic aneurysm of unspecified site, without rupture
I72.3	Aneurysm of iliac artery

Covered when medically necessary for a diagnosis from Table 2

Procedure Codes							
34710*	34830	34831	34832				

Table 2

MEDICAL POLICY

POLICY TITLE	ENDOVASCULAR GRAFTS FOR ABDOMINAL AORTIC ANEURYSMS
POLICY NUMBER	MP 1.090

ICD-10-CM Diagnosis Codes	Description
I71.00	Dissection of unspecified site of aorta
I71.02	Dissection of abdominal aorta
I71.03	Dissection of thoracoabdominal aorta
I71.30	Abdominal aortic aneurysm, ruptured, unspecified
I71.33	Infrarenal abdominal aortic aneurysm, ruptured
I71.40	Abdominal aortic aneurysm, without rupture, unspecified
I71.43	Infrarenal abdominal aortic aneurysm, without rupture
I71.9	Aortic aneurysm of unspecified site, without rupture
I72.3*	Aneurysm of iliac artery

* This is the only procedure code that includes ICD10 I72.3.

IX. REFERENCES

[TOP](#)

1. Blum U, Voshage G, Lammer J, et al. Endoluminal stent-grafts for infrarenal abdominal aortic aneurysms. *N Engl J Med.* Jan 2 1997;336(1):13-20. PMID 8970935
2. Ernst CB. Current therapy for infrarenal aortic aneurysms. *N Engl J Med.* Jan 2 1997;336(1):59-60. PMID 8970942
3. White RA, Donayre CE, Walot I, et al. Modular bifurcation endoprosthesis for treatment of abdominal aortic aneurysms. *Ann Surg.* Sep 1997;226(3):381-389; discussion 389-391. PMID 9339944
4. Zarins CK, White RA, Schwarten D, et al. AneuRx stent graft versus open surgical repair of abdominal aortic aneurysms: multicenter prospective clinical trial. *J Vasc Surg.* Feb 1999;29(2):292-305; discussion 306-298. PMID 9950987
5. Blue Cross and Blue Shield Association. *Endovascular Stent-Grafts for Abdominal Aortic Aneurysm Repair. TEC Assessment Program.* 2001;16(2).
6. *Endovascular aneurysm repair versus open repair in patients with abdominal aortic aneurysm (EVAR trial 1): randomised controlled trial.* *Lancet.* Jun 2005; 365(9478): 2179-86. PMID 15978925
7. Blankensteijn JD, de Jong SE, Prinssen M, et al. Two-year outcomes after conventional or endovascular repair of abdominal aortic aneurysms. *N Engl J Med.* Jun 9 2005;352(23):2398-2405. PMID 15944424
8. *Endovascular aneurysm repair and outcome in patients unfit for open repair of abdominal aortic aneurysm (EVAR trial 2): randomised controlled trial.* *Lancet.* Jun 2005; 365(9478): 2187-92. PMID 15978926
9. Greenhalgh RM, Brown LC, Kwong GP, et al. Comparison of endovascular aneurysm repair with open repair in patients with abdominal aortic aneurysm (EVAR trial 1), 30-day operative mortality results: randomised controlled trial. *Lancet.* Sep 4-10 2004;364(9437):843-848. PMID 15351191

MEDICAL POLICY

POLICY TITLE	ENDOVASCULAR GRAFTS FOR ABDOMINAL AORTIC ANEURYSMS
POLICY NUMBER	MP 1.090

10. Prinssen M, Verhoeven EL, Buth J, et al. A randomized trial comparing conventional and endovascular repair of abdominal aortic aneurysms. *N Engl J Med.* Oct 14 2004;351(16):1607-1618. PMID 15483279
11. Anderson PL, Arons RR, Moskowitz AJ, et al. A statewide experience with endovascular abdominal aortic aneurysm repair: rapid diffusion with excellent early results. *J Vasc Surg.* Jan 2004;39(1):10-19. PMID 14718804
12. Hua HT, Cambria RP, Chuang SK, et al. Early outcomes of endovascular versus open abdominal aortic aneurysm repair in the National Surgical Quality Improvement Program-Private Sector (NSQIP-PS). *J Vasc Surg.* Mar 2005;41(3):382-389. PMID 15838467
13. Lee WA, Carter JW, Upchurch G, et al. Perioperative outcomes after open and endovascular repair of intact abdominal aortic aneurysms in the United States during 2001. *J Vasc Surg.* Mar 2004;39(3):491-496. PMID 14981436
14. Paravastu SC, Jayarajasingam R, Cottam R, et al. Endovascular repair of abdominal aortic aneurysm. *Cochrane Database Syst Rev.* Jan 23 2014;1(1):CD004178. PMID 24453068
15. Powell JT, Sweeting MJ, Ulug P, et al. Meta-analysis of individual-patient data from EVAR-1, DREAM, OVER and ACE trials comparing outcomes of endovascular or open repair for abdominal aortic aneurysm over 5 years. *Br J Surg.* Feb 2017;104(3):166-178. PMID 28160528
16. Biancari F, Catania A, D'Andrea V. Elective endovascular vs. open repair for abdominal aortic aneurysm in patients aged 80 years and older: systematic review and meta-analysis. *Eur J Vasc Endovasc Surg.* Nov 2011;42(5):571-576. PMID 21820922
17. Ulug P, Sweeting MJ, von Allmen RS, et al. Morphological suitability for endovascular repair, non-intervention rates, and operative mortality in women and men assessed for intact abdominal aortic aneurysm repair: systematic reviews with meta-analysis. *Lancet.* Jun 24 2017;389(10088):2482-2491. PMID 28455148
18. Lederle FA, Freischlag JA, Kyriakides TC, et al. Long-term comparison of endovascular and open repair of abdominal aortic aneurysm. *N Engl J Med.* Nov 22 2012;367(21):1988-1997. PMID 23171095
19. De Bruin JL, Baas AF, Buth J, et al. Long-term outcome of open or endovascular repair of abdominal aortic aneurysm. *N Engl J Med.* May 20 2010;362(20):1881-1889. PMID 20484396
20. Greenhalgh RM, Brown LC, Powell JT, et al. Endovascular versus open repair of abdominal aortic aneurysm. *N Engl J Med.* May 20 2010;362(20):1863-1871. PMID 20382983
21. Brown LC, Thompson SG, Greenhalgh RM, et al. Incidence of cardiovascular events and death after open or endovascular repair of abdominal aortic aneurysm in the randomized EVAR trial 1. *Br J Surg.* Jul 2011;98(7):935-942. PMID 21484775
22. Becquemin JP, Pillet JC, Lescalie F, et al. A randomized controlled trial of endovascular aneurysm repair versus open surgery for abdominal aortic aneurysms in low- to moderate-risk patients. *J Vasc Surg.* May 2011;53(5):1167-1173 e1161. PMID 21276681
23. Schermerhorn ML, Buck DB, O'Malley AJ, et al. Long-term outcomes of abdominal aortic aneurysm in the Medicare population. *N Engl J Med.* Jul 23 2015;373(4):328-338. PMID 26200979

MEDICAL POLICY

POLICY TITLE	ENDOVASCULAR GRAFTS FOR ABDOMINAL AORTIC ANEURYSMS
POLICY NUMBER	MP 1.090

24. Liang NL, Reitz KM, Makaroun MS, et al. Comparable perioperative mortality outcomes in younger patients undergoing elective open and endovascular abdominal aortic aneurysm repair. *J Vasc Surg.* May 2018;67(5):1404-1409 e1402. PMID 29097041
25. Krenzien F, Matia I, Wiltberger G, et al. Outcome after open surgery repair in endovascular-suitable patients with ruptured abdominal aortic aneurysms. *Vasa.* Nov 2013;42(6):442-448. PMID 24220121
26. Sweeting MJ, Balm R, Desgranges P, et al. Individual-patient meta-analysis of three randomized trials comparing endovascular versus open repair for ruptured abdominal aortic aneurysm. *Br J Surg.* Sep 2015;102(10):1229-1239. PMID 26104471
27. Badger S, Forster R, Blair PH, et al. Endovascular treatment for ruptured abdominal aortic aneurysm. *Cochrane Database Syst Rev.* May 26 2017;5:CD005261. PMID 28548204
28. Powell JT, Sweeting MJ, Thompson MM, et al. Endovascular or open repair strategy for ruptured abdominal aortic aneurysm: 30 day outcomes from IMPROVE randomised trial. *BMJ.* Jan 13 2014; 348: f7661. PMID 244189500
29. Braithwaite B, Cheshire NJ, Greenhalgh RM, et al. Endovascular strategy or open repair for ruptured abdominal aortic aneurysm: one-year outcomes from the IMPROVE randomized trial. *Eur Heart J.* Aug 14 2015; 36(31): 2061-2069. PMID 25855369
30. Reimerink JJ, Hoornweg LL, Vahl AC, et al. Endovascular repair versus open repair of ruptured abdominal aortic aneurysms: a multicenter randomized controlled trial. *Ann Surg.* Aug 2013;258(2):248-256. PMID 23549424
31. Desgranges P, Kobeiter H, Katsahian S, et al. ECAR (Endovasculaire ou Chirurgie dans les Aneurysmes aorto-iliaques Rompus): A French randomized controlled trial of endovascular versus open surgical repair of ruptured aorto-iliac aneurysms. *Eur J Vasc Endovasc Surg.* Sep 2015;50(3):303-310. PMID 26001320
32. Edwards ST, Schermerhorn ML, O'Malley AJ, et al. Comparative effectiveness of endovascular versus open repair of ruptured abdominal aortic aneurysm in the Medicare population. *J Vasc Surg.* Mar 2014;59(3):575-582. PMID 24342064
33. Filardo G, Powell JT, Martinez MA, et al. Surgery for small asymptomatic abdominal aortic aneurysms. *Cochrane Database Syst Rev.* Mar 14 2012;3(3):CD001835. PMID 22419281
34. Cao P, De Rango P, Verzini F, et al. Comparison of surveillance versus aortic endografting for small aneurysm repair (CAESAR): results from a randomised trial. *Eur J Vasc Endovasc Surg.* Jan 2011;41(1):13-25. PMID 20869890
35. Ouriel K, Clair DG, Kent KC, et al. Endovascular repair compared with surveillance for patients with small abdominal aortic aneurysms. *J Vasc Surg.* May 2010;51(5):1081-1087. PMID 20304589
36. De Rango P, Verzini F, Parlani G, et al. Quality of life in patients with small abdominal aortic aneurysm: the effect of early endovascular repair versus surveillance in the CAESAR trial. *Eur J Vasc Endovasc Surg.* Mar 2011;41(3):324-331. PMID 21145269
37. Greenhalgh RM, Brown LC, Powell JT, et al. Endovascular repair of aortic aneurysm in patients physically ineligible for open repair. *N Engl J Med.* May 20 2010;362(20):1872-1880. PMID 20382982
38. Wilt TJ, Lederle FA, Macdonald R, et al. Comparison of endovascular and open surgical repairs for abdominal aortic aneurysm. *Evid Rep Technol Assess (Full Rep).* Aug 2006(144):1-113. PMID 17764213

MEDICAL POLICY

POLICY TITLE	ENDOVASCULAR GRAFTS FOR ABDOMINAL AORTIC ANEURYSMS
POLICY NUMBER	MP 1.090

39. Sweeting MJ, Patel R, Powell JT, et al. Endovascular repair of abdominal aortic aneurysm in patients physically ineligible for open repair: very long-term follow-up in the EVAR-2 randomized controlled trial. *Ann Surg.* Nov 2017;266(5):713-719. PMID 28742684
40. Rooke TW, Hirsch AT, Misra S, et al. 2011 ACCF/AHA focused update of the guideline for the management of patients with peripheral artery disease (updating the 2005 guideline): a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol.* Nov 1 2011;58(19):2020-2045. PMID 21963765
41. Hirsch AT, Haskal ZJ, Hertzler NR, et al. ACC/AHA 2005 guidelines for the management of patients with peripheral arterial disease (lower extremity, renal, mesenteric, and abdominal aortic): executive summary a collaborative report from the American Association for Vascular Surgery/Society for Vascular Surgery, Society for Cardiovascular Angiography and Interventions, Society for Vascular Medicine and Biology, Society of Interventional Radiology, and the ACC/AHA Task Force on Practice Guidelines (Writing Committee to Develop Guidelines for the Management of Patients With Peripheral Arterial Disease) endorsed by the American Association of Cardiovascular and Pulmonary Rehabilitation; National Heart, Lung, and Blood Institute; Society for Vascular Nursing; TransAtlantic Inter-Society Consensus; and Vascular Disease Foundation. *J Am Coll Cardiol.* Mar 21 2006;47(6):1239-1312. PMID 16545667
42. Walker TG, Kalva SP, Yeddula K, et al. Clinical practice guidelines for endovascular abdominal aortic aneurysm repair: written by the Standards of Practice Committee for the Society of Interventional Radiology and endorsed by the Cardiovascular and Interventional Radiological Society of Europe and the Canadian Interventional Radiology Association. *J Vasc Interv Radiol.* Nov 2010;21(11):1632-1655. PMID 20884242
43. Chaikof EL, Dalman RL, Eskandari MK, et al. The Society for Vascular Surgery practice guidelines on the care of patients with an abdominal aortic aneurysm. *J Vasc Surg.* Jan 2018;67(1):2-77.e72. PMID 29268916
44. Bulder RMA, Bastiaannet E, Hamming JF, et al. Meta-analysis of long-term survival after elective endovascular or open repair of abdominal aortic aneurysm. *Br J Surg.* Apr 2019; 106(5): 523-533. PMID 30883709
45. Chen ZG, Tan SP, Diao YP, et al. The long-term outcomes of open and endovascular repair for abdominal aortic aneurysm: A meta-analysis. *Asian J Surg.* Oct 2019; 42(10): 899-906. PMID 30914154
46. Li B, Khan S, Salata K, et al. A systematic review and meta-analysis of the long-term outcomes of endovascular versus open repair of abdominal aortic aneurysm. *J Vasc Surg.* Sep 2019; 70(3): 954-969.e30. PMID 31147117
47. Lederle FA, Kyriakides TC, Stroupe KT, et al. Open versus Endovascular Repair of Abdominal Aortic Aneurysm. *N Engl J Med.* May 30 2019; 380(22): 2126-2135. PMID 31141634
48. van Beek SC, Reimerink JJ, Vahl AC, et al. Outcomes after open repair for ruptured abdominal aortic aneurysms in patients with friendly versus hostile aortoiliac anatomy. *Eur J Vasc Endovasc Surg.* Apr 2014; 47(4): 380-7. PMID 24485844

MEDICAL POLICY

POLICY TITLE	ENDOVASCULAR GRAFTS FOR ABDOMINAL AORTIC ANEURYSMS
POLICY NUMBER	MP 1.090

49. Soden PA, Zettervall SL, Ultee KH, et al. Outcomes for symptomatic abdominal aortic aneurysms in the American College of Surgeons National Surgical Quality Improvement Program. *J Vasc Surg.* 2016;64(2):297-305. doi:10.1016/j.jvs.2016.02.055

50. Wanhainen A, et al., European Society for Vascular Surgery (ESVS) 2019 Clinical Practice Guidelines on the Management of Abdominal Aorto-iliac Artery Aneurysms, *European Journal of Vascular and Endovascular Surgery* (2018)

51. Scheinert D, Schröder M., Steinkamp H, Ludwig J, Biamino G. Treatment of iliac artery aneurysms by percutaneous implantation of stent grafts. *Circulation.* 2000;102(suppl_3). doi:10.1161/circ.102.suppl_3.iii-253

52. Uberoi R, Tsetis D, Shrivastava V, Morgan R, Belli AM; Subcommittee on Reporting Standards for Arterial Aneurysms of The Society for Vascular Surgery. Standard of practice for the interventional management of isolated iliac artery aneurysms. *Cardiovasc Intervent Radiol.* 2011;34(1):3-13. doi:10.1007/s00270-010-0055-0

53. Kirkwood, M. Iliac Artery Aneurysm. In: *UpToDate Online Journal [serial online]. Updated 07/26/22*

54. Kirkwood, M. Surgical and endovascular repair of Iliac artery aneurysm. In: *UpToDate Online Journal [serial online]. Updated 04/05/21*

55. Isselbacher, E., Chen, E. P., Brown-Zimmerman, M. M., Bray, B. E., Braverman, A. C., Bolen, M. A., Beck, A. W., Augoustides, J. G., Black, J. H., & Preventza, O., et al(2022, November 2). 2022 ACC/AHA guideline for the diagnosis and management of aortic ... *Circulation.* <https://www.ahajournals.org/doi/full/10.1161/CIR.0000000000001106>

56. *Blue Cross Blue Shield Association Medical Policy Reference Manual. 7.01.67 Endovascular Stent Grafts for Abdominal Aortic Aneurysms. June 2023.*

X. POLICY HISTORY

[TOP](#)

MP 1.090	01/01/2020 Administrative Update. Added new codes 34717 & 34718.
	02/21/2020 Consensus Review. Policy statement unchanged. References reviewed. Coding reviewed.
	01/19/2021 Consensus Review. Policy statement unchanged. Background and references updated.
	08/01/2022 Administrative Update. ICD10 codes updated
	08/05/2022 Consensus Review Policy statement unchanged. Updated background to include iliac artery aneurysms. Codes 34712 and 34713 added. ICD10 code I72.3 added. References update.
	05/18/2023 Minor Review. Moved statement “treatment of aneurysms that do meet the recommended threshold for surgery in patients who are ineligible for open repair due to physical limitations or other factors” from INV to MN. Updated background, references. No coding changes.
	05/17/2024 Consensus Review. No change to policy statement. References updated. Coding reviewed, no changes.

MEDICAL POLICY

POLICY TITLE	ENDOVASCULAR GRAFTS FOR ABDOMINAL AORTIC ANEURYSMS
POLICY NUMBER	MP 1.090

[Top](#)

Health care benefit programs issued or administered by Capital Blue Cross and/or its subsidiaries, Capital Advantage Insurance Company[®], Capital Advantage Assurance Company[®] and Keystone Health Plan[®] Central. Independent licensees of the Blue Cross BlueShield Association. Communications issued by Capital Blue Cross in its capacity as administrator of programs and provider relations for all companies.