

MEDICAL POLICY

POLICY TITLE	GENETIC TESTING FOR EPILEPSY
POLICY NUMBER	MP 2.262

CLINICAL BENEFIT	☐ MINIMIZE SAFETY RISK OR CONCERN. ☒ MINIMIZE HARMFUL OR INEFFECTIVE INTERVENTIONS. ☐ ASSURE APPROPRIATE LEVEL OF CARE. ☐ ASSURE APPROPRIATE DURATION OF SERVICE FOR INTERVENTIONS. ☒ ASSURE THAT RECOMMENDED MEDICAL PREREQUISITES HAVE BEEN MET. ☐ ASSURE APPROPRIATE SITE OF TREATMENT OR SERVICE.
Effective Date:	RETIRED 7/1/2026

[POLICY
RATIONALE
CODING INFORMATION](#)

[PRODUCT VARIATIONS
DEFINITIONS
REFERENCES](#)

[DESCRIPTION/BACKGROUND
DISCLAIMER
POLICY HISTORY](#)

I. POLICY

Genetic testing for genes associated with infantile- and early-childhood onset epilepsy syndromes in individuals with infantile- and early-childhood-onset epilepsy syndromes in which epilepsy is the core clinical symptom (see Policy Guidelines section) may be considered **medically necessary** if positive test results may:

1. Lead to changes in medication management; AND/OR
2. Lead to changes in diagnostic testing such that alternative potentially invasive tests are avoided; AND/OR
3. Lead to changes in reproductive decision-making.

Genetic testing for epilepsy is considered **investigational** for all other situations. There is insufficient evidence to support a general conclusion concerning the health outcomes or benefits associated with this procedure.

Policy Guidelines

Included Tests and Conditions

This policy addresses testing for epilepsy that is possibly genetic. In 2010, the International League Against Epilepsy classified epilepsy as having underlying genetic cause or etiology when, as best understood, the epilepsy is the direct result of a known or presumed genetic defect and seizures are the core symptom of the disorder and for which there is no structural or metabolic defect predisposing to epilepsy. The updated 2017 ILAE classification system does not discuss epilepsy with a genetic cause.

This policy also addresses the rare epilepsy syndromes that present in infancy or early childhood, in which epilepsy is the core clinical symptom (e.g., Dravet syndrome, early infantile epileptic encephalopathy, generalized epilepsy with febrile seizures plus, epilepsy and intellectual disability limited to females, nocturnal frontal lobe epilepsy, and others). Other

MEDICAL POLICY

POLICY TITLE	GENETIC TESTING FOR EPILEPSY
POLICY NUMBER	MP 2.262

clinical manifestations may be present in these syndromes but are generally secondary to the epilepsy itself.

Excluded Tests and Conditions

This policy does not address testing for genetic syndromes that have a wider range of symptomatology, of which seizures may be one, such as the neurocutaneous disorders (e.g., neurofibromatosis, tuberous sclerosis) or genetic syndromes associated with cerebral malformations or abnormal cortical development, or metabolic or mitochondrial disorders. Genetic testing for these syndromes may be specifically addressed in other policies (see Cross-reference section).

Testing that is limited to genotyping of *CYP450* genes is addressed separately (MP 2.234).

This policy does *not* address the use of genotyping for the HLA-B*1502 allelic variant in patients of Asian ancestry prior to considering drug treatment with carbamazepine due to risks of severe dermatologic reactions. This testing is recommended by the U.S. Food and Drug Administration (FDA) labeling for carbamazepine.

This policy also does *not* address the use of testing for variants in the mitochondrial DNA polymerase gamma (*POLG*) gene in patients with clinically suspected mitochondrial disorders prior to initiation of therapy with valproate. Valproate's label contains a black box warning related to increased risk of acute liver failure associated with the use of valproate in patients with *POLG* gene-related hereditary neurometabolic syndromes. FDA labeling states: "Valproate is contraindicated in patients known to have mitochondrial disorders caused by variants in mitochondrial DNA polymerase γ (*POLG*; e.g., Alpers-Huttenlocher Syndrome) and children under two years of age who are suspected of having a *POLG*-related disorder.

For positions on whole exome and whole genome sequencing for the diagnosis of neurodevelopmental disorders refer to MP 2.324

Medically Necessary Statement Definitions and Testing Strategy

The medically necessary statement refers to epilepsy syndromes that present in infancy or early childhood, are severe, and are characterized by epilepsy as the primary manifestation, without associated metabolic or brain structural abnormalities. As defined by the International League Against Epilepsy, these include epileptic encephalopathies, which are electroclinical syndrome associated with a high probability of encephalopathic features that present or worsen after the onset of epilepsy. Other clinical manifestations, including developmental delay and/or intellectual disability, may be present secondary to the epilepsy itself. Specific clinical syndromes based on the International League Against Epilepsy classification include:

- Dravet syndrome (also known as severe myoclonic epilepsy in infancy [SMEI] or polymorphic myoclonic epilepsy in infancy [PMEI])
- EFMR syndrome (epilepsy limited to females with intellectual disability)
- Epileptic encephalopathy with continuous spike-and-wave during sleep
- GEFS+ syndrome (generalized epilepsies with febrile seizures plus)
- Ohtahara syndrome (also known as early infantile epileptic encephalopathy with burst suppression pattern)
- Landau-Kleffner syndrome

MEDICAL POLICY

POLICY TITLE	GENETIC TESTING FOR EPILEPSY
POLICY NUMBER	MP 2.262

- West syndrome
- Glucose transporter type 1 deficiency syndrome.

Variants in a large number of genes have been associated with early-onset epilepsies. Some of these are summarized in Table PG1.

Table PG1. Single Genes Associated with Epileptic Syndromes

Syndrome	Associated Genes
Dravet syndrome	<i>SCN1A, SCN9A, GABRA1, STXBP1, PCDH19, SCN1B, CHD2, HCN1</i>
Epilepsy limited to females with intellectual disability	<i>PCDH19</i>
Epileptic encephalopathy with continuous spike-and-wave during sleep	<i>GRIN2A</i>
Genetic epilepsy with febrile seizures plus	<i>SCN1A, SCN9A</i>
Early infantile epileptic encephalopathy with suppression burst (Ohtahara syndrome)	<i>KCNQ2, SLC25A22, STXBP1, CDKL5, ARX</i>
Landau-Kleffner syndrome	<i>GRIN2A</i>
West syndrome	<i>ARX, TSC1, TSC2, CDKL5, ALG13, MAGI2, STXBP1, SCN1A, SCN2A, GABA, GABRB3, DNM1</i>
Glucose transporter type 1 deficiency syndrome	<i>SLC2A1</i>

Application of Medically Necessary Policy Statement

Although there is no standard definition of epileptic encephalopathies, they are generally characterized by at least some of the following: (1) onset in early childhood (often in infancy); (2) refractory to therapy; (3) associated with developmental delay or regression; and (4) severe electroencephalogram (EEG) abnormalities. There is a challenge in defining the population appropriate for testing given that specific epileptic syndromes may be associated with different EEG abnormalities, which may change over time, and patients may present with severe seizures prior to the onset or recognition of developmental delay or regression. However, for this policy, the medically necessary policy statement would apply for patients with:

1. Onset of seizures in early childhood (i.e., before the age of 5 years); AND
2. Clinically severe seizures that affect daily functioning and/or interictal EEG abnormalities; AND
3. No other clinical syndrome that would potentially better explain the patient's symptoms.

MEDICAL POLICY

POLICY TITLE	GENETIC TESTING FOR EPILEPSY
POLICY NUMBER	MP 2.262

Testing Strategy

There is clinical and genetic overlap for many of the electroclinical syndromes previously discussed. If there is suspicion for a specific syndrome based on history, EEG findings, and other test results, testing should begin with targeted variant testing for the candidate gene most likely to be involved, followed by sequential testing for other candidate genes. In particular, if an *SCN1A*-associated syndrome is suspected (Dravet syndrome, GEFS+), molecular genetic testing of *SCN1A* with sequence analysis of the *SCN1A* coding region, followed by deletion/duplication analysis if a pathogenic variant is not identified, should be obtained.

Given the genetic heterogeneity of early-onset epilepsy syndromes, a testing strategy that uses a multigene panel may be considered reasonable. In these cases, panels should meet the criteria outlined in MP 2.323 (General Approach to Evaluating the Utility of Genetic Panels). Criteria for use of whole exome sequencing are outlined in MP 2.324 (Whole Exome and Whole Genome Sequencing for Diagnosis of Genetic Disorders).

Genetics Nomenclature Update

The Human Genome Variation Society nomenclature is used to report information on variants found in DNA and serves as an international standard in DNA diagnostics. It is being implemented for genetic testing medical evidence review updates starting in 2017 (see Table PG2). The Society's nomenclature is recommended by the Human Variome Project, the Human Genome Organization, and by the Human Genome Variation Society itself.

The American College of Medical Genetics and Genomics (ACMG) and Association for Molecular Pathology (AMP) standards and guidelines for interpretation of sequence variants represent expert opinion from ACMG, AMP, and the College of American Pathologists. These recommendations primarily apply to genetic tests used in clinical laboratories, including genotyping, single genes, panels, exomes, and genomes. Table PG3 shows the recommended standard terminology—"pathogenic," "likely pathogenic," "uncertain significance," "likely benign," and "benign"—to describe variants identified that cause Mendelian disorders.

Table PG2. Nomenclature to Report on Variants Found in DNA

Previous	Updated	Definition
Mutation	Disease-associated variant	Disease-associated change in the DNA sequence
	Variant	Change in the DNA sequence
	Familial variant	Disease-associated variant identified in a proband for use in subsequent targeted genetic testing in first-degree relatives

Table PG3. ACMG-AMP Standards and Guidelines for Variant Classification

Variant Classification	Definition
Pathogenic	Disease-causing change in the DNA sequence
Likely pathogenic	Likely disease-causing change in the DNA sequence

MEDICAL POLICY

POLICY TITLE	GENETIC TESTING FOR EPILEPSY
POLICY NUMBER	MP 2.262

Variant of uncertain significance	Change in DNA sequence with uncertain effects on disease
Likely benign	Likely benign change in the DNA sequence
Benign	Benign change in the DNA sequence

ACMG: American College of Medical Genetics and Genomics; AMP: Association for Molecular Pathology.

Genetic Counseling

Genetic counseling is primarily aimed at patients who are at risk for inherited disorders, and experts recommend formal genetic counseling in most cases when genetic testing for an inherited condition is considered. The interpretation of the results of genetic tests and the understanding of risk factors can be very difficult and complex. Therefore, genetic counseling will assist individuals in understanding the possible benefits and harms of genetic testing, including the possible impact of the information on the individual's family. Genetic counseling may alter the utilization of genetic testing substantially and may reduce inappropriate testing. Genetic counseling should be performed by an individual with experience and expertise in genetic medicine and genetic testing methods.

Cross-References:

MP 2.234 Cytochrome P450 Genotype Guided Treatment Strategy
 MP 2.276 Genetic Testing for *FMR1* Variants (Including Fragile X Syndrome)
 MP 2.323 General Approach to Evaluating the Utility of Genetic Panels
 MP 2.324 Whole Exome and Whole Genome Sequencing for Diagnosis of Genetic Disorders
 MP 2.326 General Approach to Genetic Testing

II. PRODUCT VARIATIONS

[TOP](#)

This policy is only applicable to certain programs and products administered by Capital Blue Cross and subject to benefit variations as discussed in Section VI. Please see additional information below.

FEP PPO: Refer to FEP Medical Policy Manual. The FEP Medical Policy manual can be found at:

<https://www.fepblue.org/benefit-plans/medical-policies-and-utilization-management-guidelines/medical-policies>.

III. DESCRIPTION/BACKGROUND

[TOP](#)

Epilepsy

Epilepsy is defined as the occurrence of two or more unprovoked seizures. It is a common neurologic disorder, with approximately 3% of the population developing the disorder over their entire lifespan.

MEDICAL POLICY

POLICY TITLE	GENETIC TESTING FOR EPILEPSY
POLICY NUMBER	MP 2.262

Sudden unexplained death in epilepsy (SUDEP) is defined as a sudden, unexpected, nontraumatic, and non-drowning death in patients with epilepsy, excluding documented status epilepticus, with no cause of death identified following comprehensive postmortem evaluation.

Classification

Epilepsy is heterogeneous in etiology and clinical expression and can be classified in a variety of ways. Most commonly, classification is done by the clinical phenotype, i.e., the type of seizures that occur. In 2017, the International League Against Epilepsy (ILAE) updated its classification system that is widely used for clinical care and research purposes (see Table 1). Classification of seizures can also be done on the basis of age of onset: neonatal, infancy, childhood, and adolescent/adult.

Table 1. Classification of Seizure Disorders by Type

Focal Onset (including aware and impaired awareness)	Generalized Onset	Unknown Onset	Unclassified
Motor onset • Automatisms • Atonic^a • Clonic • Epileptic spasms^a • Hyperkinetic • Myoclonic • Tonic	Motor • Tonic-clonic • Clonic • Tonic • Myoclonic • Myoclonic-tonic-clonic • Myoclonic-atonic • Atonic • Epileptic Spasms	Motor • Tonic-clonic • Epileptic Spasms	
Nonmotor Onset • Autonomic • Behavior Arrest • Cognitive • Emotional Sensory	Nonmotor (absence) • Typical • Atypical • Myoclonic • Eyelid Myoclonia	Nonmotor • Behavior Arrest	
Focal to bilateral tonic-clonic			

Adapted from Fisher et al (2017) ^a Degree of awareness usually is not specified.

MEDICAL POLICY

POLICY TITLE	GENETIC TESTING FOR EPILEPSY
POLICY NUMBER	MP 2.262

Although genetic epilepsies are not discussed in the 2017 ILAE report², a 2010 ILAE report³, identified genetic epilepsies as conditions in which the seizures are a direct result of a known or presumed genetic defect(s). Genetic epilepsies are characterized by recurrent unprovoked seizures in patients who do not have demonstrable brain lesions or metabolic abnormalities. In addition, seizures are the core symptom of the disorder, and other symptomatology is not present, except as a direct result of seizures. This is differentiated from genetically determined conditions in which seizures are part of a larger syndrome, such as tuberous sclerosis, fragile X syndrome, or Rett syndrome.

The review focuses on the category of genetic epilepsies in which seizures are the primary clinical manifestation. This category does not include syndromes that have multiple clinical manifestations, of which seizures may be one. Examples of syndromes that include seizures are Rett syndrome and tuberous sclerosis. Genetic testing for these syndromes will not be assessed herein but may be included in separate reviews that specifically address genetic testing for that syndrome.

Genetic epilepsies can be further broken down by type of seizures. For example, genetic generalized epilepsy refers to patients who have convulsive (grand mal) seizures, while genetic absence epilepsy refers to patients with nonconvulsive (absence) seizures. The disorders are also sometimes classified by the age of onset.

The category of genetic epilepsies includes a number of rare epilepsy syndromes that present in infancy or early childhood. These syndromes are characterized by epilepsy as the primary manifestation, without associated metabolic or brain structural abnormalities. They are often severe and sometimes refractory to medication treatment. They may involve other clinical manifestations such as developmental delay and/or intellectual disability, which in many cases are thought to be caused by frequent uncontrolled seizures. In these cases, the epileptic syndrome may be classified as an epileptic encephalopathy, which is described by ILAE as disorders in which the epileptic activity itself may contribute to severe cognitive and behavioral impairments above and beyond what might be expected from the underlying pathology alone and that these can worsen over time. A partial list of severe early-onset epilepsy syndromes is as follows:

- Dravet syndrome
- EFMR syndrome (epilepsy limited to females with mental retardation)
- Nocturnal frontal lobe epilepsy
- GEFS+ syndrome (genetic epilepsy with febrile seizures plus)
- EIEE syndrome (early infantile epileptic encephalopathy with suppression burst)
- West syndrome
- Ohtahara syndrome

Dravet syndrome falls on a spectrum of *SCN1A*-related seizure disorders, which includes febrile seizures at the mild end to Dravet syndrome and intractable childhood epilepsy with generalized tonic-clonic seizures at the severe end. The spectrum may be associated with multiple seizure phenotypes, with a broad spectrum of severity; more severe seizure disorders may be associated with cognitive impairment, or deterioration. Ohtahara syndrome is a severe early-onset epilepsy syndrome characterized by intractable tonic spasms, other seizures, interictal

MEDICAL POLICY

POLICY TITLE	GENETIC TESTING FOR EPILEPSY
POLICY NUMBER	MP 2.262

electroencephalography abnormalities, and developmental delay. It may be secondary to structural abnormalities but has been associated with variants in the *STXBP1* gene in rare cases. West syndrome is an early-onset seizure disorder associated with infantile spasms and the characteristic electroencephalography finding of hypsarrhythmia. Other seizure disorders presenting early in childhood may have a genetic component but are characterized by a more benign course, including benign familial neonatal seizures and benign familial infantile seizures.

Genetic Etiology

Most genetic epilepsies are primarily believed to involve multifactorial inheritance patterns. This follows the concept of a threshold effect, in which any particular genetic defect may increase the risk of epilepsy but is not by itself causative. A combination of risk-associated genes, together with environmental factors, determines whether the clinical phenotype of epilepsy occurs. In this model, individual genes that increase the susceptibility to epilepsy have a relatively weak impact. Multiple genetic defects, and/or a particular combination of genes, probably increase the risk by a greater amount. However, it is not well understood how many abnormal genes are required to exceed the threshold to cause clinical epilepsy, nor is it understood which combination of genes may increase the risk more than others.

Early-onset epilepsy syndromes may be single-gene disorders. Because of the small amount of research available, the evidence base for these rare syndromes is incomplete, and new variants are currently being frequently discovered.

Some of the most common genes associated with genetic epileptic syndromes are listed in Table 2.

Table 2. Selected Genes Most Commonly Associated with Genetic Epilepsy

Genes	Physiologic Function
<i>KCNQ2</i>	Potassium channel
<i>KCNQ3</i>	Potassium channel
<i>SCN1A</i>	Sodium channel α -subunit
<i>SCN2A</i>	Sodium channel α -subunit
<i>SCN1B</i>	Sodium channel β -subunit
<i>GABRG2</i>	γ -aminobutyrate A-type subunit
<i>GABRA1</i>	γ -aminobutyrate A-type subunit
<i>GABRD</i>	γ -aminobutyrate subunit
<i>CHRNA2</i>	Acetylcholine receptor $\alpha 2$ subunit
<i>CHRNA4</i>	Acetylcholine receptor $\alpha 4$ subunit
<i>CHRNA2</i>	Acetylcholine receptor $\beta 2$ subunit
<i>STXBP1</i>	Synaptic vesicle release
<i>ARX</i>	Homeobox gene
<i>PCDH19</i>	Protocadherin cell-cell adhesion
<i>EFHC1</i>	Calcium homeostasis
<i>CACNB4</i>	Calcium channel subunit
<i>CLCN2</i>	Chloride channel
<i>LG11</i>	G-protein component

MEDICAL POLICY

POLICY TITLE	GENETIC TESTING FOR EPILEPSY
POLICY NUMBER	MP 2.262

Adapted from Williams and Battaglia (2013).

For the severe early epilepsy syndromes, the disorders most frequently reported to be associated with single-gene variants include generalized epilepsies with febrile seizures plus syndrome (associated with *SCN1A*, *SCN1B*, and *GABRG2* variants), Dravet syndrome (associated with *SCN1A* variants, possibly modified by *SCN9A* variants), and epilepsy and intellectual disability limited to females (associated with *PCDH19* variants). Ohtahara syndrome has been associated with variants in *STXBP1* in cases where patients have no structural or metabolic abnormalities. West syndrome is often associated with chromosomal abnormalities or tuberous sclerosis or may be secondary to an identifiable infectious or metabolic cause, but when there is no underlying cause identified, it is thought to be due to a multifactorial genetic predisposition.

Targeted testing for individual genes is available. Several commercial epilepsy genetic panels are also available. The number of genes included in the tests varies widely, from about 50 to over 450. The panels frequently include genes for other disorders such as neural tube defects, lysosomal storage disorders, cardiac channelopathies, congenital disorders of glycosylation, metabolic disorders, neurologic syndromes, and multisystemic genetic syndromes. Some panels are designed to be comprehensive while other panels target specific subtypes of epilepsy. Chambers et al (2016) reviewed comprehensive epilepsy panels from 7 U.S.-based clinical laboratories and found that between 1% and 4% of panel contents were genes not known to be associated with primary epilepsy. Between 1% and 70% of the genes included on an individual panel were not on any other panel.

Treatment

The condition is generally chronic, requiring treatment with one or more medications to adequately control symptoms. Seizures can be controlled by antiepileptic medications in most cases, but some patients are resistant to medications, and further options such as surgery, vagus nerve stimulation, and/or the ketogenic diet can be used.

Pharmacogenomics

Another area of interest for epilepsy is the pharmacogenomics of antiepileptic medications. There are a wide variety of these medications, from numerous different classes. The choice of medications and the combinations of medications for patients who require treatment with more than 1 agent is complex. Approximately one-third of patients are considered refractory to medications, defined as inadequate control of symptoms with a single medication. These patients often require escalating doses and/or combinations of different medications. At present, selection of agents is driven by the clinical phenotype of seizures but has a large trial-and-error component in many refractory cases. The current focus of epilepsy pharmacogenomics is in detecting genetic markers that identify patients likely to be refractory to the most common medications. This may lead to directed treatment that will result in a more efficient process for medication selection, and potentially more effective control of symptoms.

Of note, genotyping for the *HLA-B*1502* allelic variant in patients of Asian ancestry, prior to considering drug treatment with carbamazepine due to risks of severe dermatologic reactions, is recommended by the U.S. Food and Drug Administration labeling for carbamazepine.

MEDICAL POLICY

POLICY TITLE	GENETIC TESTING FOR EPILEPSY
POLICY NUMBER	MP 2.262

Regulatory Status

Clinical laboratories may develop and validate tests in-house and market them as a laboratory service; laboratory-developed tests must meet the general regulatory standards of the Clinical Laboratory Improvement Amendments. Commercially available genetic tests for epilepsy are available under the auspices of the Clinical Laboratory Improvement Amendments. Laboratories that offer laboratory-developed tests must be licensed by the Clinical Laboratory Improvement Amendments for high-complexity testing. To date, the U.S. Food and Drug Administration has chosen not to require any regulatory review of this test.

IV. RATIONALE

[TOP](#)

SUMMARY OF EVIDENCE

For individuals who have infantile- or early-childhood-onset epileptic encephalopathy who receive testing for genes associated with epileptic encephalopathies, the evidence includes a systematic review and meta-analysis, prospective, and retrospective cohort studies describing the testing yield. Relevant outcomes are test accuracy and validity, symptoms, quality of life, functional outcomes, medication use, resource utilization, and treatment-related morbidity. For Dravet syndrome, which appears to have the largest body of associated literature, the sensitivity of testing for *SCN1A* disease-associated variants is high (~80%). For other early-onset epileptic encephalopathies, the true clinical sensitivity and specificity of testing are not well-defined. However, studies reporting on the overall testing yield in populations with epileptic encephalopathies and early-onset epilepsy have reported detection rates for clinically significant variants ranging from 7.5% to 57%. The clinical utility of genetic testing occurs primarily when there is a positive test for a known pathogenic variant. The presence of a pathogenic variant may lead to targeted medication management, avoidance of other diagnostic tests, and/or informed reproductive planning. The evidence is sufficient to determine that the technology results in a meaningful improvement in the net health outcome.

For individuals who have presumed genetic epilepsy who receive testing for genetic variants associated with genetic epilepsies, the evidence includes prospective and retrospective cohort studies describing testing yields. Relevant outcomes are test accuracy and validity, changes in reproductive decision making, symptoms, quality of life, functional outcomes, medication use, resource utilization, and treatment-related morbidity. For most genetic epilepsies, which are thought to have a complex, multifactorial basis, the association between specific genetic variants and the risk of epilepsy is uncertain. Despite a large body of literature on associations between genetic variants and epilepsies, the clinical validity of genetic testing is poorly understood. Published literature is characterized by weak and inconsistent associations, which have not been replicated independently or by meta-analyses. A number of studies have also reported associations between genetic variants and AED treatment response, AED adverse effect risk, epilepsy phenotype, and risk of sudden unexplained death in epilepsy. The largest number of these studies is related to AED pharmacogenomics, which has generally reported some association between variants in a number of genes (including *SCN1A*, *SCN2A*, *ABCC2*, *EPHX1*, *CYP2C9*, *CYP2C19*), and AED response. Similarly, genetic associations between a number of genes and AED-related adverse events have been reported. However, no empirical evidence on the clinical utility of testing for the genetic epilepsies was identified, and the

MEDICAL POLICY

POLICY TITLE	GENETIC TESTING FOR EPILEPSY
POLICY NUMBER	MP 2.262

changes in clinical management that might occur as a result of testing are not well defined. The evidence is insufficient to determine the effects of the technology on health outcomes.

V. DEFINITIONS

[TOP](#)

N/A

VI. DISCLAIMER

[TOP](#)

Capital Blue Cross' medical policies are used to determine coverage for specific medical technologies, procedures, equipment, and services. These medical policies do not constitute medical advice and are subject to change as required by law or applicable clinical evidence from independent treatment guidelines. Treating providers are solely responsible for medical advice and treatment of members. These policies are not a guarantee of coverage or payment. Payment of claims is subject to a determination regarding the member's benefit program and eligibility on the date of service, and a determination that the services are medically necessary and appropriate. Final processing of a claim is based upon the terms of contract that applies to the members' benefit program, including benefit limitations and exclusions. If a provider or a member has a question concerning this medical policy, please contact Capital Blue Cross' Provider Services or Member Services.

VII. CODING INFORMATION

[TOP](#)

Note: This list of codes may not be all-inclusive, and codes are subject to change at any time. The identification of a code in this section does not denote coverage as coverage is determined by the terms of member benefit information. In addition, not all covered services are eligible for separate reimbursement.

Covered when medically necessary, genetic testing for epilepsy:

Procedure Codes								
0231U	0232U	81401	81403	81404	81405	81406	81407	81419
81479								

ICD-10-CM Diagnosis Codes	Description
G40.001	Localization-related (focal) (partial) idiopathic epilepsy and epileptic syndromes with seizures of localized onset, not intractable, with status epilepticus
G40.009	Localization-related (focal) (partial) idiopathic epilepsy and epileptic syndromes with seizures of localized onset, not intractable, without status epilepticus
G40.011	Localization-related (focal) (partial) idiopathic epilepsy and epileptic syndromes with seizures of localized onset, intractable, with status epilepticus
G40.019	Localization-related (focal) (partial) idiopathic epilepsy and epileptic syndromes with seizures of localized onset, intractable, without status epilepticus

MEDICAL POLICY

POLICY TITLE	GENETIC TESTING FOR EPILEPSY
POLICY NUMBER	MP 2.262

ICD-10-CM Diagnosis Codes	Description
G40.101	Localization-related (focal) (partial) symptomatic epilepsy and epileptic syndromes with simple partial seizures, not intractable, with status epilepticus
G40.109	Localization-related (focal) (partial) symptomatic epilepsy and epileptic syndromes with simple partial seizures, not intractable, without status epilepticus
G40.111	Localization-related (focal) (partial) symptomatic epilepsy and epileptic syndromes with simple partial seizures, intractable, with status epilepticus
G40.119	Localization-related (focal) (partial) symptomatic epilepsy and epileptic syndromes with simple partial seizures, intractable, without status epilepticus
G40.201	Localization-related (focal) (partial) symptomatic epilepsy and epileptic syndromes with complex partial seizures, not intractable, with status epilepticus
G40.209	Localization-related (focal) (partial) symptomatic epilepsy and epileptic syndromes with complex partial seizures, not intractable, without status epilepticus
G40.211	Localization-related (focal) (partial) symptomatic epilepsy and epileptic syndromes with complex partial seizures, intractable, with status epilepticus
G40.219	Localization-related (focal) (partial) symptomatic epilepsy and epileptic syndromes with complex partial seizures, intractable, without status epilepticus
G40.301	Generalized idiopathic epilepsy and epileptic syndromes, not intractable, with status epilepticus
G40.309	Generalized idiopathic epilepsy and epileptic syndromes, not intractable, without status epilepticus
G40.311	Generalized idiopathic epilepsy and epileptic syndromes, intractable, with status epilepticus
G40.319	Generalized idiopathic epilepsy and epileptic syndromes, intractable, without status epilepticus
G40.401	Other generalized epilepsy and epileptic syndromes, not intractable, with status epilepticus
G40.409	Other generalized epilepsy and epileptic syndromes, not intractable, without status epilepticus
G40.411	Other generalized epilepsy and epileptic syndromes, intractable, with status epilepticus
G40.419	Other generalized epilepsy and epileptic syndromes, intractable, without status epilepticus
G40.801	Other epilepsy, not intractable, with status epilepticus
G40.802	Other epilepsy, not intractable, without status epilepticus
G40.803	Other epilepsy, intractable, with status epilepticus
G40.804	Other epilepsy, intractable, without status epilepticus
G40.811	Lennox-Gastaut syndrome, not intractable, with status epilepticus
G40.812	Lennox-Gastaut syndrome, not intractable, without status epilepticus
G40.813	Lennox-Gastaut syndrome, intractable, with status epilepticus
G40.814	Lennox-Gastaut syndrome, intractable, without status epilepticus

MEDICAL POLICY

POLICY TITLE	GENETIC TESTING FOR EPILEPSY
POLICY NUMBER	MP 2.262

ICD-10-CM Diagnosis Codes	Description
G40.821	Epileptic spasms, not intractable, with status epilepticus
G40.822	Epileptic spasms, not intractable, without status epilepticus
G40.823	Epileptic spasms, intractable, with status epilepticus
G40.824	Epileptic spasms, intractable, without status epilepticus
G40.833	Dravet Syndrome, intractable with status epilepticus
G40.834	Dravet Syndrome, intractable without status epilepticus
G40.841	KCNQ2-related epilepsy, not intractable, with status epilepticus
G40.842	KCNQ2-related epilepsy, not intractable, without status epilepticus
G40.843	KCNQ2-related epilepsy, intractable, with status epilepticus
G40.844	KCNQ2-related epilepsy, intractable, without status epilepticus
G40.89	Other seizures
G40.901	Epilepsy, unspecified, not intractable, with status epilepticus
G40.909	Epilepsy, unspecified, not intractable, without status epilepticus
G40.911	Epilepsy, unspecified, intractable, with status epilepticus
G40.919	Epilepsy, unspecified, intractable, without status epilepticus
G40.A01	Absence epileptic syndrome, not intractable, with status epilepticus
G40.A09	Absence epileptic syndrome, not intractable, without status epilepticus
G40.A11	Absence epileptic syndrome, intractable, with status epilepticus
G40.A19	Absence epileptic syndrome, intractable, without status epilepticus
G40.B01	Juvenile myoclonic epilepsy, not intractable, with status epilepticus
G40.B09	Juvenile myoclonic epilepsy, not intractable, without status epilepticus
G40.B11	Juvenile myoclonic epilepsy, intractable, with status epilepticus
G40.B19	Juvenile myoclonic epilepsy, intractable, without status epilepticus
G40.C01	Lafora progressive myoclonus epilepsy, not intractable, with status epilepticus
G40.C09	Lafora progressive myoclonus epilepsy, not intractable, without status epilepticus
G40.C11	Lafora progressive myoclonus epilepsy, intractable, with status epilepticus
G40.C19	Lafora progressive myoclonus epilepsy, intractable, without status epilepticus

VIII. REFERENCES

[TOP](#)

1. Williams CA, Battaglia A. Molecular biology of epilepsy genes. *Exp Neurol*. Jun 2013; 244: 51-8. PMID 22178301
2. Fisher RS, Cross JH, French JA, et al. Operational classification of seizure types by the International League Against Epilepsy: Position Paper of the ILAE Commission for Classification and Terminology. *Epilepsia*. Apr 2017; 58(4): 522-530. PMID 28276060

MEDICAL POLICY

POLICY TITLE	GENETIC TESTING FOR EPILEPSY
POLICY NUMBER	MP 2.262

3. Berg AT, Berkovic SF, Brodie MJ, et al. Revised terminology and concepts for organization of seizures and epilepsies: report of the ILAE Commission on Classification and Terminology, 2005-2009. *Epilepsia*. Apr 2010; 51(4): 676-85. PMID 20196795
4. Merwick A, O'Brien M, Delanty N. Complex single gene disorders and epilepsy. *Epilepsia*. Sep 2012; 53 Suppl 4: 81-91. PMID 22946725
5. Miller IO, Sotero de Menezes MA. SCN1A-Related Seizure Disorders. In: Adam MP, Ardinger HH, Pagon RA, et al., eds. *GeneReviews*. Seattle, WA: University of Washington; 2014.
6. Petrovski S, Kwan P. Unraveling the genetics of common epilepsies: approaches, platforms, and caveats. *Epilepsy Behav*. Mar 2013; 26(3): 229-33. PMID 23103323
7. Helbig I, Lowenstein DH. Genetics of the epilepsies: where are we and where are we going?. *Curr Opin Neurol*. Apr 2013; 26(2): 179-85. PMID 23429546
8. Deprez L, Jansen A, De Jonghe P. Genetics of epilepsy syndromes starting in the first year of life. *Neurology*. Jan 20 2009; 72(3): 273-81. PMID 19153375
9. Chambers C, Jansen LA, Dhamija R. Review of Commercially Available Epilepsy Genetic Panels. *J Genet Couns*. Apr 2016; 25(2): 213-7. PMID 26536886
10. Kwan P, Brodie MJ. Early identification of refractory epilepsy. *N Engl J Med*. Feb 03 2000; 342(5): 314-9. PMID 10660394
11. Cavalleri GL, McCormack M, Alhusaini S, et al. Pharmacogenomics and epilepsy: the road ahead. *Pharmacogenomics*. Oct 2011; 12(10): 1429-47. PMID 22008048
12. Food and Drug Administration (FDA). Label: Tegritol. 2023; <https://www.accessdata.fda.gov/scripts/cder/daf/index.cfm?event=overview.process&ApplNo=016608>. Accessed December 18, 2023.
13. Food and Drug Administration (FDA). Depakene (valproic acid) Capsules and Oral Solution, Depakote (divalproex sodium) Delayed Release and Depakote ER (Extended Release) Tablets, Depakote Sprinkle Capsules (divalproex sodium coated particles in capsules), Depacon (valproate sodium) Injection. Safety Labeling Changes Approved By FDA Center for Drug Evaluation and Research (CDER) 2015; https://www.accessdata.fda.gov/drugsatfda_docs/applletter/2016/018081Orig1s064,018082Orig1s047,018723Orig1s056,019680Orig1s043,020593Orig1s034,021168Orig1s033ltr.pdf. Accessed December 18, 2023.
14. Dymant DA, Tetreault M, Beaulieu CL, et al. Whole-exome sequencing broadens the phenotypic spectrum of rare pediatric epilepsy: a retrospective study. *Clin Genet*. Jul 2015; 88(1): 34-40. PMID 25046240
15. Thevenon J, Milh M, Feillet F, et al. Mutations in SLC13A5 cause autosomal-recessive epileptic encephalopathy with seizure onset in the first days of life. *Am J Hum Genet*. Jul 03 2014; 95(1): 113-20. PMID 24995870
16. National Center for Biotechnology Information. GTR: Genetic Testing Registry. n.d.; <https://www.ncbi.nlm.nih.gov/gtr/>. Accessed December 18, 2023.
17. Hirose S, Scheffer IE, Marini C, et al. SCN1A testing for epilepsy: application in clinical practice. *Epilepsia*. May 2013; 54(5): 946-52. PMID 23586701
18. Mulley JC, Nelson P, Guerrero S, et al. A new molecular mechanism for severe myoclonic epilepsy of infancy: exonic deletions in SCN1A. *Neurology*. Sep 26 2006; 67(6): 1094-5. PMID 17000989

MEDICAL POLICY

POLICY TITLE	GENETIC TESTING FOR EPILEPSY
POLICY NUMBER	MP 2.262

19. Wu YW, Sullivan J, McDaniel SS, et al. Incidence of Dravet Syndrome in a US Population. *Pediatrics*. Nov 2015; 136(5): e1310-5. PMID 26438699
20. Esterhuizen AI, Mefford HC, Ramesar RS, et al. Dravet syndrome in South African infants: Tools for an early diagnosis. *Seizure*. Nov 2018; 62: 99-105. PMID 30321769
21. Peng J, Pang N, Wang Y, et al. Next-generation sequencing improves treatment efficacy and reduces hospitalization in children with drug-resistant epilepsy. *CNS Neurosci Ther*. Jan 2019; 25(1): 14-20. PMID 29933521
22. Scheffer IE, Bennett CA, Gill D, et al. Exome sequencing for patients with developmental and epileptic encephalopathies in clinical practice. *Dev Med Child Neurol*. Jan 2023; 65(1): 50-57. PMID 35701389
23. Jiang T, Gao J, Jiang L, et al. Application of Trio-Whole Exome Sequencing in Genetic Diagnosis and Therapy in Chinese Children With Epilepsy. *Front Mol Neurosci*. 2021; 14: 699574. PMID 34489640
24. Kim SY, Jang SS, Kim H, et al. Genetic diagnosis of infantile-onset epilepsy in the clinic: Application of whole-exome sequencing following epilepsy gene panel testing. *Clin Genet*. Mar 2021; 99(3): 418-424. PMID 33349918
25. Palmer EE, Sachdev R, Macintosh R, et al. Diagnostic Yield of Whole Genome Sequencing After Nondiagnostic Exome Sequencing or Gene Panel in Developmental and Epileptic Encephalopathies. *Neurology*. Mar 30 2021; 96(13): e1770-e1782. PMID 33568551
26. Salinas V, Martínez N, Maturo JP, et al. Clinical next generation sequencing in developmental and epileptic encephalopathies: Diagnostic relevance of data re-analysis and variants re-interpretation. *Eur J Med Genet*. Dec 2021; 64(12): 104363. PMID 34673242
27. Sun D, Liu Y, Cai W, et al. Detection of Disease-Causing SNVs/Indels and CNVs in Single Test Based on Whole Exome Sequencing: A Retrospective Case Study in Epileptic Encephalopathies. *Front Pediatr*. 2021; 9: 635703. PMID 34055682
28. Lee J, Lee C, Park WY, et al. Genetic Diagnosis of Dravet Syndrome Using Next Generation Sequencing-Based Epilepsy Gene Panel Testing. *Ann Clin Lab Sci*. Sep 2020; 50(5): 625-637. PMID 33067208
29. Lee S, Karp N, Zapata-Aldana E, et al. Genetic Testing in Children with Epilepsy: Report of a Single-Center Experience. *Can J Neurol Sci*. Mar 2021; 48(2): 233-244. PMID 32741404
30. Lee J, Lee C, Ki CS, et al. Determining the best candidates for next-generation sequencing-based gene panel for evaluation of early-onset epilepsy. *Mol Genet Genomic Med*. Sep 2020; 8(9): e1376. PMID 32613771
31. Stödberg T, Tomson T, Barbaro M, et al. Epilepsy syndromes, etiologies, and the use of next-generation sequencing in epilepsy presenting in the first 2 years of life: A population-based study. *Epilepsia*. Nov 2020; 61(11): 2486-2499. PMID 32964447
32. Berg AT, Coryell J, Saneto RP, et al. Early-Life Epilepsies and the Emerging Role of Genetic Testing. *JAMA Pediatr*. Sep 01 2017; 171(9): 863-871. PMID 28759667
33. Møller RS, Larsen LH, Johannesen KM, et al. Gene Panel Testing in Epileptic Encephalopathies and Familial Epilepsies. *Mol Syndromol*. Sep 2016; 7(4): 210-219. PMID 27781031

MEDICAL POLICY

POLICY TITLE	GENETIC TESTING FOR EPILEPSY
POLICY NUMBER	MP 2.262

34. Trump N, McTague A, Brittain H, et al. Improving diagnosis and broadening the phenotypes in early-onset seizure and severe developmental delay disorders through gene panel analysis. *J Med Genet.* May 2016; 53(5): 310-7. PMID 26993267
35. Wirrell EC, Shellhaas RA, Joshi C, et al. How should children with West syndrome be efficiently and accurately investigated? Results from the National Infantile Spasms Consortium. *Epilepsia.* Apr 2015; 56(4): 617-25. PMID 25779538
36. Mercimek-Mahmutoglu S, Patel J, Cordeiro D, et al. Diagnostic yield of genetic testing in epileptic encephalopathy in childhood. *Epilepsia.* May 2015; 56(5): 707-16. PMID 25818041
37. Hrabik SA, Standridge SM, Greiner HM, et al. The Clinical Utility of a Single-Nucleotide Polymorphism Microarray in Patients With Epilepsy at a Tertiary Medical Center. *J Child Neurol.* Nov 2015; 30(13): 1770-7. PMID 25862739
38. Ottman R, Hirose S, Jain S, et al. Genetic testing in the epilepsies--report of the ILAE Genetics Commission. *Epilepsia.* Apr 2010; 51(4): 655-70. PMID 20100225
39. Go CY, Mackay MT, Weiss SK, et al. Evidence-based guideline update: medical treatment of infantile spasms. Report of the Guideline Development Subcommittee of the American Academy of Neurology and the Practice Committee of the Child Neurology Society. *Neurology.* Jun 12 2012; 78(24): 1974-80. PMID 22689735
40. Pellock JM, Hrachovy R, Shinnar S, et al. Infantile spasms: a U.S. consensus report. *Epilepsia.* Oct 2010; 51(10): 2175-89. PMID 20608959
41. Wilmshurst JM, Gaillard WD, Vinayan KP, et al. Summary of recommendations for the management of infantile seizures: Task Force Report for the ILAE Commission of Pediatrics. *Epilepsia.* Aug 2015; 56(8): 1185-97. PMID 26122601
42. National Institute for Health and Care Excellence. Epilepsies: diagnosis and management [CG137]. 2021; <https://www.nice.org.uk/guidance/cg137>. Accessed December 18, 2023.
43. Ream MA, Mikati MA. Clinical utility of genetic testing in pediatric drug-resistant epilepsy: a pilot study. *Epilepsy Behav.* Aug 2014; 37: 241-8. PMID 25108116
44. Hoelz H, Herdl C, Gerstl L, et al. Impact on Clinical Decision Making of Next-Generation Sequencing in Pediatric Epilepsy in a Tertiary Epilepsy Referral Center. *Clin EEG Neurosci.* Jan 2020; 51(1): 61-69. PMID 31554424
45. National Institute of Neurological Disorders and Stroke (NINDS). NINDS Common Data Elements: Epilepsy. 2020, January; <https://www.commondataelements.ninds.nih.gov/epilepsy>. Accessed December 18, 2023.
46. McKnight D, Bristow SL, Truty RM, et al. Multigene Panel Testing in a Large Cohort of Adults With Epilepsy: Diagnostic Yield and Clinically Actionable Genetic Findings. *Neurol Genet.* Feb 2022; 8(1): e650. PMID 34926809
47. Alsubaie L, Aloraini T, Amoudi M, et al. Genomic testing and counseling: The contribution of next-generation sequencing to epilepsy genetics. *Ann Hum Genet.* Nov 2020; 84(6): 431-436. PMID 32533790
48. Johannesen KM, Nikanorova N, Marjanovic D, et al. Utility of genetic testing for therapeutic decision-making in adults with epilepsy. *Epilepsia.* Jun 2020; 61(6): 1234-1239. PMID 32427350

MEDICAL POLICY

POLICY TITLE	GENETIC TESTING FOR EPILEPSY
POLICY NUMBER	MP 2.262

49. Minardi R, Licchetta L, Baroni MC, et al. Whole-exome sequencing in adult patients with developmental and epileptic encephalopathy: It is never too late. *Clin Genet*. Nov 2020; 98(5): 477-485. PMID 32725632
50. Hesse AN, Bevilacqua J, Shankar K, et al. Retrospective genotype-phenotype analysis in a 305 patient cohort referred for testing of a targeted epilepsy panel. *Epilepsy Res*. Aug 2018; 144: 53-61. PMID 29778030
51. Lindy AS, Stosser MB, Butler E, et al. Diagnostic outcomes for genetic testing of 70 genes in 8565 patients with epilepsy and neurodevelopmental disorders. *Epilepsia*. May 2018; 59(5): 1062-1071. PMID 29655203
52. Miao P, Feng J, Guo Y, et al. Genotype and phenotype analysis using an epilepsy-associated gene panel in Chinese pediatric epilepsy patients. *Clin Genet*. Dec 2018; 94(6): 512-520. PMID 30182498
53. Butler KM, da Silva C, Alexander JJ, et al. Diagnostic Yield From 339 Epilepsy Patients Screened on a Clinical Gene Panel. *Pediatr Neurol*. Dec 2017; 77: 61-66. PMID 29056246
54. Tan NC, Berkovic SF. The Epilepsy Genetic Association Database (epiGAD): analysis of 165 genetic association studies, 1996-2008. *Epilepsia*. Apr 2010; 51(4): 686-9. PMID 20074235
55. Anney RJ, Avbersek A, Balding D, et al. Genetic determinants of common epilepsies: a meta-analysis of genome-wide association studies. *Lancet Neurol*. Sep 2014; 13(9): 893-903. PMID 25087078
56. Steffens M, Leu C, Ruppert AK, et al. Genome-wide association analysis of genetic generalized epilepsies implicates susceptibility loci at 1q43, 2p16.1, 2q22.3 and 17q21.32. *Hum Mol Genet*. Dec 15 2012; 21(24): 5359-72. PMID 22949513
57. Guo Y, Baum LW, Sham PC, et al. Two-stage genome-wide association study identifies variants in CAMSAP1L1 as susceptibility loci for epilepsy in Chinese. *Hum Mol Genet*. Mar 01 2012; 21(5): 1184-9. PMID 22116939
58. Córdoba M, Consalvo D, Moron DG, et al. SLC6A4 gene variants and temporal lobe epilepsy susceptibility: a meta-analysis. *Mol Biol Rep*. Dec 2012; 39(12): 10615-9. PMID 23065262
59. Nurmohamed L, Garcia-Bournissen F, Buono RJ, et al. Predisposition to epilepsy--does the ABCB1 gene play a role?. *Epilepsia*. Sep 2010; 51(9): 1882-5. PMID 20491876
60. Kauffman MA, Moron DG, Consalvo D, et al. Association study between interleukin 1 beta gene and epileptic disorders: a HuGe review and meta-analysis. *Genet Med*. Feb 2008; 10(2): 83-8. PMID 18281914
61. Tang L, Lu X, Tao Y, et al. SCN1A rs3812718 polymorphism and susceptibility to epilepsy with febrile seizures: a meta-analysis. *Gene*. Jan 01 2014; 533(1): 26-31. PMID 24076350
62. von Podewils F, Kowoll V, Schroeder W, et al. Predictive value of EFHC1 variants for the long-term seizure outcome in juvenile myoclonic epilepsy. *Epilepsy Behav*. Mar 2015; 44: 61-6. PMID 25625532
63. Kwan P, Poon WS, Ng HK, et al. Multidrug resistance in epilepsy and polymorphisms in the voltage-gated sodium channel genes SCN1A, SCN2A, and SCN3A: correlation among phenotype, genotype, and mRNA expression. *Pharmacogenet Genomics*. Nov 2008; 18(11): 989-98. PMID 18784617

MEDICAL POLICY

POLICY TITLE	GENETIC TESTING FOR EPILEPSY
POLICY NUMBER	MP 2.262

64. Jang SY, Kim MK, Lee KR, et al. Gene-to-gene interaction between sodium channel-related genes in determining the risk of antiepileptic drug resistance. *J Korean Med Sci.* Feb 2009; 24(1): 62-8. PMID 19270815
65. Lin CH, Chou IC, Hong SY. Genetic factors and the risk of drug-resistant epilepsy in young children with epilepsy and neurodevelopment disability: A prospective study and updated meta-analysis. *Medicine (Baltimore).* Mar 26 2021; 100(12): e25277. PMID 33761731
66. Li SX, Liu YY, Wang QB. ABCB1 gene C3435T polymorphism and drug resistance in epilepsy: evidence based on 8,604 subjects. *Med Sci Monit.* Mar 23 2015; 21: 861-8. PMID 25799371
67. Song C, Li X, Mao P, et al. Impact of CYP2C19 and CYP2C9 gene polymorphisms on sodium valproate plasma concentration in patients with epilepsy. *Eur J Hosp Pharm.* Jul 2022; 29(4): 198-201. PMID 32868386
68. Zhao T, Yu J, Wang TT, et al. Impact of ABCB1 Polymorphism on Levetiracetam Serum Concentrations in Epileptic Uyghur Children in China. *Ther Drug Monit.* Dec 2020; 42(6): 886-892. PMID 32890316
69. Lu Y, Fang Y, Wu X, et al. Effects of UGT1A9 genetic polymorphisms on monohydroxylated derivative of oxcarbazepine concentrations and oxcarbazepine monotherapeutic efficacy in Chinese patients with epilepsy. *Eur J Clin Pharmacol.* Mar 2017; 73(3): 307-315. PMID 27900402
70. Hashi S, Yano I, Shibata M, et al. Effect of CYP2C19 polymorphisms on the clinical outcome of low-dose clobazam therapy in Japanese patients with epilepsy. *Eur J Clin Pharmacol.* Jan 2015; 71(1): 51-8. PMID 25323806
71. Ma CL, Wu XY, Jiao Z, et al. SCN1A, ABCC2 and UGT2B7 gene polymorphisms in association with individualized oxcarbazepine therapy. *Pharmacogenomics.* 2015; 16(4): 347-60. PMID 25823783
72. Guo Y, Yan KP, Qu Q, et al. Common variants of KCNJ10 are associated with susceptibility and anti-epileptic drug resistance in Chinese genetic generalized epilepsies. *PLoS One.* 2015; 10(4): e0124896. PMID 25874548
73. Ma CL, Wu XY, Zheng J, et al. Association of SCN1A, SCN2A and ABCC2 gene polymorphisms with the response to antiepileptic drugs in Chinese Han patients with epilepsy. *Pharmacogenomics.* Jul 2014; 15(10): 1323-36. PMID 25155934
74. Rädisch S, Dickens D, Lang T, et al. A comprehensive functional and clinical analysis of ABCC2 and its impact on treatment response to carbamazepine. *Pharmacogenomics J.* Oct 2014; 14(5): 481-7. PMID 24567120
75. Yun W, Zhang F, Hu C, et al. Effects of EPHX1, SCN1A and CYP3A4 genetic polymorphisms on plasma carbamazepine concentrations and pharmacoresistance in Chinese patients with epilepsy. *Epilepsy Res.* Dec 2013; 107(3): 231-7. PMID 24125961
76. Taur SR, Kulkarni NB, Gandhe PP, et al. Association of polymorphisms of CYP2C9, CYP2C19, and ABCB1, and activity of P-glycoprotein with response to anti-epileptic drugs. *J Postgrad Med.* 2014; 60(3): 265-9. PMID 25121365
77. Haerian BS, Roslan H, Raymond AA, et al. ABCB1 C3435T polymorphism and the risk of resistance to antiepileptic drugs in epilepsy: a systematic review and meta-analysis. *Seizure.* Jul 2010; 19(6): 339-46. PMID 20605481

MEDICAL POLICY

POLICY TITLE	GENETIC TESTING FOR EPILEPSY
POLICY NUMBER	MP 2.262

78. Sun G, Sun X, Guan L. Association of MDR1 gene C3435T polymorphism with childhood intractable epilepsy: a meta-analysis. *J Neural Transm (Vienna)*. Jul 2014; 121(7): 717-24. PMID 24553780
79. Shazadi K, Petrovski S, Roten A, et al. Validation of a multigenic model to predict seizure control in newly treated epilepsy. *Epilepsy Res*. Dec 2014; 108(10): 1797-805. PMID 25282706
80. Chung WH, Chang WC, Lee YS, et al. Genetic variants associated with phenytoin-related severe cutaneous adverse reactions. *JAMA*. Aug 06 2014; 312(5): 525-34. PMID 25096692
81. He XJ, Jian LY, He XL, et al. Association of ABCB1, CYP3A4, EPHX1, FAS, SCN1A, MICA, and BAG6 polymorphisms with the risk of carbamazepine-induced Stevens-Johnson syndrome/toxic epidermal necrolysis in Chinese Han patients with epilepsy. *Epilepsia*. Aug 2014; 55(8): 1301-6. PMID 24861996
82. Wang W, Hu FY, Wu XT, et al. Genetic susceptibility to the cross-reactivity of aromatic antiepileptic drugs-induced cutaneous adverse reactions. *Epilepsy Res*. Aug 2014; 108(6): 1041-5. PMID 24856347
83. Bagnall RD, Crompton DE, Cutmore C, et al. Genetic analysis of PHOX2B in sudden unexpected death in epilepsy cases. *Neurology*. Sep 09 2014; 83(11): 1018-21. PMID 25085640
84. Coll M, Allegue C, Partemi S, et al. Genetic investigation of sudden unexpected death in epilepsy cohort by panel target resequencing. *Int J Legal Med*. Mar 2016; 130(2): 331-9. PMID 26423924
85. Bagnall RD, Crompton DE, Petrovski S, et al. Exome-based analysis of cardiac arrhythmia, respiratory control, and epilepsy genes in sudden unexpected death in epilepsy. *Ann Neurol*. Apr 2016; 79(4): 522-34. PMID 26704558
86. Rivello JJ, Ashwal S, Hirtz D, et al. Practice parameter: diagnostic assessment of the child with status epilepticus (an evidence-based review): report of the Quality Standards Subcommittee of the American Academy of Neurology and the Practice Committee of the Child Neurology Society. *Neurology*. Nov 14 2006; 67(9): 1542-50. PMID 17101884
87. Hirtz D, Ashwal S, Berg A, et al. Practice parameter: evaluating a first nonfebrile seizure in children: report of the quality standards subcommittee of the American Academy of Neurology, The Child Neurology Society, and The American Epilepsy Society. *Neurology*. Sep 12 2000; 55(5): 616-23. PMID 10980722
88. Smith L, Malinowski J, Ceulemans S, et al. Genetic testing and counseling for the unexplained epilepsies: An evidence-based practice guideline of the National Society of Genetic Counselors. *J Genet Couns*. Apr 2023; 32(2): 266-280. PMID 36281494
89. Burgunder JM, Finsterer J, Szolnoki Z, et al. EFNS guidelines on the molecular diagnosis of channelopathies, epilepsies, migraine, stroke, and dementias. *Eur J Neurol*. May 2010; 17(5): 641-8. PMID 20298421
90. Wirrell EC, Laux L, Donner E, et al. Optimizing the Diagnosis and Management of Dravet Syndrome: Recommendations From a North American Consensus Panel. *Pediatr Neurol*. Mar 2017; 68: 18-34.e3. PMID 28284397
91. Blue Cross Blue Shield Association Medical Policy Reference Manual. 2.04.109 Genetic Testing for Epilepsy. March 2024

MEDICAL POLICY

POLICY TITLE	GENETIC TESTING FOR EPILEPSY
POLICY NUMBER	MP 2.262

IX. POLICY HISTORY

[Top](#)

MP 2.262	03/04/2019 Consensus Review. No change to policy statements. References and Policy Guidelines updated.
	03/13/2020 Consensus Review. No change to policy statements. References updated; coding reviewed.
	09/01/2020 Administrative Update. Added ICD10 codes G40.83, G40.833, G40.834
	10/16/2020 Administrative Update. Added new codes 81419 (INV) and 0232U (MN with criteria) effective 01/01/2021.
	05/26/2021 Consensus Review. No change to policy statement. References and coding reviewed.
	06/21/2022 Consensus Review. References updated. 0231U added as MN (no change in coverage for code).
	01/10/2023 Consensus Review. No change to policy statement. References reviewed and updated.
	02/26/2024 Consensus Review. Policy guidelines, cross-references, background and references updated. 81419 moved to MN coding table. Updated ICD-10 codes.
	08/15/2024 Administrative Update. Added ICD-10 codes as part of New Code. Eff 10/01/2024.
	03/10/2025 Consensus Review. Updated policy guidelines, background, and references. No changes to coding.
	07/22/2025 Administrative Update. Removed Benefit Variations Section and updated Disclaimer.
	03/06/2026 Retirement Review. EviCore delegation.

[Top](#)

Health care benefit programs issued or administered by Capital Blue Cross and/or its subsidiaries, Capital Advantage Insurance Company®, Capital Advantage Assurance Company®, and Keystone Health Plan® Central. Independent licensees of the Blue Cross BlueShield Association. Communications issued by Capital Blue Cross in its capacity as administrator of programs and provider relations for all companies.