

MEDICAL POLICY

POLICY TITLE	MOLECULAR MARKERS IN FINE NEEDLE ASPIRATES OF THE THYROID
POLICY NUMBER	MP 2.275

CLINICAL BENEFIT	☐ MINIMIZE SAFETY RISK OR CONCERN. ☒ MINIMIZE HARMFUL OR INEFFECTIVE INTERVENTIONS. ☐ ASSURE APPROPRIATE LEVEL OF CARE. ☐ ASSURE APPROPRIATE DURATION OF SERVICE FOR INTERVENTIONS. ☒ ASSURE THAT RECOMMENDED MEDICAL PREREQUISITES HAVE BEEN MET. ☐ ASSURE APPROPRIATE SITE OF TREATMENT OR SERVICE.
Effective Date:	RETIRED 7/1/2026

POLICY

For individuals who have thyroid nodules without strong clinical or radiologic findings suggestive of malignancy in whom surgical decision making would be affected by test results the use of either of the following types of molecular marker testing or gene variant analysis in fine needle aspirates of thyroid nodules with indeterminate cytologic findings (i.e., Bethesda diagnostic category III [atypia/follicular lesion of undetermined significance] or Bethesda diagnostic category IV [follicular neoplasm/suspicion for a follicular neoplasm]) may be considered **medically necessary**:

- Afirma® Genomic Sequencing Classifier; or
- ThyroSeq® Genomic Classifier; or
- In lieu of the above tests:
 - BRAF V600E, RET/PTC, RAS, PAX8/PPAR

The use of any of the following types of molecular marker testing or gene variant analysis in fine needle aspirates of thyroid nodules with indeterminate findings (Bethesda diagnostic category III [atypia/follicular lesion of undetermined significance] or Bethesda diagnostic category IV [follicular neoplasm/suspicion for a follicular neoplasm]) or suspicious findings (Bethesda diagnostic category V [suspicious for malignancy]) to rule in malignancy to guide surgical planning for initial resection rather than a 2-stage surgical biopsy followed by definitive surgery may be considered **medically necessary**:

- ThyroSeq® Genomic Classifier; or
- ThyraMIR® microRNA/ThyGeNEXT®; or
- Afirma® Genomic Sequencing Classifier; or
- In lieu of the above tests:
 - BRAF V600E, RET/PTC, RAS, PAX8/PPAR

Gene expression classifiers, genetic variant analysis, and molecular marker testing in fine needle aspirates of the thyroid not meeting criteria outlined above, including but not limited to use of, ThyroSeq CRC, Thyroid GuidePx, and single-gene *TERT* testing are considered **investigational** as there is insufficient evidence to support a general conclusion concerning the health outcomes or benefits associated with this procedure.

MEDICAL POLICY

POLICY TITLE	MOLECULAR MARKERS IN FINE NEEDLE ASPIRATES OF THE THYROID
POLICY NUMBER	MP 2.275

*If coverage of a test is requested, but is not listed above, please refer to **MP 2.259 Expanded Molecular Panel Testing of Cancers to Identify Targeted Therapies** for additional guidance.*

Policy Guidelines

In individuals who do not undergo surgical biopsy or thyroidectomy on the basis of gene expression classifier or molecular marker results, regular active surveillance is indicated.

Use of molecular marker testing based on fine needle aspirate of a thyroid nodule to rule in malignancy prior to surgical biopsy may guide surgical planning, particularly factors such as choice of surgical facility provider to ensure that the capability is available to conduct a frozen section pathologic reading during surgical biopsy so that surgical approach may be adjusted accordingly in a single surgery.

GENETICS NOMENCLATURE UPDATE

The Human Genome Variation Society nomenclature is used to report information on variants found in DNA and serves as an international standard in DNA diagnostics. It is being implemented for genetic testing medical evidence review updates starting in 2017 (see Table PG1). The Society's nomenclature is recommended by the Human Variome Project, the Human Genome Organization, and by the Human Genome Variation Society itself. The American College of Medical Genetics and Genomics and the Association for Molecular Pathology standards and guidelines for interpretation of sequence variants represent expert opinion from both organizations, in addition to the College of American Pathologists. These recommendations primarily apply to genetic tests used in clinical laboratories, including genotyping, single genes, panels, exomes, and genomes. Table PG2 shows the recommended standard terminology-"pathogenic," "likely pathogenic," "uncertain significance," "likely benign," and "benign"-to describe variants identified that cause Mendelian disorders.

Table PG1. Nomenclature to Report on Variants Found in DNA

Previous	Updated	Definition
Mutation	Disease-associated variant	Disease-associated change in the DNA sequence
	Variant	Change in the DNA sequence
	Familial variant	Disease-associated variant identified in a proband for use in subsequent targeted genetic testing in first-degree relatives

Table PG2. ACMG-AMP Standards and Guidelines for Variant Classification

Variant Classification	Definition
Pathogenic	Disease-causing change in the DNA sequence
Likely pathogenic	Likely disease-causing change in the DNA sequence
Variant of uncertain significance	Change in DNA sequence with uncertain effects on disease
Likely benign	Likely benign change in the DNA sequence
Benign	Benign change in the DNA sequence

MEDICAL POLICY

POLICY TITLE	MOLECULAR MARKERS IN FINE NEEDLE ASPIRATES OF THE THYROID
POLICY NUMBER	MP 2.275

ACMG: American College of Medical Genetics and Genomics; AMP: Association for Molecular Pathology.

GENETIC COUNSELING

Experts recommend formal genetic counseling for patients who are at risk for inherited disorders and who wish to undergo genetic testing. Interpreting the results of genetic tests and understanding risk factors can be difficult for some patients; genetic counseling helps individuals understand the impact of genetic testing, including the possible effects the test results could have on the individual or their family members. It should be noted that genetic counseling may alter the utilization of genetic testing substantially and may reduce inappropriate testing; further, genetic counseling should be performed by an individual with experience and expertise in genetic medicine and genetic testing methods.

Cross-Reference:

MP 2.259 Molecular Panel Testing of Cancers to Identify Targeted Therapies

PRODUCT VARIATIONS

This policy is only applicable to certain programs and products administered by Capital Blue Cross and subject to benefit variations. Please see additional information below.

FEP PPO - Refer to FEP Medical Policy Manual. The FEP Medical Policy manual can be found at: <https://www.fepblue.org/benefit-plans/medical-policies-and-utilization-management-guidelines/medical-policies>

DESCRIPTION/BACKGROUND

THYROID NODULES

Thyroid nodules are common, present in 5% to 7% of the U.S. adult population; however, most are benign, and most cases of thyroid cancer are curable surgically when detected early.

Diagnosis

Sampling thyroid cells by fine needle aspirate (FNA) is currently the most accurate procedure to distinguish benign thyroid lesions from malignant ones, reducing the rate of unnecessary thyroid surgery for patients with benign nodules and triaging patients with thyroid cancer to appropriate surgery.

About 60% to 70% of thyroid nodules are classified cytologically as benign, and 4% to 10% of nodules are cytologically deemed malignant. However, the remaining 20% to 30% have equivocal findings, usually due to overlapping cytologic features between benign and malignant nodules; these nodules usually require surgery for a final diagnosis. Thyroid FNA cytology is classified by Bethesda System criteria into the following groups: nondiagnostic; benign; follicular lesion of undetermined significance (FLUS) or atypia of undetermined significance (AUS); follicular neoplasm (or suspicious for follicular neoplasm); suspicious for malignancy; and malignant. Lesions with FNA cytology in the atypia of undetermined significance or follicular

MEDICAL POLICY

POLICY TITLE	MOLECULAR MARKERS IN FINE NEEDLE ASPIRATES OF THE THYROID
POLICY NUMBER	MP 2.275

neoplasm of undetermined significance or follicular neoplasm categories are often considered indeterminate.

Management

There is some individualization of management for patients with FNA-indeterminate nodules, but many patients will require a surgical biopsy, typically thyroid lobectomy, with intraoperative pathology. Consultation would typically be the next step in diagnosis. Approximately 80% of patients with indeterminate cytology undergo surgical resection; postoperative evaluation has revealed a malignancy rate ranging from 6% to 30%, making this a clinical process with very low specificity. Thus, if analysis of FNA samples could reliably identify the risk of malignancy as low, there is potential for patients to avoid surgical biopsy.

Preoperative planning of optimal surgical management in patients with equivocal cytologic results is challenging, because different thyroid malignancies require different surgical procedures (e.g., unilateral lobectomy vs total or subtotal thyroidectomy with or without lymph node dissection) depending on several factors, including histologic subtype and risk-stratification strategies (tumor size, patient age). If a diagnosis cannot be made intraoperatively, a lobectomy is typically performed, and, if on postoperative histology the lesion is malignant, a second surgical intervention may be necessary for completion thyroidectomy.

THYROID CANCER

Most thyroid cancers originate from thyroid follicular cells and include well-differentiated papillary thyroid carcinoma (PTC; 80% of all thyroid cancers) and follicular carcinoma (15%). Poorly differentiated and anaplastic thyroid carcinomas are uncommon and can arise de novo or from preexisting well-differentiated papillary or follicular carcinomas. Medullary thyroid carcinoma originates from parafollicular or C cells and accounts for about 3% of all thyroid cancers.

The diagnosis of malignancy in the case of PTC is primarily based on cytologic features. If FNA in a case of PTC is indeterminate, surgical biopsy with intraoperative pathology consultation is most often diagnostic, although its efficacy and therefore its use will vary across institutions, surgeons, and pathologists. In 2016, reclassification of encapsulated follicular-variant PTC as a noninvasive follicular tumor with papillary-like nuclei was proposed and largely adopted; this classification removes the word *carcinoma* from the diagnosis to acknowledge the indolent behavior of these tumors.

For follicular carcinoma, the presence of invasion of the tumor capsule or blood vessels is diagnostic, and cannot be determined by cytology, because tissue sampling is necessary to observe these histologic characteristics. Intraoperative diagnosis of follicular carcinoma is challenging and often not feasible because extensive sampling of the tumor and capsule is usually necessary and performed on postoperative, permanent sections.

New approaches for improving the diagnostic accuracy of thyroid FNA include variant analysis for somatic genetic alterations, to more accurately classify which patients need to proceed to surgery (and may include the extent of surgery necessary), and a gene expression classifier to identify patients who do not need surgery and can be safely followed.

MEDICAL POLICY

POLICY TITLE	MOLECULAR MARKERS IN FINE NEEDLE ASPIRATES OF THE THYROID
POLICY NUMBER	MP 2.275

Genetic Variants Associated with Thyroid Cancer

A number of genetic variants have been discovered in thyroid cancer. The most common 4 gene variants are *BRAF* and *RAS* single nucleotide variants (SNVs) and *RET/PTC* and *PAX8/PPAR* rearrangements.

Papillary carcinomas carry SNVs of the *BRAF* and *RAS* genes, as well as *RET/PTC* and *TRK* rearrangements, all of which can activate the mitogen-activated protein kinase pathway. These mutually exclusive variants are found in more than 70% of papillary carcinomas. *BRAF* SNVs are highly specific for PTC. Follicular carcinomas harbor either *RAS* SNVs or *PAX8/PPAR* rearrangements. These variants have been identified in 70% to 75% of follicular carcinomas. Genetic alterations involving the PI3K/AKT signaling pathway also occur in thyroid tumors, although they are rare in well-differentiated thyroid cancers and have a higher prevalence in less differentiated thyroid carcinomas. Additional variants known to occur in poorly differentiated and anaplastic carcinomas involve the *TP53* and *CTNNB1* genes. Medullary carcinomas, which can be familial or sporadic, frequently possess SNVs located in the *RET* gene.

Studies have evaluated the association between various genes and cancer phenotype in individuals with diagnosed thyroid cancer.

Telomerase reverse transcriptase (*TERT*) promoter variants occur with varying frequency in different thyroid cancer subtypes. Overall, *TERT* C228T or C250T variants have been reported in approximately 15% of thyroid cancers, with higher rates in the undifferentiated and anaplastic subtypes compared with the well-differentiated subtypes. *TERT* variants are associated with several demographic and histopathologic features such as older age and advanced TNM stage. *TERT* promoter variants have been reported to be independent predictors of disease recurrence and cancer-related mortality in well-differentiated thyroid cancer. Also, the co-occurrence of *BRAF* or *RAS* variants with *TERT* or *TP53* variants may identify a subset of thyroid cancers with unfavorable outcomes.

Molecular Diagnostic Testing

Variant Detection and Rearrangement Testing

SNVs in specific genes, including *BRAF*, *RAS*, and *RET*, and evaluation for rearrangements associated with thyroid cancers can be accomplished with Sanger sequencing or pyrosequencing or with real-time polymerase chain reaction (PCR) of single or multiple genes or by next-generation sequencing (NGS) panels. Panel tests for genes associated with thyroid cancer, with varying compositions, are also available. For example, Quest Diagnostics offers a Thyroid Cancer Mutation Panel, which includes *BRAF* and *RAS* variant analysis and testing for *RET/PTC* and *PAX8/PPAR* rearrangements.

The ThyroSeq Genomic Classifier (Sonic Healthcare) is an NGS panel of 112 genes for four main classes of molecular alterations, including mutations, gene fusions, copy number alterations, and gene expression alterations. The results are processed by a unique genomic classifier (GC) and reported as positive or negative. ThyGeNEXT is an NGS panel that sequences 10 genes and identifies specific gene variants and translocations associated with

MEDICAL POLICY

POLICY TITLE	MOLECULAR MARKERS IN FINE NEEDLE ASPIRATES OF THE THYROID
POLICY NUMBER	MP 2.275

thyroid cancer. ThyGeNEXT is intended to be used in conjunction with the ThyraMIRv2 microRNA expression test when the initial ThyGeNEXT test is negative.

Gene Expression Profiling

Genetic alterations associated with thyroid cancer can be assessed using gene expression profiling, which refers to the analysis of messenger RNA (mRNA) expression levels of many genes simultaneously. Several gene expression profiling tests are available and stratify tissue from thyroid nodules biologically.

The Afirma Gene Expression Classifier (Afirma GSC; Veracyte) analyzes the expression of 142 different genes to determine patterns associated with benign findings on surgical biopsy. It is designed to evaluate thyroid nodules that have an “indeterminate” classification on FNA as a method to select patients (“rule out”) who are at low risk for cancer. In 2017, Veracyte migrated the Afirma GEC microarray analysis to a next-generation RNA sequencing platform and now markets the Afirma Gene Sequencing Classifier (Afirma GSC) which evaluates 10,196 genes with 1115 core genes.

Other gene expression profiles have been reported in investigational settings but have not been widely validated or used commercially (e.g., Barros-Filho et al [2015], Zheng et al [2015]); they are not addressed in this review.

ThyraMIRv2 is a microRNA expression–based classifier intended for use in thyroid nodules with indeterminate cytology on FNA following a negative result from the ThyGeNEXT Thyroid Oncogene Panel.

Algorithmic Testing

Algorithmic testing involves the use of 2 or more tests in a prespecified sequence, with a subsequent test automatically obtained depending on results of an earlier test.

Algorithmic Testing Using ThyGeNEXT and ThyraMIRv2

The ThyGeNEXT Thyroid Oncogene Panel (Interpace Diagnostics) is an NGS panel designed to assess patients with indeterminate thyroid FNA results. It includes sequencing of 10 genes and 38 RNA fusions. ThyGeNEXT has replaced the predicate ThyGenXThyroid test.

ThyraMIRv2 (Interpace Diagnostics) is a microRNA expression–based classifier intended for use in thyroid nodules with indeterminate cytology on FNA following a negative result from the ThyGeNEXT Thyroid Oncogene Panel.

The testing strategy for combined ThyGeNEXT and ThyraMIRv2 testing is first to predict malignancy. A positive result on ThyGeNEXT would “rule in” patients for surgical resection. The specific testing results from a ThyGeNEXT positive test would be used to inform preoperative planning when positive. For a ThyGeNEXT negative result, the reflex testing involves the ThyraMIRv2 microRNA expression test to “rule out” for a surgical biopsy procedure given the high negative predictive value of the second test. Patients with a negative result from the ThyraMIRv2 test would be followed with active surveillance and avoid a surgical biopsy.

Recurrence

MEDICAL POLICY

POLICY TITLE	MOLECULAR MARKERS IN FINE NEEDLE ASPIRATES OF THE THYROID
POLICY NUMBER	MP 2.275

ThyroSeq Cancer Risk Classifier (CRC) and Thyroid GuidePx are tests for recurrence. ThyroSeq CRC is to be used in FNAs from cytologically malignant (Bethesda VI) nodules and in resected thyroid cancers to predict risk of cancer recurrence. Thyroid GuidePx is used in papillary thyroid cancer and identifies a low-risk group with <5% recurrence rate.

REGULATORY STATUS

Clinical laboratories may develop and validate tests in-house and market them as a laboratory service; laboratory-developed tests must meet the general regulatory standards of the Clinical Laboratory Improvement Amendments. Thyroid variant testing and gene expression classifiers are available under the auspices of the Clinical Laboratory Improvement Amendments. Laboratories that offer laboratory-developed tests must be licensed by the Clinical Laboratory Improvement Amendments for high-complexity testing. To date, the U.S. Food and Drug Administration has chosen not to require any regulatory review of this test.

Table 2 provides a summary of commercially available molecular diagnostic tests for indeterminate thyroid pathology.

Table 2. Summary of Molecular Tests for Indeterminate Thyroid Cytopathology FNA Specimens

Test	Predicate	Methodology	Analyte(s)	Report
Afirma® GSC	Afirma® GEC	mRNA gene expression	1115 genes	Benign/suspicious
ThyroSeq GC	ThyroSeq	Next-generation sequencing	112 genes	Specific gene variant/translocation
ThyGeNEXT®	ThyGenX®, miRInform® ^a	Next-generation sequencing	10 genes and 38 gene fusions	Specific gene variant/translocation
ThyraMIR®v2	ThyraMIR®	microRNA expression	11 microRNAs	Negative/positive
RosettaGX™ Reveal		microRNA expression	24 microRNAs	☐ Benign ☐ Suspicious for malignancy ☐ High risk for medullary carcinoma

FNA: fine needle aspirate; NGS: next-generation sequencing; PCR: polymerase chain reaction.

^a The miRInform® test is the predicate test to ThyGenX® and is not commercially available.

RATIONALE

SUMMARY OF EVIDENCE

To determine which patients need thyroid resection, many physicians will perform a cytologic examination of FNA samples from a thyroid lesion; however, this method has diagnostic

MEDICAL POLICY

POLICY TITLE	MOLECULAR MARKERS IN FINE NEEDLE ASPIRATES OF THE THYROID
POLICY NUMBER	MP 2.275

limitations. As a result, assays using molecular markers have been developed to improve the accuracy of thyroid FNA biopsies.

For individuals with thyroid nodule(s) and indeterminate findings on FNA who receive FNA sample testing with molecular tests to rule out malignancy and to avoid surgical biopsy or resection, the evidence includes a prospective clinical validity study with the Afirma GSC and a chain of evidence to support clinical utility. The relevant outcomes are disease-specific survival, test accuracy and validity, morbid events, and resource utilization. In a multicenter validation study, the Afirma GSC was reported to have a high (NPV 96%; 95% CI, 90%-99%). These results are consistent with an earlier study on the Afirma GEC in the same study population. In other multicenter and single-center studies, there is suggestive evidence that rates of malignancy are low in Afirma patients who are classified as benign, but the exact NPV is unknown. The available evidence suggests that the decisions a physician makes regarding surgery are altered by Afirma GEC/GSC results; however, it should be noted that long-term follow-up of patients with thyroid nodules who avoided surgery based on GEC results is limited. A chain of evidence can be constructed to establish the potential for clinical utility with GEC testing in cytologically indeterminate lesions, but there is only a single study of the marketed test reporting a true NPV. Clinical input, obtained in 2017, supported the use of the previous version of the Afirma test in FNA of thyroid nodules with indeterminate cytologic findings to rule out malignancy and avoid surgical biopsy with an acceptably low trade-off in missed malignancy. The evidence is sufficient to determine that the technology improves the net health outcome. For individuals with thyroid nodule(s) and indeterminate findings on FNA who receive FNA sample testing with molecular markers to rule in malignancy and to guide surgical planning, the evidence includes prospective and retrospective studies of clinical validity. Relevant outcomes are disease-specific survival, test accuracy and validity, morbid events, and resource utilization. Variant analysis has the potential to improve the accuracy of an equivocal FNA of the thyroid and may play a role in preoperative risk stratification and surgical planning. Single-center studies have suggested that testing for a panel of genetic variants associated with thyroid cancer may allow for the appropriate selection of patients for surgical management for the initial resection. Prospective studies in additional populations are needed to validate these results. Although the presence of certain variants may predict more aggressive malignancies, the management changes that would occur as a result of identifying higher risk tumors, are not well established. Clinical input, obtained in 2017, considered ThyraMIR microRNA/ThyGenX, Afirma BRAF after Afirma GEC, and Afirma MTC after Afirma GEC to provide a clinically meaningful improvement for patients with cytologic findings suspicious for malignancy to guide surgical planning for the initial resection. The evidence is sufficient to determine that the technology results in a meaningful improvement in the net health outcome.

For individuals with thyroid nodule(s) and indeterminate findings on FNA who receive FNA sample testing with molecular markers to rule out malignancy and avoid surgical biopsy or to rule in malignancy for surgical planning, the evidence includes multiple retrospective and prospective clinical validation studies for the ThyroSeq v2 or v3 test and 2 retrospective clinical validation studies that used a predicate test 17-variant panel (miRInform) test to the current ThyGenX and ThyraMIR. The relevant outcomes are disease-specific survival, test accuracy and validity, morbid events, and resource utilization. In a retrospective validation study on FNA samples, the 17-variant panel (miRInform) test and ThyraMIR had a sensitivity of 89%, and an

MEDICAL POLICY

POLICY TITLE	MOLECULAR MARKERS IN FINE NEEDLE ASPIRATES OF THE THYROID
POLICY NUMBER	MP 2.275

NPV of 94%. A prospective clinical validation study of ThyroSeq v3 reported an NPV of 97% and PPV of 68%. No studies were identified demonstrating the diagnostic characteristics of the marketed ThyGenX. No studies were identified demonstrating evidence of direct outcome improvements. A chain of evidence for the ThyroSeq v3 test and combined ThyGenX and ThyraMIR testing would rely on establishing clinical validity. Clinical input, obtained in 2017, considered ThyroSeq v2 to provide a clinically meaningful improvement for patients with indeterminate cytologic findings to rule out malignancy and avoid surgical biopsy and in patients with cytologic findings suspicious for malignancy to guide surgical planning for the initial resection. The evidence is sufficient to determine that the technology results in a meaningful improvement in the net health outcome.

DEFINITIONS

N/A

DISCLAIMER

Capital Blue Cross' medical policies are used to determine coverage for specific medical technologies, procedures, equipment, and services. These medical policies do not constitute medical advice and are subject to change as required by law or applicable clinical evidence from independent treatment guidelines. Treating providers are solely responsible for medical advice and treatment of members. These policies are not a guarantee of coverage or payment. Payment of claims is subject to a determination regarding the member's benefit program and eligibility on the date of service, and a determination that the services are medically necessary and appropriate. Final processing of a claim is based upon the terms of contract that applies to the members' benefit program, including benefit limitations and exclusions. If a provider or a member has a question concerning this medical policy, please contact Capital Blue Cross' Provider Services or Member Services.

CODING INFORMATION

Note: This list of codes may not be all-inclusive, and codes are subject to change at any time. The identification of a code in this section does not denote coverage as coverage is determined by the terms of member benefit information. In addition, not all covered services are eligible for separate reimbursement.

RosettaGX Reveal, Thyroseq CRC, Thyroid GuidePx and single-gene *TERT* testing are considered investigational:

Procedure Codes							
81345	81479*	0287U	0362U				

*Can be used to bill for RosettaGX Reveal which is INV

MEDICAL POLICY

POLICY TITLE	MOLECULAR MARKERS IN FINE NEEDLE ASPIRATES OF THE THYROID
POLICY NUMBER	MP 2.275

BRAF V600E, RET/PTC, RAS, PAX8/PPAR, Afirma Genomic Sequencing Classifier, ThyroSeq Genomic Classifier, or ThyraMIR® microRNA/ThyGeNEXT® are covered when medically necessary:

Procedure Codes							
0018U	0026U	0245U	81210	81275	81276	81311	81401
81403	81445	81479*	81546				

*Can be used to bill for RET/PTC Rearrangement which may be MN

ICD-10-CM Diagnosis Codes	Description
D44.0	Neoplasm of uncertain behavior of thyroid gland

REFERENCES

1. Adeniran AJ, Theoharis C, Hui P, et al. Reflex BRAF testing in thyroid fine-needle aspiration biopsy with equivocal and positive interpretation: a prospective study. *Thyroid*. Jul 2011; 21(7): 717-23. PMID 21568726
2. Chudova D, Wilde JJ, Wang ET, et al. Molecular classification of thyroid nodules using high-dimensionality genomic data. *J Clin Endocrinol Metab*. Dec 2010; 95(12): 5296-304. PMID 20826580
3. Nikiforov YE, Seethala RR, Tallini G, et al. Nomenclature Revision for Encapsulated Follicular Variant of Papillary Thyroid Carcinoma: A Paradigm Shift to Reduce Overtreatment of Indolent Tumors. *JAMA Oncol*. Aug 01 2016; 2(8): 1023-9. PMID 27078145
4. Nikiforov YE. Molecular diagnostics of thyroid tumors. *Arch Pathol Lab Med*. May 2011; 135(5): 569-77. PMID 21526955
5. Han PA, Kim HS, Cho S, et al. Association of BRAF V600E Mutation and MicroRNA Expression with Central Lymph Node Metastases in Papillary Thyroid Cancer: A Prospective Study from Four Endocrine Surgery Centers. *Thyroid*. Apr 2016; 26(4): 532-42. PMID 26950846
6. Yip L, Nikiforova MN, Yoo JY, et al. Tumor genotype determines phenotype and disease-related outcomes in thyroid cancer: a study of 1510 patients. *Ann Surg*. Sep 2015; 262(3): 519-25; discussion 524-5. PMID 26258321
7. Lin JD, Fu SS, Chen JY, et al. Clinical Manifestations and Gene Expression in Patients with Conventional Papillary Thyroid Carcinoma Carrying the BRAF(V600E) Mutation and BRAF Pseudogene. *Thyroid*. May 2016; 26(5): 691-704. PMID 26914762
8. Alzahrani AS, Alsaadi R, Murugan AK, et al. TERT Promoter Mutations in Thyroid Cancer. *Horm Cancer*. Jun 2016; 7(3): 165-77. PMID 26902827
9. Landa I, Ganly I, Chan TA, et al. Frequent somatic TERT promoter mutations in thyroid cancer: higher prevalence in advanced forms of the disease. *J Clin Endocrinol Metab*. Sep 2013; 98(9): E1562-6. PMID 23833040

MEDICAL POLICY

POLICY TITLE	MOLECULAR MARKERS IN FINE NEEDLE ASPIRATES OF THE THYROID
POLICY NUMBER	MP 2.275

10. Liu X, Bishop J, Shan Y, et al. Highly prevalent TERT promoter mutations in aggressive thyroid cancers. *Endocr Relat Cancer*. Aug 2013; 20(4): 603-10. PMID 23766237
11. Liu T, Wang N, Cao J, et al. The age- and shorter telomere-dependent TERT promoter mutation in follicular thyroid cell-derived carcinomas. *Oncogene*. Oct 16 2014; 33(42): 4978-84. PMID 24141777
12. Xing M, Liu R, Liu X, et al. BRAF V600E and TERT promoter mutations cooperatively identify the most aggressive papillary thyroid cancer with highest recurrence. *J Clin Oncol*. Sep 01 2014; 32(25): 2718-26. PMID 25024077
13. Song YS, Lim JA, Choi H, et al. Prognostic effects of TERT promoter mutations are enhanced by coexistence with BRAF or RAS mutations and strengthen the risk prediction by the ATA or TNM staging system in differentiated thyroid cancer patients. *Cancer*. May 01 2016; 122(9): 1370-9. PMID 26969876
14. Nikiforova MN, Wald AI, Roy S, et al. Targeted next-generation sequencing panel (ThyroSeq) for detection of mutations in thyroid cancer. *J Clin Endocrinol Metab*. Nov 2013; 98(11): E1852-60. PMID 23979959
15. Sonic Healthcare USA. Thyroseq. Thyroid Genomic Classifier. 2024. <https://www.thyroseq.com/physicians/>. Accessed June 13, 2024.
16. Barros-Filho MC, Marchi FA, Pinto CA, et al. High Diagnostic Accuracy Based on CLDN10, HMGA2, and LAMB3 Transcripts in Papillary Thyroid Carcinoma. *J Clin Endocrinol Metab*. Jun 2015; 100(6): E890-9. PMID 25867809
17. Zheng B, Liu J, Gu J, et al. A three-gene panel that distinguishes benign from malignant thyroid nodules. *Int J Cancer*. Apr 01 2015; 136(7): 1646-54. PMID 25175491
18. Diggans J, Kim SY, Hu Z, et al. Machine learning from concept to clinic: reliable detection of BRAF V600E DNA mutations in thyroid nodules using high-dimensional RNA expression data. *Pac Symp Biocomput*. 2015: 371-82. PMID 25592597
19. Tessler FN, Middleton WD, Grant EG, et al. ACR Thyroid Imaging, Reporting and Data System (TI-RADS): White Paper of the ACR TI-RADS Committee. *J Am Coll Radiol*. May 2017; 14(5): 587-595. PMID 28372962
20. Lee E, Terhaar S, McDaniel L, et al. Diagnostic performance of the second-generation molecular tests in the assessment of indeterminate thyroid nodules: A systematic review and meta-analysis. *Am J Otolaryngol*. 2022; 43(3): 103394. PMID 35241290
21. Nasr CE, Andrioli M, Endo M, et al. Real-World Performance of the Afirma Genomic Sequencing Classifier (GSC)-A Meta-analysis. *J Clin Endocrinol Metab*. May 17 2023; 108(6): 1526-1532. PMID 36470585
22. Patel KN, Angell TE, Babiarz J, et al. Performance of a Genomic Sequencing Classifier for the Preoperative Diagnosis of Cytologically Indeterminate Thyroid Nodules. *JAMA Surg*. Sep 01 2018; 153(9): 817-824. PMID 29799911
23. Alexander EK, Kennedy GC, Baloch ZW, et al. Preoperative diagnosis of benign thyroid nodules with indeterminate cytology. *N Engl J Med*. Aug 23 2012; 367(8): 705-15. PMID 22731672
24. Livhits MJ, Zhu CY, Kuo EJ, et al. Effectiveness of Molecular Testing Techniques for Diagnosis of Indeterminate Thyroid Nodules: A Randomized Clinical Trial. *JAMA Oncol*. Jan 01 2021; 7(1): 70-77. PMID 33300952

MEDICAL POLICY

POLICY TITLE	MOLECULAR MARKERS IN FINE NEEDLE ASPIRATES OF THE THYROID
POLICY NUMBER	MP 2.275

25. Kim NE, Raghunathan RS, Hughes EG, et al. Bethesda III and IV Thyroid Nodules Managed Nonoperatively After Molecular Testing With Afirma GSC or Thyroseq v3. *J Clin Endocrinol Metab.* Aug 18 2023; 108(9): e698-e703. PMID 36995878
26. Santhanam P, Khthir R, Gress T, et al. Gene expression classifier for the diagnosis of indeterminate thyroid nodules: a meta-analysis. *Med Oncol.* Feb 2016; 33(2): 14. PMID 26749587
27. Liu Y, Pan B, Xu L, et al. The Diagnostic Performance of Afirma Gene Expression Classifier for the Indeterminate Thyroid Nodules: A Meta-Analysis. *Biomed Res Int.* 2019; 2019: 7150527. PMID 31531363
28. Angell TE, Frates MC, Medici M, et al. Afirma Benign Thyroid Nodules Show Similar Growth to Cytologically Benign Nodules During Follow-Up. *J Clin Endocrinol Metab.* Nov 2015; 100(11): E1477-83. PMID 26353010
29. Sipos JA, Blevins TC, Shea HC, et al. LONG-TERM NONOPERATIVE RATE OF THYROID NODULES WITH BENIGN RESULTS ON THE AFIRMA GENE EXPRESSION CLASSIFIER. *Endocr Pract.* Jun 2016; 22(6): 666-72. PMID 26789352
30. Deaver KE, Haugen BR, Pozdeyev N, et al. Outcomes of Bethesda categories III and IV thyroid nodules over 5 years and performance of the Afirma gene expression classifier: A single-institution study. *Clin Endocrinol (Oxf).* Aug 2018; 89(2): 226-232. PMID 29791966
31. Valderrabano P, Hallanger-Johnson JE, Thapa R, et al. Comparison of Postmarketing Findings vs the Initial Clinical Validation Findings of a Thyroid Nodule Gene Expression Classifier: A Systematic Review and Meta-analysis. *JAMA Otolaryngol Head Neck Surg.* Sep 01 2019; 145(9): 783-792. PMID 31318389
32. Harrell RM, Eyerly-Webb SA, Golding AC, et al. STATISTICAL COMPARISON OF AFIRMA GSC AND AFIRMA GEC OUTCOMES IN A COMMUNITY ENDOCRINE SURGICAL PRACTICE: EARLY FINDINGS. *Endocr Pract.* Feb 2019; 25(2): 161-164. PMID 30383497
33. Polavarapu P, Fingeret A, Yuil-Valdes A, et al. Comparison of Afirma GEC and GSC to Nodules Without Molecular Testing in Cytologically Indeterminate Thyroid Nodules. *J Endocr Soc.* Nov 01 2021; 5(11): bvab148. PMID 34708178
34. Kandil E, Metz TA, Issa PP, et al. Diagnostic Performance of Afirma and Interpace Diagnostics Genetic Testing in Indeterminate Thyroid Nodules: A Single Center Study. *Cancers (Basel).* Mar 31 2023; 15(7). PMID 37046759
35. Duick DS, Klopner JP, Diggans JC, et al. The impact of benign gene expression classifier test results on the endocrinologist-patient decision to operate on patients with thyroid nodules with indeterminate fine-needle aspiration cytopathology. *Thyroid.* Oct 2012; 22(10): 996-1001. PMID 22873825
36. Aragon Han P, Olson MT, Fazeli R, et al. The impact of molecular testing on the surgical management of patients with thyroid nodules. *Ann Surg Oncol.* Jun 2014; 21(6): 1862-9. PMID 24522987
37. Noureldine SI, Olson MT, Agrawal N, et al. Effect of Gene Expression Classifier Molecular Testing on the Surgical Decision-Making Process for Patients With Thyroid Nodules. *JAMA Otolaryngol Head Neck Surg.* Dec 2015; 141(12): 1082-8. PMID 26606459
38. Abeykoon JP, Mueller L, Dong F, et al. The Effect of Implementing Gene Expression Classifier on Outcomes of Thyroid Nodules with Indeterminate Cytology. *Horm Cancer.* Aug 2016; 7(4): 272-8. PMID 27102883

MEDICAL POLICY

POLICY TITLE	MOLECULAR MARKERS IN FINE NEEDLE ASPIRATES OF THE THYROID
POLICY NUMBER	MP 2.275

39. Chaudhary S, Hou Y, Shen R, et al. Impact of the Afirma Gene Expression Classifier Result on the Surgical Management of Thyroid Nodules with Category III/IV Cytology and Its Correlation with Surgical Outcome. *Acta Cytol.* 2016; 60(3): 205-10. PMID 27344463
40. Cibas ES, Ali SZ. The Bethesda System for Reporting Thyroid Cytopathology. *Thyroid.* Nov 2009; 19(11): 1159-65. PMID 19888858
41. Lithwick-Yanai G, Dromi N, Shtabsky A, et al. Multicentre validation of a microRNA-based assay for diagnosing indeterminate thyroid nodules utilising fine needle aspirate smears. *J Clin Pathol.* Jun 2017; 70(6): 500-507. PMID 27798083
42. Walts AE, Sacks WL, Wu HH, et al. A retrospective analysis of the performance of the RosettaGX® Reveal™ thyroid miRNA and the Afirma Gene Expression Classifiers in a cohort of cytologically indeterminate thyroid nodules. *Diagn Cytopathol.* Nov 2018; 46(11): 901-907. PMID 30353692
43. Fnais N, Soobiah C, Al-Qahtani K, et al. Diagnostic value of fine needle aspiration BRAF(V600E) mutation analysis in papillary thyroid cancer: a systematic review and meta-analysis. *Hum Pathol.* Oct 2015; 46(10): 1443-54. PMID 26232865
44. Kloos RT, Monroe RJ, Traweek ST, et al. A Genomic Alternative to Identify Medullary Thyroid Cancer Preoperatively in Thyroid Nodules with Indeterminate Cytology. *Thyroid.* Jun 2016; 26(6): 785-93. PMID 26992356
45. Xing M, Clark D, Guan H, et al. BRAF mutation testing of thyroid fine-needle aspiration biopsy specimens for preoperative risk stratification in papillary thyroid cancer. *J Clin Oncol.* Jun 20 2009; 27(18): 2977-82. PMID 19414674
46. Yin DT, Yu K, Lu RQ, et al. Clinicopathological significance of TERT promoter mutation in papillary thyroid carcinomas: a systematic review and meta-analysis. *Clin Endocrinol (Oxf).* Aug 2016; 85(2): 299-305. PMID 26732020
47. Bernet V, Hupart KH, Parangi S, et al. AACE/ACE disease state commentary: molecular diagnostic testing of thyroid nodules with indeterminate cytopathology. *Endocr Pract.* Apr 2014; 20(4): 360-3. PMID 24727662
48. Yip L, Wharry LI, Armstrong MJ, et al. A clinical algorithm for fine-needle aspiration molecular testing effectively guides the appropriate extent of initial thyroidectomy. *Ann Surg.* Jul 2014; 260(1): 163-8. PMID 24901361
49. Ferris RL, Baloch Z, Bernet V, et al. American Thyroid Association Statement on Surgical Application of Molecular Profiling for Thyroid Nodules: Current Impact on Perioperative Decision Making. *Thyroid.* Jul 2015; 25(7): 760-8. PMID 26058403
50. Nikiforova MN, Mercurio S, Wald AI, et al. Analytical performance of the ThyroSeq v3 genomic classifier for cancer diagnosis in thyroid nodules. *Cancer.* Apr 15 2018; 124(8): 1682-1690. PMID 29345728
51. Steward DL, Carty SE, Sippel RS, et al. Performance of a Multigene Genomic Classifier in Thyroid Nodules With Indeterminate Cytology: A Prospective Blinded Multicenter Study. *JAMA Oncol.* Feb 01 2019; 5(2): 204-212. PMID 30419129
52. Valderrabano P, Khazai L, Leon ME, et al. Evaluation of ThyroSeq v2 performance in thyroid nodules with indeterminate cytology. *Endocr Relat Cancer.* Mar 2017; 24(3): 127-136. PMID 28104680
53. Taye A, Gurciullo D, Miles BA, et al. Clinical performance of a next-generation sequencing assay (ThyroSeq v2) in the evaluation of indeterminate thyroid nodules. *Surgery.* Jan 2018; 163(1): 97-103. PMID 29154079

MEDICAL POLICY

POLICY TITLE	MOLECULAR MARKERS IN FINE NEEDLE ASPIRATES OF THE THYROID
POLICY NUMBER	MP 2.275

54. Moses W, Weng J, Sansano I, et al. Molecular testing for somatic mutations improves the accuracy of thyroid fine-needle aspiration biopsy. *World J Surg.* Nov 2010; 34(11): 2589-94. PMID 20703476
55. Ohori NP, Nikiforova MN, Schoedel KE, et al. Contribution of molecular testing to thyroid fine-needle aspiration cytology of "follicular lesion of undetermined significance/atypia of undetermined significance". *Cancer Cytopathol.* Feb 25 2010; 118(1): 17-23. PMID 20099311
56. Beaudenon-Huibregtse S, Alexander EK, Guttler RB, et al. Centralized molecular testing for oncogenic gene mutations complements the local cytopathologic diagnosis of thyroid nodules. *Thyroid.* Oct 2014; 24(10): 1479-87. PMID 24811481
57. Li W, Justice-Clark T, Cohen MB. The utility of ThyroSeq ® in the management of indeterminate thyroid nodules by fine-needle aspiration. *Cytopathology.* Jul 2021; 32(4): 505-512. PMID 33914382
58. Labourier E, Shifrin A, Busseniers AE, et al. Molecular Testing for miRNA, mRNA, and DNA on Fine-Needle Aspiration Improves the Preoperative Diagnosis of Thyroid Nodules With Indeterminate Cytology. *J Clin Endocrinol Metab.* Jul 2015; 100(7): 2743-50. PMID 25965083
59. Gharib H, Papini E, Garber JR, et al. AMERICAN ASSOCIATION OF CLINICAL ENDOCRINOLOGISTS, AMERICAN COLLEGE OF ENDOCRINOLOGY, AND ASSOCIAZIONE MEDICI ENDOCRINOLOGI MEDICAL GUIDELINES FOR CLINICAL PRACTICE FOR THE DIAGNOSIS AND MANAGEMENT OF THYROID NODULES--2016 UPDATE. *Endocr Pract.* May 2016; 22(5): 622-39. PMID 27167915
60. Haugen BR, Alexander EK, Bible KC, et al. 2015 American Thyroid Association Management Guidelines for Adult Patients with Thyroid Nodules and Differentiated Thyroid Cancer: The American Thyroid Association Guidelines Task Force on Thyroid Nodules and Differentiated Thyroid Cancer. *Thyroid.* Jan 2016; 26(1): 1-133. PMID 26462967
61. National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology: Thyroid Carcinoma. Version 4.2024.
62. Ross DS. Evaluation and management of thyroid nodules with indeterminate cytology. In: *UpToDate Online Journal* [serial online]. Waltham, MA: UpToDate; updated August 29, 2024. Literature review current through August 2024.

POLICY HISTORY

MP 2.275	03/30/2020 Consensus Review. Nomenclature updated. Gene Expression updated to replace Afirma GEC to Afirma GSC throughout policy. No change to policy statement. References reviewed and updated.
	09/08/2020 Administrative Update. New codes added 0204U; 02085U
	12/11/2020 Administrative Update. Deleted code 81545. Effective 01/01/2021.
	02/01/2021 Administrative Update. Added new code 0245U as covered with criteria. Effective 04/01/2021.

MEDICAL POLICY

POLICY TITLE	MOLECULAR MARKERS IN FINE NEEDLE ASPIRATES OF THE THYROID
POLICY NUMBER	MP 2.275

	11/15/2021 Minor Review. Wording “in patients who have the following characteristics” removed from first paragraph in policy statement. No change to criteria. Description/Background updated. References reviewed and updated.
	10/19/2022 Minor Review. Updated investigational statement to include Thyroseq CRC Thyroid GuidePx. Reviewed and updated coding and references. Added NCCN statement.
	09/22/2023 Minor Review. Updated names of tests to current versions. Added option for non-proprietary testing for specific mutations (BRAF V600E, RET/PTC, RAS, PAX8/PPAR). Added note within policy statement regarding MP 2.259. Updated background, coding, and references.
	06/07/2024 Administrative Update. Removed 0204U as it has been deleted. Effective 07/01/2024.
	09/29/2024 Consensus Review. Updated clinical benefit, background and references. No changes to coding.
	08/15/2025 Consensus Review. RosettaGX test removed from policy as the test has been discontinued.
	03/04/2026 Retirement Review. Evicore Delegation.

Health care benefit programs issued or administered by Capital Blue Cross and/or its subsidiaries, Capital Advantage Insurance Company®, Capital Advantage Assurance Company® and Keystone Health Plan® Central. Independent licensees of the BlueCross BlueShield Association. Communications issued by Capital Blue Cross in its capacity as administrator of programs and provider relations for all companies.