

POLICY TITLE	ORTHOPEDIC APPLICATIONS OF PLATELET- RICH PLASMA
POLICY NUMBER	MP-4.039

Original Issue Date (Created):	2/1/2016
Most Recent Review Date (Revised):	8/19/2020
Effective Date:	12/1/2020

[POLICY RATIONALE](#)
[DISCLAIMER](#)
[POLICY HISTORY](#)

[PRODUCT VARIATIONS](#)
[DEFINITIONS](#)
[CODING INFORMATION](#)

[DESCRIPTION/BACKGROUND](#)
[BENEFIT VARIATIONS](#)
[REFERENCES](#)

I. POLICY

Use of platelet-rich plasma is considered **investigational** for all orthopedic indications. This includes, but is not limited to, use in the following situations:

- Primary use (injection) for the following conditions:
 - Achilles tendinopathy
 - Lateral epicondylitis
 - Plantar fasciitis
 - Osteochondral lesions
 - Osteoarthritis.
- Adjunctive use in the following surgical procedures:
 - Anterior cruciate ligament reconstruction
 - Hip fracture
 - Long-bone nonunion
 - Patellar tendon repair
 - Rotator cuff repair
 - Spinal fusion
 - Subacromial decompression surgery
 - Total knee arthroplasty.

Cross-reference:

MP-2.033 Recombinant and Autologous Platelet-Derived Growth Factors as Treatment of Wound Healing and Other Non-Orthopedic Conditions

MP-2.061 Prolotherapy

MP-2.080 Orthopedic Applications of Stem Cell Therapy (including Allograft and Bone Substitute Products Used with Autologous Bone Marrow)

POLICY TITLE	ORTHOPEDIC APPLICATIONS OF PLATELET- RICH PLASMA
POLICY NUMBER	MP-4.039

II. PRODUCT VARIATIONS

[TOP](#)

This policy is only applicable to certain programs and products administered by Capital BlueCross please see additional information below, and subject to benefit variations as discussed in Section VI below.

III. DESCRIPTION/BACKGROUND

[TOP](#)

A variety of growth factors have been found to play a role in wound healing, including platelet-derived growth factors (PDGFs), epidermal growth factor, fibroblast growth factors, transforming growth factors, and insulin-like growth factors. Autologous platelets are a rich source of PDGF, transforming growth factors that function as a mitogen for fibroblasts, smooth muscle cells, osteoblasts, and vascular endothelial growth factors. Recombinant PDGF has also been extensively investigated for clinical use in wound healing (see MP-2.033).

Autologous platelet concentrate suspended in plasma, also known as platelet-rich plasma (PRP), can be prepared from samples of centrifuged autologous blood. Exposure to a solution of thrombin and calcium chloride degranulates platelets, releasing the various growth factors. The polymerization of fibrin from fibrinogen creates a platelet gel, which can then be used as an adjunct to surgery with the intent of promoting hemostasis and accelerating healing. In the operating room setting, PRP has been investigated as an adjunct to a variety of periodontal, reconstructive, and orthopedic procedures. For example, bone morphogenetic proteins are a type of transforming growth factors, and thus PRP has been used in conjunction with bone-replacement grafting (using either autologous grafts or bovine-derived xenograft) in periodontal and maxillofacial surgeries. Alternatively, PRP may be injected directly into various tissues. PRP injections have been proposed as a primary treatment of miscellaneous conditions, such as epicondylitis, plantar fasciitis, and Dupuytren contracture.

Injection of PRP for tendon and ligament pain is theoretically related to prolotherapy (discussed in MP-2.061). However, prolotherapy differs in that it involves injection of chemical irritants that are intended to stimulate inflammatory responses and induce release of endogenous growth factors.

PRP is distinguished from fibrin glues or sealants, which have been used as a surgical adjunct to promote local hemostasis at incision sites. Fibrin glue is created from platelet-poor plasma and consists primarily of fibrinogen. Commercial fibrin glues are created from pooled homologous human donors; Tisseel® (Baxter) and Hemaseel® (Haemacure Corp) are examples of commercially available fibrin sealants. Autologous fibrin sealants can be created from platelet-poor plasma. This evidence review does not address the use of fibrin sealants.

REGULATORY STATUS

The U.S. Food and Drug Administration (FDA) regulates human cells and tissues intended for implantation, transplantation, or infusion through the Center for Biologics Evaluation and Research, under Code of Federal Regulation, title 21, parts 1270 and 1271. Blood products such

MEDICAL POLICY

POLICY TITLE	ORTHOPEDIC APPLICATIONS OF PLATELET- RICH PLASMA
POLICY NUMBER	MP-4.039

as PRP are included in these regulations. Under these regulations, certain products including blood products such as PRP are exempt and therefore do not follow the traditional FDA regulatory pathway. To date, the FDA has not attempted to regulate activated PRP.

A number of PRP preparation systems are available, many of which were cleared for marketing by the FDA through the 510(k) process for producing platelet-rich preparations intended to be mixed with bone graft materials to enhance the bone grafting properties in orthopedic practices. The use of PRP outside of this setting (eg, an office injection) would be considered off-label. The Aurix System™ (previously called AutoloGel™; Cytomedix) and SafeBlood® (SafeBlood Technologies) are two related but distinct autologous blood-derived preparations that can be used at the bedside for immediate application. Both AutoloGel™ and SafeBlood® have been specifically marketed for wound healing. Other devices may be used during surgery (eg, Medtronic Electromedics, Elmd-500 Autotransfusion system, the Plasma Saver device, the SmartPREPO [Harvest Technologies] device). The Magellan™ Autologous Platelet Separator System (Medtronic Sofamor Danek) includes a disposable kit for use with the Magellan™ Autologous Platelet Separator portable tabletop centrifuge. GPS®II (BioMet Biologics), a gravitational platelet separation system, was cleared for marketing by the FDA through the 510(k) process for use as disposable separation tube for centrifugation and a dual cannula tip to mix the platelets and thrombin at the surgical site. Filtration or plasmapheresis may also be used to produce platelet-rich concentrates. The use of different devices and procedures can lead to variable concentrations of activated platelets and associated proteins, increasing variability between studies of clinical efficacy.

IV. RATIONALE

[TOP](#)

SUMMARY OF EVIDENCE

Primary Treatment for Tendinopathies

For individuals with tendinopathy who receive PRP injections, the evidence includes multiple randomized controlled trials (RCTs) and systematic reviews with meta-analyses. Relevant outcomes are symptoms, functional outcomes, health status measures, quality of life, and treatment-related morbidity. Findings from meta-analyses of RCTs have been mixed and have generally found that PRP did not have a statistically and/or clinically significant impact on symptoms (ie, pain) or functional outcomes. The evidence is insufficient to determine the effects of the technology on health outcomes.

Primary Treatment for Non-Tendon Soft Tissue Injury or Inflammation

For individuals with non-tendon soft tissue injury or inflammation (eg, plantar fasciitis) who receive PRP injections, the evidence includes six small RCTs, multiple prospective observational studies, and a systematic review. Relevant outcomes are symptoms, functional outcomes, health status measures, quality of life, and treatment-related morbidity. The systematic review, which identified three RCTs on PRP for plantar fasciitis, did not pool study findings. Results among the six RCTs were inconsistent. The largest RCT showed that treatment using PRP compared with corticosteroid injection resulted in statistically significant improvement in pain and disability, but

MEDICAL POLICY

POLICY TITLE	ORTHOPEDIC APPLICATIONS OF PLATELET- RICH PLASMA
POLICY NUMBER	MP-4.039

not quality of life. Larger RCTs are still needed to address important uncertainties in efficacy and safety. The evidence is insufficient to determine the effects of the technology on health outcomes.

Primary Treatment for Osteochondral Lesions

For individuals with osteochondral lesions who receive PRP injections, the evidence includes an open-labeled quasi-randomized study. Relevant outcomes are symptoms, functional outcomes, health status measures, quality of life, and treatment-related morbidity. The quasi-randomized study found a statistically significant greater impact on outcomes in the PRP group than in the hyaluronic acid group. Limitations of the evidence base include lack of adequately randomized studies, lack of blinding, lack of sham controls, and comparison only to an intervention of uncertain efficacy. The evidence is insufficient to determine the effects of the technology on health outcomes.

Primary Treatment for Knee or Hip Osteoarthritis

For individuals with knee or hip OA who receive PRP injections, the evidence includes multiple RCTs and systematic reviews. Relevant outcomes are symptoms, functional outcomes, health status measures, quality of life, and treatment-related morbidity. Most trials have compared platelet-rich plasma with hyaluronic acid for knee osteoarthritis. Systematic reviews have generally found that platelet-rich plasma was more effective than placebo or hyaluronic acid in reducing pain and improving function. However, systematic review authors have noted that their findings should be interpreted with caution due to important limitations including significant residual statistical heterogeneity, questionable clinical significance, and high risk of bias in study conduct. RCTs with follow-up durations of at least 12 months published subsequent to the systematic reviews found statistically significantly greater 12 month reductions in the Western Ontario and McMaster Universities Osteoarthritis Index scores, but these findings were also limited by important study conduct flaws including potential inadequate control for selection bias and unclear blinding. Also, benefits were not maintained at 5 years. Also, using hyaluronic acid as a comparator is questionable, because the evidence demonstrating the benefit of hyaluronic acid treatment for osteoarthritis is not robust. The single RCT evaluating hip osteoarthritis reported statistically significant reductions in visual analog scale scores for pain, with no difference in functional scores. Additional studies comparing platelet-rich plasma with placebo and with alternatives other than hyaluronic acid are needed to determine the efficacy of platelet-rich plasma for knee and hip osteoarthritis. Studies are also needed to determine the optimal protocol for delivering platelet-rich plasma. The evidence is insufficient to determine the effects of the technology on health outcomes.

Adjunct to Surgery

For individuals with anterior cruciate ligament (ACL) reconstruction who receive PRP injections plus orthopedic surgery, the evidence includes two systematic reviews of multiple RCTs and prospective studies. Relevant outcomes are symptoms, functional outcomes, health status measures, quality of life, morbid events, resource utilization, and treatment-related morbidity. Only one of the two systematic reviews conducted a meta-analysis; it showed that adjunctive PRP treatment did not result in a significant effect on International Knee Documentation Committee

POLICY TITLE	ORTHOPEDIC APPLICATIONS OF PLATELET- RICH PLASMA
POLICY NUMBER	MP-4.039

scores, a patient-reported, knee-specific outcome measure that assesses pain and functional activity. Individual trials have shown mixed results. The evidence is insufficient to determine the effects of the technology on health outcomes

For individuals with hip fracture who receive PRP injections plus orthopedic surgery, the evidence includes an open-labeled RCT. Relevant outcomes are symptoms, functional outcomes, health status measures, quality of life, morbid events, resource utilization, and treatment-related morbidity. The single open-labeled RCT failed to show a statistically significant reduction in the need for surgical revision with the addition of PRP treatment. The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals with long bone nonunion who receive PRP injections plus orthopedic surgery, the evidence includes three RCTs. Relevant outcomes are symptoms, functional outcomes, health status measures, quality of life, morbid events, resource utilization, and treatment-related morbidity. One trial with a substantial risk of bias failed to show significant differences in patient-reported or clinician-assessed functional outcome scores between those who received PRP plus allogenic bone graft and those who received only allogenic bone graft. While the trial showed a statistically significant increase in the proportion of bones that healed in patients receiving PRP in a modified intention-to-treat analysis, the results did not differ in the intention-to-treat analysis. The second RCT, which compared PRP with recombinant human bone morphogenetic protein-7, also failed to show any clinical or radiologic benefits of PRP over morphogenetic protein. The third RCT reported no difference in the number of unions or time to union in patients receiving PRP injections vs no treatment. The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals with rotator cuff repair who receive PRP injections plus orthopedic surgery, the evidence includes multiple RCTs and systematic reviews. The relevant outcomes are symptoms, functional outcomes, health status measures, quality of life, morbid events, resource utilization, and treatment-related morbidity. Although systematic reviews consistently found significant reductions in pain with platelet-rich plasma at 12 months, important study conduct and relevance weaknesses limit interpretation of these findings. Additionally, the pain reductions with platelet-rich plasma were not maintained in longer-term studies. Further, the systematic reviews and meta-analyses failed to show a statistically and/or clinically significant impact on other outcomes. Findings of subsequently published small, single-center RCTs were consistent with the systematic reviews. The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals with spinal fusion who receive PRP injections plus orthopedic surgery, the evidence includes two controlled prospective studies. Relevant outcomes are symptoms, functional outcomes, health status measures, quality of life, morbid events, resource utilization, and treatment-related morbidity. The two studies failed to show any statistically significant

POLICY TITLE	ORTHOPEDIC APPLICATIONS OF PLATELET- RICH PLASMA
POLICY NUMBER	MP-4.039

differences in fusion rates between the PRP arm and the control arm. The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals undergoing spinal fusion who receive platelet-rich plasma injections, the evidence includes a single small RCT and a few observational studies. Relevant outcomes include symptoms, functional outcomes, health status measures, quality of life, morbid events, resource utilization, and treatment-related morbidity. Studies have generally failed to show a statistically and/or clinically significant impact on symptoms (ie, pain). The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals with subacromial decompression surgery who receive PRP injections plus orthopedic surgery, the evidence includes a small RCT. Relevant outcomes are symptoms, functional outcomes, health status measures, quality of life, morbid events, resource utilization, and treatment-related morbidity. A single small RCT failed to show a reduction in self-assessed or physician-assessed spinal instability scores with PRP injections. However, subjective pain, use of pain medications, and objective measures of range of motion showed clinically significant improvements with PRP. Larger trials are required to confirm these benefits. The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals with total knee arthroplasty (TKA) who receive PRP injections plus orthopedic surgery, the evidence includes a small RCT. The relevant outcomes are symptoms, functional outcomes, health status measures, quality of life, morbid events, resource utilization, and treatment-related morbidity. The RCT showed no significant differences between the PRP and untreated control groups in bleeding, range of motion, swelling around the knee joint, muscle power recovery, pain, or Knee Society Score and Knee Injury and Osteoarthritis Outcome Score. The evidence is insufficient to determine the effects of the technology on health outcomes.

V. DEFINITIONS

[TOP](#)

N/A

VI. BENEFIT VARIATIONS

[TOP](#)

The existence of this medical policy does not mean that this service is a covered benefit under the member's health benefit plan. Benefit determinations should be based in all cases on the applicable health benefit plan language. Medical policies do not constitute a description of benefits. A member's health benefit plan governs which services are covered, which are excluded, which are subject to benefit limits and which require preauthorization. There are different benefit plan designs in each product administered by Capital BlueCross. Members and providers should consult the member's health benefit plan for information or contact Capital BlueCross for benefit information.

POLICY TITLE	ORTHOPEDIC APPLICATIONS OF PLATELET- RICH PLASMA
POLICY NUMBER	MP-4.039

VII. DISCLAIMER

[TOP](#)

Capital BlueCross’s medical policies are developed to assist in administering a member’s benefits, do not constitute medical advice and are subject to change. Treating providers are solely responsible for medical advice and treatment of members. Members should discuss any medical policy related to their coverage or condition with their provider and consult their benefit information to determine if the service is covered. If there is a discrepancy between this medical policy and a member’s benefit information, the benefit information will govern. If a provider or a member has a question concerning the application of this medical policy to a specific member’s plan of benefits, please contact Capital BlueCross’ Provider Services or Member Services. Capital BlueCross considers the information contained in this medical policy to be proprietary and it may only be disseminated as permitted by law.

VIII. CODING INFORMATION

[TOP](#)

Note: This list of codes may not be all-inclusive, and codes are subject to change at any time. The identification of a code in this section does not denote coverage as coverage is determined by the terms of member benefit information. In addition, not all covered services are eligible for separate reimbursement.

Investigational therefore, not covered when used for orthopedic applications:

CPT Codes®								
0232T	86999							

Current Procedural Terminology (CPT) copyrighted by American Medical Association. All Rights Reserved.

IX. REFERENCES

[TOP](#)

1. Crovetti G, Martinelli G, Issi M, et al. Platelet gel for healing cutaneous chronic wounds. *Transfus Apher Sci.* Apr 2004;30(2):145-151. PMID 15062754.
2. Eppley BL, Woodell JE, Higgins J. Platelet quantification and growth factor analysis from platelet-rich plasma: implications for wound healing. *Plast Reconstr Surg.* Nov 2004;114(6):1502-1508. PMID 15509939.
3. Keyv SV, Jacobson MS. Comparison of methods for point of care preparation of autologous platelet gel. *J Extra Corpor Technol.* Mar 2004;36(1):28-35. PMID 15095838.
4. Castillo TN, Pouliot MA, Kim HJ, et al. Comparison of growth factor and platelet concentration from commercial platelet-rich plasma separation systems. *Am J Sports Med.* Feb 2011;39(2):266-271. PMID 21051428.
5. Mazzucco L, Balbo V, Cattana E, et al. Not every PRP-gel is born equal. Evaluation of growth factor availability for tissues through four PRP-gel preparations: Fibrinet, RegenPRP-Kit, Plateltex and one manual procedure. *Vox Sang.* Aug 2009;97(2):110-118. PMID 19392780.

POLICY TITLE	ORTHOPEDIC APPLICATIONS OF PLATELET- RICH PLASMA
POLICY NUMBER	MP-4.039

6. Hsu WK, Mishra A, Rodeo SR, et al. Platelet-rich plasma in orthopaedic applications: evidence-based recommendations for treatment. *J Am Acad Orthop Surg. Dec 2013;21(12):739-748. PMID 24292930.*
7. Johal H, Khan M, Yung SP et al. Impact of Platelet-Rich Plasma Use on Pain in Orthopaedic Surgery: A Systematic Review and Meta-analysis. *Sports Health. 2019 Jul;11(4). PMID 31136726*
7. Miller LE, Parrish WR, Roides B, et al. Efficacy of platelet-rich plasma injections for symptomatic tendinopathy: systematic review and meta-analysis of randomised injection-controlled trials. *BMJ Open Sport Exerc Med. Nov 6 2017;3(1):e000237. PMID 29177072.*
8. Tsikopoulos K, Tsikopoulos I, Simeonidis E, et al. The clinical impact of platelet-rich plasma on tendinopathy compared to placebo or dry needling injections: A meta-analysis. *Phys Ther Sport. Jan 2016;17:87-94. PMID 26621224.*
9. Balasubramaniam U, Dissanayake R, Annabell L. Efficacy of platelet-rich plasma injections in pain associated with chronic tendinopathy: A systematic review. *Phys Sportsmed. Jul 2015;43(3):253-261. PMID 25599747.*
10. Andia I, Latorre PM, Gomez MC, et al. Platelet-rich plasma in the conservative treatment of painful tendinopathy: a systematic review and meta-analysis of controlled studies. *Br Med Bull. Jun 2014;110(1):99-115. PMID 24795364.*
11. Gupta PK, Acharya A, Khanna V et al. PRP versus steroids in a deadlock for efficacy: long-term stability versus short-term intensity-results from a randomised trial. *Musculoskelet Surg. 2019 Aug. PMID 31448392*
12. Scott A, LaPrade RF, Harmon KG et al. Platelet-Rich Plasma for Patellar Tendinopathy: A Randomized Controlled Trial of Leukocyte-Rich PRP or Leukocyte-Poor PRP Versus Saline. *Am J Sports Med. 2019 Jun;47(7). PMID 31038979*
13. Fitzpatrick J, Bulsara MK, O'Donnell J et al. Leucocyte-Rich Platelet-Rich Plasma Treatment of Gluteus Medius and Minimus Tendinopathy: A Double-Blind Randomized Controlled Trial With 2-Year Follow-up. *Am J Sports Med. 2019 Apr;47(5). PMID 30840831*
14. Martin JI, Atilano L, Bully P et al. Needle tenotomy with PRP versus lidocaine in epicondylopathy: clinical and ultrasonographic outcomes over twenty months. *Skeletal Radiol. 2019 Sep;48(9). PMID 30826853*
15. Franceschi F, Papalia R, Franceschetti E, et al. Platelet-rich plasma injections for chronic plantar fasciopathy: a systematic review. *Br Med Bull. Dec 2014;112(1):83-95. PMID 25239050.*
16. Monto RR. Platelet-rich plasma efficacy versus corticosteroid injection treatment for chronic severe plantar fasciitis. *Foot Ankle Int. Apr 2014;35(4):313-318. PMID 24419823.*
17. Peerbooms JC, Lodder P, den Oudsten BL et al. Positive Effect of Platelet-Rich Plasma on Pain in Plantar Fasciitis: A Double-Blind Multicenter Randomized Controlled Trial. *Am J Sports Med. 2019 Nov;47(13). PMID 31603721*

POLICY TITLE	ORTHOPEDIC APPLICATIONS OF PLATELET- RICH PLASMA
POLICY NUMBER	MP-4.039

18. Shetty SH, Dhond A, Arora M, et al. Platelet-Rich Plasma Has Better Long-Term Results Than Corticosteroids or Placebo for Chronic Plantar Fasciitis: Randomized Control Trial. *J Foot Ankle Surg.* Jan 2019;58(1):42-46. PMID 30448183
19. Johnson-Lynn S, Cooney A, Ferguson D et al. A Feasibility Study Comparing Platelet-Rich Plasma Injection With Saline for the Treatment of Plantar Fasciitis Using a Prospective, Randomized Trial Design. *Foot Ankle Spec.* 2019 Apr;12(2). PMID 29779399
20. Mei-Dan O, Carmont MR, Laver L, et al. Platelet-rich plasma or hyaluronate in the management of osteochondral lesions of the talus. *Am J Sports Med.* Mar 2012;40(3):534-541. PMID 22253252
21. Laudy AB, Bakker EW, Rekers M, et al. Efficacy of platelet-rich plasma injections in osteoarthritis of the knee: a systematic review and meta-analysis. *Br J Sports Med.* May 2015;49(10):657-672. PMID 25416198.
22. Chang KV, Hung CY, Aliwarga F, et al. Comparative effectiveness of platelet-rich plasma injections for treating knee joint cartilage degenerative pathology: a systematic review and meta-analysis. *Arch Phys Med Rehabil.* Mar 2014;95(3):562-575. PMID 24291594.
23. Meheux CJ, McCulloch PC, Lintner DM, et al. Efficacy of intra-articular platelet-rich plasma injections in knee osteoarthritis: a systematic review. *Arthroscopy.* Mar 2016;32(3):495-505. PMID 26432430.
24. Lai LP, Stitik TP, Foye PM, et al. Use of platelet-rich plasma in intra-articular knee injections for osteoarthritis: a systematic review. *PM R.* Jun 2015;7(6):637-648. PMID 25687110.
25. Cole BJ, Karas V, Hussey K, et al. Hyaluronic acid versus platelet-rich plasma. *Am J Sports Med.* Feb 2017;45(2):339-346. PMID 28146403.
26. Duymus TM, Mutlu S, Dernek B, et al. Choice of intra-articular injection in treatment of knee osteoarthritis: platelet-rich plasma, hyaluronic acid or ozone options. *Knee Surg Sports Traumatol Arthrosc.* Feb 2017;25(2):485-492. PMID 27056686.
27. Kanchanatawan W, Arirachakaran A, Chaijenkij K, et al. Short-term outcomes of platelet-rich plasma injection for treatment of osteoarthritis of the knee. *Knee Surg Sports Traumatol Arthrosc.* May 2016;24(5):1665-1677. PMID 26387122.
28. Xu Z, Luo J, Huang X, et al. Efficacy of platelet-rich plasma in pain and self-report function in knee osteoarthritis: a best-evidence synthesis. *Am J Phys Med Rehabil.* Nov 2017;96(11):793-800. PMID 28398969.
29. Huang Y, Liu X, Xu X et al. Intra-articular injections of platelet-rich plasma, hyaluronic acid or corticosteroids for knee osteoarthritis : A prospective randomized controlled study. *Orthopade.* 2019 Mar;48(3). PMID 30623236
30. Di Martino A, Di Matteo B, Papio T et al. Platelet-Rich Plasma Versus Hyaluronic Acid Injections for the Treatment of Knee Osteoarthritis: Results at 5 Years of a Double-Blind, Randomized Controlled Trial. *Am J Sports Med.* 2019 Feb;47(2). PMID 30545242
31. Lin KY, Yang CC, Hsu CJ et al. Intra-articular Injection of Platelet-Rich Plasma Is Superior to Hyaluronic Acid or Saline Solution in the Treatment of Mild to Moderate Knee

POLICY TITLE	ORTHOPEDIC APPLICATIONS OF PLATELET- RICH PLASMA
POLICY NUMBER	MP-4.039

- Osteoarthritis: A Randomized, Double-Blind, Triple-Parallel, Placebo-Controlled Clinical Trial. Arthroscopy. 2019 Jan;35(1). PMID 30611335*
32. Patel S, Dhillon MS, Aggarwal S, et al. Treatment with platelet-rich plasma is more effective than placebo for knee osteoarthritis: a prospective, double-blind, randomized trial. *Am J Sports Med.* Feb 2013;41(2):356-364. PMID 23299850.
 33. Tubach F, Ravaud P, Baron G, et al. Evaluation of clinically relevant changes in patient reported outcomes in knee and hip osteoarthritis: the minimal clinically important improvement. *Ann Rheum Dis.* Jan 2005;64(1):29-33. PMID 15208174.
 34. Smith PA. Autologous conditioned plasma injections provide safe and efficacious treatment for knee osteoarthritis: an FDA-sanctioned, randomized, double-blind, placebo-controlled clinical trial. *Am J Sports Med.* Apr 2016;44(4):884-891. PMID 26831629.
 35. Dallari D, Stagni C, Rani N, et al. Ultrasound-guided injection of platelet-rich plasma and hyaluronic acid, separately and in combination, for hip osteoarthritis: a randomized controlled study. *Am J Sports Med.* Mar 2016;44(3):664-671. PMID 26797697.
 36. Trueba Vasavilbaso C, Rosas Bello CD, Medina Lopez E, et al. Benefits of different postoperative treatments in patients undergoing knee arthroscopic debridement. *Open Access Rheumatol.* Sep 25 2017;9:171-179. PMID 29026341
 37. Figueroa D, Figueroa F, Calvo R, et al. Platelet-rich plasma use in anterior cruciate ligament surgery: systematic review of the literature. *Arthroscopy.* May 2015;31(5):981-988. PMID 25595696
 38. Nin JR, Gasque GM, Azcarate AV, et al. Has platelet-rich plasma any role in anterior cruciate ligament allograft healing? *Arthroscopy.* Nov 2009;25(11):1206-1213. PMID 19896041
 39. Griffin XL, Achten J, Parsons N, et al. Platelet-rich therapy in the treatment of patients with hip fractures: a single centre, parallel group, participant-blinded, randomised controlled trial. *BMJ Open.* Jun 25 2013;3(6). PMID 23801709.
 40. Griffin XL, Wallace D, Parsons N, et al. Platelet rich therapies for long bone healing in adults. *Cochrane Database Syst Rev.* Jul 11 2012;7(7):CD009496. PMID 22786528.
 41. Calori GM, Tagliabue L, Gala L, et al. Application of rhBMP-7 and platelet-rich plasma in the treatment of long bone non-unions: a prospective randomised clinical study on 120 patients. *Injury.* Dec 2008;39(12):1391-1402. PMID 19027898.
 42. Dallari D, Savarino L, Stagni C, et al. Enhanced tibial osteotomy healing with use of bone grafts supplemented with platelet gel or platelet gel and bone marrow stromal cells. *J Bone Joint Surg Am.* Nov 2007;89(11):2413- 2420. PMID 17974883.
 43. Samuel G, Menon J, Thimmaiah S, et al. Role of isolated percutaneous autologous platelet concentrate in delayed union of long bones. *Eur J Orthop Surg Traumatol.* Nov 22 2017. PMID 29167980.
 44. Wang Y, Han C, Hao J et al. Efficacy of platelet-rich plasma injections for treating Achilles tendonitis : Systematic review of high-quality randomized controlled trials. *Orthopade.* 2019 Sep;48(9). PMID 30937491

POLICY TITLE	ORTHOPEDIC APPLICATIONS OF PLATELET- RICH PLASMA
POLICY NUMBER	MP-4.039

45. Moraes VY, Lenza M, Tamaoki MJ, et al. Platelet-rich therapies for musculoskeletal soft tissue injuries. *Cochrane Database Syst Rev.* Dec 23 2013;12(12):CD010071. PMID 24363098.
46. Zhao JG, Zhao L, Jiang YX, et al. Platelet-rich plasma in arthroscopic rotator cuff repair: a meta-analysis of randomized controlled trials. *Arthroscopy.* Jan 2015;31(1):125-135. PMID 25278352.
47. Yang J, Sun Y, Xu P, et al. Can patients get better clinical outcomes by using PRP in rotator cuff repair: a meta- analysis of randomized controlled trials. *J Sports Med Phys Fitness.* Nov 2016;56(11):1359-1367. PMID 26473444.
48. Cai YZ, Zhang C, Lin XJ. Efficacy of platelet-rich plasma in arthroscopic repair of full-thickness rotator cuff tears: a meta-analysis. *J Shoulder Elbow Surg.* Dec 2015;24(12):1852-1859. PMID 26456434.
49. Chen X, Jones IA, Park C, et al. The efficacy of platelet-rich plasma on tendon and ligament healing: a systematic review and meta-analysis with bias assessment. *Am J Sports Med.* Dec 1 2017;363546517743746. PMID 29268037.
50. Fu CJ, Sun JB, Bi ZG, et al. Evaluation of platelet-rich plasma and fibrin matrix to assist in healing and repair of rotator cuff injuries: a systematic review and meta-analysis. *Clin Rehabil.* Feb 2017;31(2):158-172. PMID 26928856.
51. Saltzman BM, Jain A, Campbell KA, et al. Does the use of platelet-rich plasma at the time of surgery improve clinical outcomes in arthroscopic rotator cuff repair when compared with control cohorts? A systematic review of meta-analyses. *Arthroscopy.* May 2016;32(5):906-918. PMID 26725454.
52. Walsh MR, Nelson BJ, Braman JP, et al. Platelet-rich plasma in fibrin matrix to augment rotator cuff repair: a prospective, single-blinded, randomized study with 2-year follow-up. *J Shoulder Elbow Surg.* 2018 Sep;27(9):1553-1563. PMID: 29996980.
53. Malavolta EA, Gracitelli MEC, Assuncao JH, et al. Clinical and Structural Evaluations of Rotator Cuff Repair With and Without Added Platelet-Rich Plasma at 5-Year Follow-up: A Prospective Randomized Study. *Am J Sports Med.* Nov 2018;46(13):3134-3141. PMID 30234999
54. Snow M, Hussain F, Pagkalos J et al. The Effect of Delayed Injection of Leukocyte-Rich Platelet-Rich Plasma (LR-PRP) Following Rotator Cuff Repair on Patient Function: A randomized Double-Blind Controlled Trial. *Arthroscopy.* 2019 Nov. PMID 31784365
55. Kubota G, Kamoda H, Orita S et al. Platelet-rich plasma enhances bone union in posterolateral lumbar fusion: A prospective randomized controlled trial. *Spine J.* 2019 Feb;19(2). PMID 28735763
56. Ebert JR, Wang A, Smith A, et al. A midterm evaluation of postoperative platelet-rich plasma injections on arthroscopic supraspinatus repair: a randomized controlled trial. *Am J Sports Med.* Nov 2017;45(13):2965-2974. PMID 28806095.

MEDICAL POLICY

POLICY TITLE	ORTHOPEDIC APPLICATIONS OF PLATELET- RICH PLASMA
POLICY NUMBER	MP-4.039

57. Carreon LY, Glassman SD, Anekstein Y, et al. Platelet gel (AGF) fails to increase fusion rates in instrumented posterolateral fusions. *Spine (Phila Pa 1976)*. May 1 2005;30(9):E243-246; discussion E247. PMID 15864142.

58. Tsai CH, Hsu HC, Chen YJ, et al. Using the growth factors-enriched platelet glue in spinal fusion and its efficiency. *J Spinal Disord Tech*. Jun 2009;22(4):246-250. PMID 19494743.

59. Everts PA, Devilee RJ, Brown Mahoney C, et al. Exogenous application of platelet-leukocyte gel during open subacromial decompression contributes to improved patient outcome. A prospective randomized double-blind study. *Eur Surg Res*. Nov 2008;40(2):203-210. PMID 17998780.

60. Morishita M, Ishida K, Matsumoto T, et al. Intraoperative platelet-rich plasma does not improve outcomes of total knee arthroplasty. *J Arthroplasty*. Dec 2014;29(12):2337-2341. PMID 24851794.

61. American Academy of Orthopaedic Surgeons. *Treatment of osteoarthritis of the knee*. 2013; www.aaos.org. Accessed August 20, 2020.

62. American Academy of Orthopaedic Surgeons A. *Management of Osteoarthritis of the Hip - Evidence-Based Clinical Practice Guideline*. 2017; https://www.aaos.org/uploadedFiles/PreProduction/Quality/Guidelines_and_Reviews/OA%20Hip%20CPG_3.13.17.pdf. Accessed August 20, 2020.

63. National Institute for Health and Care Excellence (NICE). *Autologous blood injection for tendinopathy [IPG438]*. 2013; <https://www.nice.org.uk/guidance/ipg438>. Accessed August 20, 2020.

64. National Institute for Health and Care Excellence. *Autologous blood injection for plantar fasciitis [IPG437]*. 2013; <https://www.nice.org.uk/guidance/ipg437>. Accessed August 20, 2020.

65. National Institute for Health and Care Excellence (NICE). *Platelet-rich plasma injections for osteoarthritis of the knee: Interventional procedure guidance [IPG637]*. 2019; <https://www.nice.org.uk/guidance/ipg637>. Accessed August 20, 2020.

66. Blue Cross Blue Shield Association Medical Policy Reference Manual. 2.01.98, *Orthopedic Applications of Platelet-Rich Plasma*. May 2020.

X. POLICY HISTORY

[TOP](#)

MP-4.039	CAC 9/29/15 New policy. BCBSA adopted. Policy created on the orthopedic applications of platelet-rich plasma that were previously described in MP-2.033 Recombinant and Autologous Platelet-Derived Growth Factors as a Treatment of Wound Healing and Other Conditions. All indications are considered investigational. FEP variation added. Codes applied.
	CAC 11/29/16 Consensus review. Policy statements unchanged. Variation reformatting completed. FEP variation updated. Cross Reference,

POLICY TITLE	ORTHOPEDIC APPLICATIONS OF PLATELET- RICH PLASMA
POLICY NUMBER	MP-4.039

	Description/Background, Rationale, and Reference sections updated. Coding Reviewed.
	12/19/17 Consensus review. No changes to the policy statements. Background, rationale and references updated.
	11/26/18 Consensus review. No changes to the policy statements. Background, and references updated. Rationale revised.
	9/30/19 Consensus review. No changes to the policy statements. Background, summary of evidence, and references updated. FEP variation removed since archived.
	8/19/2020 Consensus review. No change to policy statement. Rationale and References updated.

[Top](#)

Health care benefit programs issued or administered by Capital BlueCross and/or its subsidiaries, Capital Advantage Insurance Company®, Capital Advantage Assurance Company® and Keystone Health Plan® Central. Independent licensees of the BlueCross BlueShield Association. Communications issued by Capital BlueCross in its capacity as administrator of programs and provider relations for all companies.