

MEDICAL POLICY

POLICY TITLE	LOW INTENSITY PULSED ULTRASOUND FRACTURE HEALING DEVICE
POLICY NUMBER	MP 6.021

CLINICAL BENEFIT	<input type="checkbox"/> MINIMIZE SAFETY RISK OR CONCERN. <input checked="" type="checkbox"/> MINIMIZE HARMFUL OR INEFFECTIVE INTERVENTIONS. <input type="checkbox"/> ASSURE APPROPRIATE LEVEL OF CARE. <input type="checkbox"/> ASSURE APPROPRIATE DURATION OF SERVICE FOR INTERVENTIONS. <input type="checkbox"/> ASSURE THAT RECOMMENDED MEDICAL PREREQUISITES HAVE BEEN MET. <input type="checkbox"/> ASSURE APPROPRIATE SITE OF TREATMENT OR SERVICE.
Effective Date:	10/1/2024

[POLICY RATIONALE](#)
[DISCLAIMER](#)
[POLICY HISTORY](#)

[PRODUCT VARIATIONS](#)
[DEFINITIONS](#)
[CODING INFORMATION](#)

[DESCRIPTION/BACKGROUND](#)
[BENEFIT VARIATIONS](#)
[REFERENCES](#)

I. POLICY

Low-intensity pulsed ultrasound is considered **not medically necessary** for treatment of the following:

- Fresh fractures (surgically managed or non-surgically managed).
- Fracture nonunion and delayed union fractures.
- Stress fractures, osteotomy, and distraction osteogenesis.

There is insufficient evidence to support a conclusion concerning the health outcomes or benefits associated with these procedures.

Policy Guidelines

Fresh (acute) fracture

There is no standard definition for a “fresh” fracture. A fracture is most commonly defined as fresh for 7 days after the fracture occurs but there is definition variability. For example, 1 study defined fresh as less than 5 days after fracture while another defined fresh as up to 10 days post fracture. Most fresh closed fractures heal without complications using standard fracture care (i.e., closed reduction and cast immobilization).

Nonunion

There is no consensus on the definition of nonunions. One definition is a failure of progression of fracture healing for at least 3 consecutive months (and at least 6 months post-fracture) accompanied by clinical symptoms of delayed/nonunion (pain, difficulty weight bearing.).

The definition of nonunion used in U.S. Food and Drug Administration labeling suggests that nonunion is considered established when the fracture site shows no visibly progressive signs of healing, without providing guidance on the timeframe of observation. The following selection

MEDICAL POLICY

POLICY TITLE	LOW INTENSITY PULSED ULTRASOUND FRACTURE HEALING DEVICE
POLICY NUMBER	MP 6.021

criteria are consistent with those proposed for electrical stimulation as a treatment of nonunions (see evidence review 7.01.07):

- At least 3 months have passed since the date of the fracture, and
- serial radiographs have confirmed that no progressive signs of healing have occurred, and
- the fracture gap is 1 cm or less, and
- the individual can be adequately immobilized and based on age, is likely to comply with nonweight bearing.

Delayed Union

Delayed union is defined as a decelerating healing process as determined by serial radiographs, together with a lack of clinical and radiologic evidence of union, bony continuity, or bone reaction at the fracture site for no less than 3 months from the index injury or the most recent intervention.

Cross-reference:

MP 1.024 Electrical Bone Growth Stimulation of the Appendicular Skeleton

MP 1.150 Electrical Stimulation of the Spine as an Adjunct to Spinal Fusion Procedures

II. PRODUCT VARIATIONS

[TOP](#)

This policy is only applicable to certain programs and products administered by Capital Blue Cross and subject to benefit variations as discussed in Section VI. Please see additional information below.

FEP PPO - Refer to FEP Medical Policy Manual. The FEP Medical Policy manual can be found at: <https://www.fepblue.org/benefit-plans/medical-policies-and-utilization-management-guidelines/medical-policies>

III. DESCRIPTION/BACKGROUND

[TOP](#)

Low-intensity pulsed ultrasound (LIPUS) has been investigated as a technique to accelerate healing of fresh fractures, surgically treated closed fractures, delayed unions, nonunions, stress fractures, osteotomy sites, and distraction osteogenesis. LIPUS is administered using a transducer applied to the skin surface overlying the fracture site.

Bone Fractures

An estimated 178 million new fractures were reported worldwide in 2019. Most bone fractures heal spontaneously over the course of several months following standard fracture care (closed reduction, if necessary, followed by immobilization with casting or splinting). However, approximately 5% to 10% of all fractures have delayed healing, resulting in continued morbidity and increased utilization of health care services. Factors contributing to a nonunion include

MEDICAL POLICY

POLICY TITLE	LOW INTENSITY PULSED ULTRASOUND FRACTURE HEALING DEVICE
POLICY NUMBER	MP 6.021

which bone is fractured, fracture site, degree of bone loss, time since injury, extent of soft tissue injury, and patient factors (e.g., smoking, diabetes, systemic disease).

Fracture Nonunion

There is no standard definition of a fracture nonunion. The Food and Drug Administration has defined nonunion as when "a minimum of 9 months has elapsed since injury and the fracture site shows no visibly progressive signs of healing for a minimum of 3 months." Other definitions cite 3 to 6 months of time from the original injury, or simply when serial radiographs fail to show any further healing. These definitions do not reflect the underlying conditions in fractures that affect healing, such as the degree of soft tissue damage, alignment of the bone fragments, vascularity, and quality of the underlying bone stock.

Delayed Union

Delayed union is generally considered a failure to heal between 3- and 9-months post fracture, after which the fracture site would be considered a nonunion. Delayed union may also be defined as a decelerating bone healing process, as identified in serial radiographs. (In contrast, nonunion serial radiographs show no evidence of healing.) It is important to include both radiographic and clinical criteria to determine fracture healing status. Clinical criteria include the lack of ability to bear weight, fracture pain, and tenderness on palpation.

Treatment

Low-intensity pulsed ultrasound (LIPUS) has been proposed to accelerate healing of fractures. LIPUS is believed to alter the molecular and cellular mechanisms involved in each stage of the healing process (inflammation, soft callus formation, hard callus formation, and bone remodeling). The mechanism of action at the cellular level is not precisely known, but it is theorized that LIPUS may stimulate the production or the activities of the following compounds that contribute to the bone healing process: cyclooxygenase-2, collagenase, integrin proteins, calcium, chondroblasts, mesenchymal cells, fibroblasts, and osteoblasts.

LIPUS treatment is self-administered, once daily for 20 minutes, until the fracture has healed, usually for 5 months.

REGULATORY STATUS

In 1994, the Sonic Accelerated Fracture Healing System (SAFHS®; renamed Exogen 2000® and since 2006, Exogen 4000+; Bioventus) was approved by the U.S. Food and Drug Administration through the premarket approval process for treatment of fresh, closed, posteriorly displaced distal radius (Colles) fractures, and fresh, closed, or grade I open tibial diaphysis fractures in skeletally mature individuals when these fractures are orthopedically managed by closed reduction and cast immobilization. In February 2000, the labeled indication was expanded to include the treatment of established nonunions, excluding skull and vertebra. Food and Drug Administration product code: LPQ.

MEDICAL POLICY

POLICY TITLE	LOW INTENSITY PULSED ULTRASOUND FRACTURE HEALING DEVICE
POLICY NUMBER	MP 6.021

IV. RATIONALE

[TOP](#)

Summary of Evidence

For individuals who have fresh fractures (surgically or non-surgically managed) who receive low-intensity pulsed ultrasound (LIPUS), the evidence includes randomized controlled trials (RCTs) and several meta-analysis. Relevant outcomes are symptoms, morbid events, functional outcomes, and quality of life. The evidence base has recently evolved with the publication of a large RCT and meta-analysis significantly shifting the weight of the evidence. Conclusions based on several earlier small RCTs, rated at high risk of bias, showed a potential benefit of LIPUS; however, the large RCT published in 2016, rated at low risk of bias, showed no benefit. A 2017 meta-analysis including only trials with low risk of bias found no difference in days to full weight bearing, pain reduction, or days to radiographic healing. Similarly, the overall results of the meta-analysis found no significant difference in return to work, subsequent operations, or adverse effect. The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals who have fracture nonunion or delayed union fracture who receive LIPUS, the evidence includes only lower quality studies including a small systematic review in scaphoid nonunions, a meta-analysis of nonunion in various locations, a meta-analysis in individuals with specific risk factors, 2 low-quality RCTs, and 1 observational comparative study. Relevant outcomes are symptoms, morbid events, functional outcomes, and quality of life. Of the 2 RCTs, one did not include functional outcomes. The second RCT had a small sample size and did not describe the randomization procedure. The observational study reported similar healing rates with low-intensity pulsed ultrasound and surgery, although the retrospective nature of the study, limits meaningful interpretation of these results. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have stress fractures, osteotomy sites, or distraction osteogenesis who receive low-intensity pulsed ultrasound (LIPUS) as an adjunct to routine care, the evidence includes only lower quality studies including small RCTs. Relevant outcomes are symptoms, morbid events, functional outcomes, and quality of life. Results do not generally include functional outcomes and results across various outcomes, primarily time to radiographic healing, are inconsistent. The meta-analysis of 3 trials using low-intensity pulsed ultrasound for distraction osteogenesis reported no statistically significant differences in physiological or functional outcomes. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

V. DEFINITIONS

[TOP](#)

NA

VI. BENEFIT VARIATIONS

[TOP](#)

The existence of this medical policy does not mean that this service is a covered benefit under the member's health benefit plan. Benefit determinations should be based in all cases on the

MEDICAL POLICY

POLICY TITLE	LOW INTENSITY PULSED ULTRASOUND FRACTURE HEALING DEVICE
POLICY NUMBER	MP 6.021

applicable health benefit plan language. Medical policies do not constitute a description of benefits. A member's health benefit plan governs which services are covered, which are excluded, which are subject to benefit limits, and which require preauthorization. There are different benefit plan designs in each product administered by Capital Blue Cross. Members and providers should consult the member's health benefit plan for information or contact Capital Blue Cross for benefit information.

VII. DISCLAIMER

[TOP](#)

Capital Blue Cross' medical policies are developed to assist in administering a member's benefits, do not constitute medical advice and are subject to change. Treating providers are solely responsible for medical advice and treatment of members. Members should discuss any medical policy related to their coverage or condition with their provider and consult their benefit information to determine if the service is covered. If there is a discrepancy between this medical policy and a member's benefit information, the benefit information will govern. If a provider or a member has a question concerning the application of this medical policy to a specific member's plan of benefits, please contact Capital Blue Cross' Provider Services or Member Services. Capital Blue Cross considers the information contained in this medical policy to be proprietary and it may only be disseminated as permitted by law.

VIII. CODING INFORMATION

[TOP](#)

Note: This list of codes may not be all-inclusive, and codes are subject to change at any time. The identification of a code in this section does not denote coverage as coverage is determined by the terms of member benefit information. In addition, not all covered services are eligible for separate reimbursement.

Low-intensity pulsed ultrasound is considered Not Medically Necessary: therefore, not covered:

Procedure Codes							
20979	E0760						

IX. REFERENCES

[TOP](#)

1. Mayr E, Rudzki MM, Rudzki M, et al. [Does low intensity, pulsed ultrasound speed healing of scaphoid fractures?]. *Handchir Mikrochir Plast Chir.* Mar 2000; 32(2): 115-22. PMID 10857066
2. Buza JA, 3rd, Einhorn T. Bone healing in 2016. *Clin Cases Miner Bone Metab.* May-Aug 2016; 13(2):101-105. PMID 27920804
3. Bhandari M, Fong K, Sprague S, et al. Variability in the definition and perceived causes of delayed unions and nonunions: a cross-sectional, multinational survey of orthopaedic surgeons. *J Bone Joint Surg Am.* Aug 1 2012; 94(15):e1091-1096. PMID 22854998
4. Schandelmaier S, Kaushal A, Lytvyn L, et al. Low-intensity pulsed ultrasound for bone healing: systematic review of randomized controlled trials. *BMJ.* Feb 22 2017; 356:j656. PMID 28348110

MEDICAL POLICY

POLICY TITLE	LOW INTENSITY PULSED ULTRASOUND FRACTURE HEALING DEVICE
POLICY NUMBER	MP 6.021

5. Seger EW, Jauregui JJ, Horton SA, et al. Low-intensity pulsed ultrasound for nonoperative treatment of scaphoid nonunions: a meta-analysis. *Hand (N Y)*. Apr 01 2017;1558944717702470. PMID 28391752
6. Lou S, Lv H, Li Z, et al. The effects of low-intensity pulsed ultrasound on fresh fracture: A meta-analysis. *Medicine (Baltimore)*. Sep 2017; 96(39):e8181. PMID 28953676
7. Leighton R, Watson JT, Giannoudis P, et al. Healing of fracture nonunions treated with low-intensity pulsed ultrasound (LIPUS): A systematic review and meta-analysis. *Injury*. Jul 2017; 48(7):1339-1347. PMID 28532896
8. Griffin XL, Parsons N, Costa ML, et al. Ultrasound and shockwave therapy for acute fractures in adults. *Cochrane Database Syst Rev*. Jun 23 2014; 6(6):CD008579. PMID 24956457
9. Busse JW, Kaur J, Mollon B, et al. Low-intensity pulsed ultrasonography for fractures: systematic review of randomised controlled trials. *BMJ*. Feb 27 2009; 338:b351. PMID 19251751
10. Blue Cross and Blue Shield Association Technology Evaluation Center (TEC). Ultrasound accelerated fracture healing. TEC Assessments 1995; Volume 10: Tab 14.
11. Schortinghuis J, Bronckers AL, Stegenga B, et al. Ultrasound to stimulate early bone formation in a distraction gap: a double-blind randomised clinical pilot trial in the edentulous mandible. *Arch Oral Biol*. Apr 2005; 50(4):411- 420. PMID 15748694
12. Schortinghuis J, Bronckers AL, Gravendeel J, et al. The effect of ultrasound on osteogenesis in the vertically distracted edentulous mandible: a double-blind trial. *Int J Oral Maxillofac Surg*. Nov 2008; 37(11):1014-1021. PMID 18757179
13. Strauss E, Ryaby JP, McCabe J. Treatment of Jones' fractures of the foot with adjunctive use of low-pulsed ultrasound stimulation. *J Orthop Trauma*. 1999; 13(4):310
14. Busse JW, Bhandari M, Einhorn TA, et al. Re-evaluation of low-intensity pulsed ultrasound in treatment of tibial fractures (TRUST): randomized clinical trial. *BMJ*. Oct 25 2016; 355:i5351. PMID 27797787
15. Tarride JE, Hopkins RB, Blackhouse G, et al. Low-intensity pulsed ultrasound for treatment of tibial fractures: an economic evaluation of the TRUST study. *Bone Joint J*. Nov 2017; 99-B (11):1526-1532. PMID 29092994
16. Emami A, Petren-Mallmin M, Larsson S. No effect of low-intensity ultrasound on healing time of intramedullary fixed tibial fractures. *J Orthop Trauma*. May 1999; 13(4):252-257. PMID 10342350
17. Lubbert PH, van der Rijt RH, Hoorntje LE, et al. Low-intensity pulsed ultrasound (LIPUS) in fresh clavicle fractures: a multi-centre double-blind randomised controlled trial. *Injury*. Dec 2008; 39(12):1444-1452. PMID 18656872
18. Schofer MD, Block JE, Aigner J, et al. Improved healing response in delayed unions of the tibia with low-intensity pulsed ultrasound: results of a randomized sham-controlled trial. *BMC Musculoskelet Disord*. Oct 08 2010; 11:229. PMID 20932272
19. Ricardo M. The effect of ultrasound on the healing of muscle-pediculated bone graft in scaphoid non-union. *Int Orthop*. Apr 2006; 30(2):123-127. PMID 16474939
20. Nolte, PP, Anderson, RR, Strauss, EE, Wang, ZZ, Hu, LL, Xu, ZZ, Steen, RR. Heal rate of metatarsal fractures: A propensity-matching study of patients treated with low-intensity pulsed ultrasound (LIPUS) vs. surgical and other treatments. *Injury*, 2016 Nov 5; 47(11). PMID 27641221

MEDICAL POLICY

POLICY TITLE	LOW INTENSITY PULSED ULTRASOUND FRACTURE HEALING DEVICE
POLICY NUMBER	MP 6.021

21. Rue JP, Armstrong DW, 3rd, Frassica FJ, et al. The effect of pulsed ultrasound in the treatment of tibial stress fractures. *Orthopedics*. Nov 2004; 27(11):1192-1195. PMID 15566133
22. Urita A, Iwasaki N, Kondo M, et al. Effect of low-intensity pulsed ultrasound on bone healing at osteotomy sites after forearm bone shortening. *J Hand Surg Am*. Mar 2013; 38(3):498-503. PMID 23375786
23. Dudda M, Hauser J, Muhr G, et al. Low-intensity pulsed ultrasound as a useful adjuvant during distraction osteogenesis: a prospective, randomized controlled trial. *J Trauma*. Nov 2011; 71(5):1376-1380. PMID 22071933
24. Salem KH, Schmelz A. Low-intensity pulsed ultrasound shortens the treatment time in tibial distraction osteogenesis. *Int Orthop*. Jul 2014; 38(7):1477-1482. PMID 24390009
25. El-Mowafi H, Mohsen M. The effect of low-intensity pulsed ultrasound on callus maturation in tibial distraction osteogenesis. *Int Orthop*. Apr 2005; 29(2):121-124. PMID 15685456
26. Tsumaki N, Kakiuchi M, Sasaki J, et al. Low-intensity pulsed ultrasound accelerates maturation of callus in patients treated with opening-wedge high tibial osteotomy by hemicallosis. *J Bone Joint Surg Am*. Nov 2004; 86-A (11):2399-2405. PMID 15523009
27. Lou S, Lv H, Li Z, Tang P, Wang Y. Effect of low-intensity pulsed ultrasound on distraction osteogenesis: a systematic review and meta-analysis of randomized controlled trials. *J Ortho Surg and Research* 2018; 13(1)205. PMID 30119631
28. MAGIC: Making GRADE the Irresistible Choice. n.d.
29. Poolman RW, Agoritsas T, Siemieniuk RA, et al. Low-intensity pulsed ultrasound (LIPUS) for bone healing: a clinical practice guideline. *BMJ*. Feb 21 2017; 356:j576. PMID 28228381
30. National Institute for Health and Care Excellence (NICE). Low-intensity pulsed ultrasound to promote healing of fresh fractures at high risk of non-healing [IPG622]. 2018
31. National Institute for Health and Care Excellence (NICE). Low-intensity pulsed ultrasound to promote healing of delayed-union and non-union fractures [IPG623]. 2018
32. National Institute for Health and Care Excellence (NICE). EXOGEN ultrasound bone healing system for long bone fractures with non-union or delayed healing [MTG12]. 2013 (Updated 2019)
33. Centers for Medicare & Medicaid Services. National Coverage Decision for Osteogenic Stimulators (150.2). 2005
34. UpToDate: General principles of definitive fracture management. Beutler A, Titus S. In: *UpToDate Online Journal* [serial online]. Waltham, MA: UpToDate; updated: Nov 17, 2022. Literature review current through Feb 2023
35. UpToDate: Overview of stress fractures. deWeber K. In: *UpToDate Online Journal* [serial online]. Waltham, MA: UpToDate; updated: Jun 23, 2022. Literature review current through Feb 2023
36. Nicholson JA, Tsang STJ, MacGillivray TJ, Perks F, Simpson AHRW. What is the role of ultrasound in fracture management?. *Bone Joint Res*. 2019;8(7):304-312. doi:10.1302/2046-3758.87.BJR-2018-0215.R2
37. McDaniel M, Eltman NR, Pan J, Swanson RL 2nd. Evaluation of Low-Intensity Pulsed Ultrasound on Stress Fractures to Reduce the Time to Return to Sport or Activity in the

MEDICAL POLICY

POLICY TITLE	LOW INTENSITY PULSED ULTRASOUND FRACTURE HEALING DEVICE
POLICY NUMBER	MP 6.021

Physically Active Population: A Systematic Review. Cureus. 2023;15(11):e49129. PMID 38024090

38. Palanisamy P, Alam M, Li S, Chow SKH, Zheng Y. Low-Intensity pulsed ultrasound stimulation for bone fractures healing. *Journal of Ultrasound in Medicine.* 2021;41(3):547-563. doi:10.1002/jum.15738
39. Bawale R, Segmeister M, Sinha S, Shariff S, Singh B. Experience of an isolated use of low-intensity pulsed ultrasound therapy on fracture healing in established non-unions: a prospective case series. *J Ultrasound.* 2021;24(3):249-252. PMID 32356220 PMCID PMC8363719
40. American Academy of Orthopaedic Surgeons, Board of Directors TAA of OS. *Treatment of Clavicle Fractures Evidence-Based Clinical Practice Guideline.*; 2022.
41. Blue Cross Blue Shield Association Medical Policy Reference Manual 1.01.05 *Ultrasound Accelerated Fracture Healing Device.* April, 2024

X. POLICY HISTORY

[Top](#)

MP 6.021	06/26/2020 Consensus Review. No change to policy statements, references updated.
	03/11/2021 Consensus Review. No change to policy statement. References updated.
	04/20/2022 Consensus Review. No change to policy statement. Coding table format updated. References reviewed and updated.
	03/29/2023 Consensus Review. No change to policy statements. Background and rationale updated. References reviewed and updated. Coding reviewed.
	06/04/2024 Consensus Review. No change to policy statement. References reviewed and updated. No coding changes.

[Top](#)

Health care benefit programs issued or administered by Capital Blue Cross and/or its subsidiaries, Capital Advantage Insurance Company®, Capital Advantage Assurance Company®, and Keystone Health Plan® Central. Independent licensees of the Blue Cross BlueShield Association. Communications issued by Capital Blue Cross in its capacity as administrator of programs and provider relations for all companies.