

MEDICAL POLICY

POLICY TITLE	TRIGGER POINT AND TENDER POINT INJECTIONS
POLICY NUMBER	MP 2.072

CLINICAL BENEFIT	☐ MINIMIZE SAFETY RISK OR CONCERN. ☒ MINIMIZE HARMFUL OR INEFFECTIVE INTERVENTIONS. ☐ ASSURE APPROPRIATE LEVEL OF CARE. ☒ ASSURE APPROPRIATE DURATION OF SERVICE FOR INTERVENTIONS. ☐ ASSURE THAT RECOMMENDED MEDICAL PREREQUISITES HAVE BEEN MET. ☐ ASSURE APPROPRIATE SITE OF TREATMENT OR SERVICE.
Effective date:	8/1/2026

POLICY

Trigger point injections with anesthetic and/or corticosteroid may be considered **medically necessary** for the treatment of myofascial pain syndrome when all of the following criteria have been met:

- There is a regional pain complaint in the expected distribution of referral pain from a trigger point, **AND**
- There is spot tenderness in a palpable taut band in a muscle, **AND**
- There is restricted range of motion, **AND**
- Conservative therapy (e.g., physical therapy, active exercises, ultrasound, heating or cooling, massage, activity modification, or pharmacotherapy) for 6 weeks fails or is not feasible, **AND**
- Trigger point injections are provided as a component of a comprehensive therapy program, **AND**
- No more than 4 injections total are given in a rolling 12-month period

Trigger point and tender point injections are considered **investigational** for all other indications including the treatment of myofascial pain syndrome not meeting the criteria above, complex regional pain syndrome, abdominal wall pain, and fibromyalgia. There is insufficient evidence to support a general conclusion concerning the health outcomes or benefits associated with this procedure.

Ultrasound and other imaging guidance of trigger point injections is considered **investigational** as there is insufficient evidence to support a general conclusion concerning the health outcomes or benefits associated with this procedure.

Cross-References:

MP 4.014 Epidural Steroid Injections for Back Pain

MP 4.050 Facet Joint Injections and Medial Branch Blocks

MP 4.041 Dry Needling of Trigger Points for Myofascial Pain

MP 8.012 Neural Therapy

MEDICAL POLICY

POLICY TITLE	TRIGGER POINT AND TENDER POINT INJECTIONS
POLICY NUMBER	MP 2.072

PRODUCT VARIATIONS

This policy is only applicable to certain programs and products administered by Capital Blue Cross and subject to benefit variations. Please see additional information below.

FEP PPO - Refer to FEP Medical Policy Manual. The FEP Medical Policy manual can be found at: <https://www.fepblue.org/benefit-plans/medical-policies-and-utilization-management-guidelines/medical-policies>.

DESCRIPTION/BACKGROUND

Trigger points are discrete, focal, hyperirritable spots within a taut band of skeletal muscle fibers that produce local and/or referred pain when stimulated. Tender points also produce local pain when stimulated but lack the taut band of tissue and hyperirritability when palpated. Injection of an anesthetic agent or botulinum toxin into trigger points and tender points is being evaluated for the management of a variety of pain syndromes.

Trigger Points

Trigger points are discrete, focal, hyperirritable spots within a taut band of skeletal muscle fibers that produce local and/or referred pain when stimulated. Trigger points are associated with local ischemia and hypoxia, a significantly lowered pH, local and referred pain, and altered muscle activation patterns.

Treatment

Trigger point injections with local anesthetic, saline, steroid, or botulinum toxin type A are a potential treatment for pain associated with trigger points. Alternative nonpharmacologic treatment modalities for trigger point pain include manual techniques, massage, acupressure, ultrasonography, application of heat or ice, diathermy, transcutaneous electrical nerve stimulation, and spray cooling with manual stretch.

Associated Disorders

Myofascial Pain Syndrome

Myofascial pain syndrome is a chronic regional pain disorder caused by the activation of at least one trigger point in muscles, tendons, or muscle fascia. It can cause local or referred pain, tightness, tenderness, stiffness and limitation of movement, muscle weakness, and often autonomic phenomena. The severity of symptoms and degree of functional impairment vary. Some individuals will have few trigger points with mild symptoms and no functional impairment, while others will have multiple satellite trigger points, widespread and severe pain, and major functional impairments. Conditions that can lead to myofascial pain syndrome include chronic repetitive minor muscle strain, poor posture, systemic disease, strain, sprain, enthesopathy, and arthritis. Management of chronic myofascial pain typically includes behavioral and pharmacologic approaches and physical therapy. Injection of a local anesthetic or botulinum toxin has also been reported.

MEDICAL POLICY

POLICY TITLE	TRIGGER POINT AND TENDER POINT INJECTIONS
POLICY NUMBER	MP 2.072

Complex Regional Pain Syndrome

Complex regional pain syndrome (previously called sympathetic dystrophy) refers to a chronic and disabling condition characterized by persistent pain that is disproportionate to the extent and duration of the primary injury and that is not restricted to the distribution of a specific peripheral nerve. Complex regional pain syndrome occurs most commonly following wrist fracture, but may follow many other types of injury, even when the preceding injury is relatively minor. Complex regional pain syndrome may also occur when there is no known injury. Complex regional pain syndrome is classified into type I when a specific nerve lesion has not been identified and type II when there is an identifiable nerve lesion. The pain may consist of thermal or mechanical allodynia (pain that occurs from a stimulus that normally does not elicit a painful response such as light touch or warmth) dysesthesia (a constant or ongoing unpleasant or electrical sensation of pain), and/or hyperalgesia (an exaggerated response to normally painful stimuli). Management of complex regional pain syndrome includes oral and topical pharmacotherapy, physical therapy, psychological therapies, and interventional procedures such as regional anesthetic blocks, sympathetic blocks, or spinal cord stimulation. Amputation of the affected limb has also been performed.

Abdominal Wall Pain

A source of chronic abdominal wall pain is anterior cutaneous nerve entrapment syndrome, which typically presents as sharp and focal abdominal pain, and is often found near a scar. One hypothesis is that anterior cutaneous nerve entrapment syndrome results from the entrapment and ischemia of an anterior cutaneous branch of a thoracic nerve as it passes through the rectus abdominus muscle. Anterior wall pain can be distinguished from intra-abdominal pain by documenting that pain increases with maneuvers that tense the abdominal muscles. It has also been proposed that abdominal wall pain may be due to a myofascial trigger point in the rectus abdominus muscle.

Tender Points

Tender points are focal areas of hyperalgesia that tend to occur at muscle-tendon junctions. Tender points are differentiated from trigger points due to the absence of a taut band of muscle tissue or local hyperirritability ("jump response") when palpated.

Despite the lack of local hyperirritability or a palpable band of tissue, some practitioners have treated tender points with injections of local anesthetic, corticosteroids, or botulinum toxin, similar to the treatment of trigger points.

Associated Disorders

Fibromyalgia

Fibromyalgia is a chronic condition characterized by widespread pain with hyperalgesia and allodynia. Constitutional symptoms such as fatigue, impaired cognition, and disrupted sleep can also occur. Early diagnostic criteria for fibromyalgia (1990) included three or more months of widespread pain above and below the waist, on both sides of the body, and along the midline, with at least 11 of 18 specific tender points. The defined bilateral areas from the American College

MEDICAL POLICY

POLICY TITLE	TRIGGER POINT AND TENDER POINT INJECTIONS
POLICY NUMBER	MP 2.072

of Rheumatology criteria are occipital, low cervical, trapezius, supraspinatus, second rib, lateral epicondyle, gluteal, greater trochanter, and knee medial fat pad. However, 2010 diagnostic criteria from the College, which were designed to facilitate diagnosis in a general practice setting, did not include a tender point exam but instead relied on the presence of widespread pain and other symptoms.

REGULATORY STATUS

Although medications used with invasive trigger point and tender point procedures are regulated by the U.S. Food and Drug Administration (FDA), trigger and tender point injections are procedures and, as such, are not subject to regulation by the FDA.

RATIONALE

Summary of Evidence

For individuals who have myofascial pain syndrome who receive trigger point injections, the evidence includes several randomized controlled trials (RCTs) and a systematic review of RCTs. Relevant outcomes are symptoms, functional outcomes, quality of life, and treatment-related morbidity. Lidocaine injections have been compared with physical therapy, lidocaine patches, sham stimulation, and dry needling. Some trials have reported that injecting lidocaine into trigger points improves subjective pain ratings to the same degree as physical therapy or lidocaine patches, but only slightly more than sham stimulation. Other trials have found that lidocaine injection was superior to dry needling on subjective pain ratings, but there was no significant benefit with lidocaine injection assessed on objective outcome measures. Randomized studies that compared bupivacaine and corticosteroid trigger point injections to standard therapies have also not found differences between groups. These results suggest a strong placebo effect of the treatment. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have complex regional pain syndrome who receive trigger point injections, the evidence includes case series. Relevant outcomes are symptoms, functional outcomes, quality of life, and treatment-related morbidity. Evidence on treatment of complex regional pain syndrome with trigger point injections is very limited, with only case series published and no recent literature identified for this treatment approach. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have abdominal wall pain who receive trigger point injections, the evidence includes RCT. Relevant outcomes are symptoms, functional outcomes, quality of life, and treatment-related morbidity. The single RCT evaluated lidocaine injections in women who had chronic pelvic pain and abdominal wall trigger points. Additional study in a larger population is needed to permit greater certainty about the efficacy of this treatment approach. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have fibromyalgia who receive tender point injections, the evidence includes an RCT. Relevant outcomes are symptoms, functional outcomes, quality of life, and treatment-related morbidity. The single RCT identified evaluated the efficacy of lidocaine injections in patients with fibromyalgia. It found a strong placebo effect, with lidocaine injection being not more

MEDICAL POLICY

POLICY TITLE	TRIGGER POINT AND TENDER POINT INJECTIONS
POLICY NUMBER	MP 2.072

effective than saline at reducing fibromyalgia pain. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

DEFINITIONS

N/A

DISCLAIMER

Capital Blue Cross' medical policies are used to determine coverage for specific medical technologies, procedures, equipment, and services. These medical policies do not constitute medical advice and are subject to change as permitted by law or applicable clinical evidence from independent treatment guidelines. Treating providers are solely responsible for medical advice and treatment of members. These policies are not a guarantee of coverage or payment. Payment of claims is subject to a determination regarding the member's benefit program and eligibility on the date of service, and a determination that the services are medically necessary and appropriate. Final processing of a claim is based upon the terms of contract that applies to the members' benefit program, including benefit limitations and exclusions. If a provider or a member has a question concerning this medical policy, please contact Capital Blue Cross' Provider Services or Member Services.

CODING INFORMATION

Note: This list of codes may not be all-inclusive, and codes are subject to change at any time. The identification of a code in this section does not denote coverage as coverage is determined by the terms of member benefit information. In addition, not all covered services are eligible for separate reimbursement.

Ultrasound and other imaging guidance of trigger point injections is considered investigational; therefore, not covered:

Procedure Codes								
76942	77002	77012	77021					

A maximum of four (4) injections (total) within a rolling 12-month period are covered when medically necessary:

Procedure Codes								
20552	20553							

ICD-10-CM Diagnosis Code	Description
M79.10	Myalgia, unspecified site
M79.12	Myalgia of auxiliary muscles, head, and neck
M79.18	Myalgia, other site

MEDICAL POLICY

POLICY TITLE	TRIGGER POINT AND TENDER POINT INJECTIONS
POLICY NUMBER	MP 2.072

REFERENCES

1. Alvarez DJ, Rockwell PG. Trigger points: diagnosis and management. *Am Fam Physician*. Feb 15 2002; 65(4): 653-60. PMID 11871683
2. O'Connell NE, Wand BM, McAuley J, et al. Interventions for treating pain and disability in adults with complex regional pain syndrome. *Cochrane Database Syst Rev*. Apr 30 2013; 2013(4): CD009416. PMID 23633371
3. Alnahhas MF, Oxentenko SC, Locke GR, et al. Outcomes of Ultrasound-Guided Trigger Point Injection for Abdominal Wall Pain. *Dig Dis Sci*. Feb 2016; 61(2): 572-7. PMID 26320087
4. Sumpton JE, Moulin DE. Fibromyalgia. *Handb Clin Neurol*. 2014; 119: 513-27. PMID 24365316
5. Wolfe F, Clauw DJ, Fitzcharles MA, et al. The American College of Rheumatology preliminary diagnostic criteria for fibromyalgia and measurement of symptom severity. *Arthritis Care Res (Hoboken)*. May 2010; 62(5): 600-10. PMID 20461783
6. Lugo LH, García HI, Rogers HL, et al. Treatment of myofascial pain syndrome with lidocaine injection and physical therapy, alone or in combination: a single blind, randomized, controlled clinical trial. *BMC Musculoskelet Disord*. Feb 24 2016; 17: 101. PMID 26911981
7. Affaitati G, Fabrizio A, Savini A, et al. A randomized, controlled study comparing a lidocaine patch, a placebo patch, and anesthetic injection for treatment of trigger points in patients with myofascial pain syndrome: evaluation of pain and somatic pain thresholds. *Clin Ther*. Apr 2009; 31(4): 705-20. PMID 19446144
8. Couto C, de Souza IC, Torres IL, et al. Paraspinal stimulation combined with trigger point needling and needle rotation for the treatment of myofascial pain: a randomized sham-controlled clinical trial. *Clin J Pain*. Mar 2014; 30(3): 214-23. PMID 23629597
9. Hong CZ. Lidocaine injection versus dry needling to myofascial trigger point. The importance of the local twitch response. *Am J Phys Med Rehabil*. 1994; 73(4): 256-63. PMID 8043247
10. Lajeunesse MD, Olivera TR, April MD, et al. Trigger Point Injection for Myofascial Pain Syndrome of the Low Back: A Partially Blinded Three-Arm Randomized Controlled Trial. *Ann Emerg Med*. Dec 2025; 86(6): 606-615. PMID 40531073
11. Brennan KL, Allen BC, Maldonado YM. Dry Needling Versus Cortisone Injection in the Treatment of Greater Trochanteric Pain Syndrome: A Noninferiority Randomized Clinical Trial. *J Orthop Sports Phys Ther*. Apr 2017; 47(4): 232-239. PMID 28257614
12. Hong CZ. Specific sequential myofascial trigger point therapy in the treatment of a patient with myofascial pain syndrome associated with reflex sympathetic dystrophy. *Australas Chiropr Osteopathy*. Mar 2000; 9(1): 7-11. PMID 17987165
13. Dunn DG. Chronic regional pain syndrome, type 1: Part II. *AORN J*. Oct 2000; 72(4): 643-51, 653; quiz 654, 656-8, 661-2. PMID 11076283
14. Ferraro MC, Cashin AG, Wand BM, et al. Interventions for treating pain and disability in adults with complex regional pain syndrome- an overview of systematic reviews. *Cochrane Database Syst Rev*. Jun 12 2023; 6(6): CD009416. PMID 37306570
15. Oor JE, Ünlü Ç, Hazebroek EJ. A systematic review of the treatment for abdominal cutaneous nerve entrapment syndrome. *Am J Surg*. Jul 2016; 212(1): 165-74. PMID 26945611

MEDICAL POLICY

POLICY TITLE	TRIGGER POINT AND TENDER POINT INJECTIONS
POLICY NUMBER	MP 2.072

16. Montenegro ML, Braz CA, Rosa-e-Silva JC, et al. Anaesthetic injection versus ischemic compression for the pain relief of abdominal wall trigger points in women with chronic pelvic pain. *BMC Anesthesiol.* Dec 01 2015; 15: 175. PMID 26628263
17. Staud R, Weyl EE, Bartley E, et al. Analgesic and anti-hyperalgesic effects of muscle injections with lidocaine or saline in patients with fibromyalgia syndrome. *Eur J Pain.* Jul 2014; 18(6): 803-12. PMID 24193993
18. Benzon HT, Connis RT, De Leon-Casasola OA, et al. Practice guidelines for chronic pain management: an updated report by the American Society of Anesthesiologists Task Force on Chronic Pain Management and the American Society of Regional Anesthesia and Pain Medicine. *Anesthesiology.* Apr 2010; 112(4): 810-33. PMID 20124882
19. Benzon HT, Elmofly D, Shankar H, et al. Use of corticosteroids for adult chronic pain interventions: sympathetic and peripheral nerve blocks, trigger point injections - guidelines from the American Society of Regional Anesthesia and Pain Medicine, the American Academy of Pain Medicine, the American Society of Interventional Pain Physicians, and the International Pain and Spine Intervention Society. *Reg Anesth Pain Med.* Mar 12 2025. PMID 39019502
20. National Institute for Health and Care Excellence. Chronic pain (primary and secondary) in over 16s: assessment of all chronic pain and management of chronic primary pain. *NICE Guideline NG193.* April 2021 (amended April 2022).

POLICY HISTORY

MP 2.072	04/29/2020 Consensus Review. No change to policy statement. References checked and updated. Coding reviewed with no changes.
	09/07/2021 Administrative Update. Addition of new ICD-10 codes. Effective date 10/01/2021.
	12/08/2021 Minor Review. Removed fibromyalgia as a medically necessary indication. FEP statement updated. Rationale and coding updated. References updated.
	07/07/2022 Minor Review. Tender Point injections changed from medically necessary to investigational. Added code 77012 as not covered. Background, Rationale and References updated.
	06/07/2023 Minor Review. Added "rolling" to the 12-month time period. Also added that it is 4 injections total. Background and References updated.
	05/28/2024 Consensus Review. No change to policy statement. Removed retired MP 3.015 Office Based Procedures Performed in a Facility from Cross Referenced policies section. References updated.
	05/30/2025 Minor Review. Chronic neck and back pain changed from medically necessary to investigational. Removed ICD10 codes G89.29, G89.4, M54.2, M54.50, M54.51 and M54.59. References added.
	09/10/2025 Administrative Update. Removed Benefit Variations Section and updated Disclaimer.
	04/09/2026 Consensus Review. No change to policy statement. Updated cross-references, rationale, and references. Coding reviewed.

MEDICAL POLICY

POLICY TITLE	TRIGGER POINT AND TENDER POINT INJECTIONS
POLICY NUMBER	MP 2.072

Health care benefit programs issued or administered by Capital Blue Cross and/or its subsidiaries, Capital Advantage Insurance Company[®], Capital Advantage Assurance Company[®] and Keystone Health Plan[®] Central. Independent licensees of the Blue Cross BlueShield Association. Communications issued by Capital Blue Cross in its capacity as administrator of programs and provider relations for all companies.