

MEDICAL POLICY

POLICY TITLE	CARDIAC HEMODYNAMIC MONITORING FOR THE MANAGEMENT OF HEART FAILURE IN THE OUTPATIENT SETTING
POLICY NUMBER	MP 2.051

CLINICAL BENEFIT	<input type="checkbox"/> MINIMIZE SAFETY RISK OR CONCERN. <input checked="" type="checkbox"/> MINIMIZE HARMFUL OR INEFFECTIVE INTERVENTIONS. <input type="checkbox"/> ASSURE APPROPRIATE LEVEL OF CARE. <input type="checkbox"/> ASSURE APPROPRIATE DURATION OF SERVICE FOR INTERVENTIONS. <input type="checkbox"/> ASSURE THAT RECOMMENDED MEDICAL PREREQUISITES HAVE BEEN MET. <input type="checkbox"/> ASSURE APPROPRIATE SITE OF TREATMENT OR SERVICE.
Effective Date:	3/1/2024

[POLICY RATIONALE](#)
[DISCLAIMER](#)
[POLICY HISTORY](#)

[PRODUCT VARIATIONS](#)
[DEFINITIONS](#)
[CODING INFORMATION](#)

[DESCRIPTION/BACKGROUND](#)
[BENEFIT VARIATIONS](#)
[REFERENCES](#)

I. POLICY

In the ambulatory care and outpatient setting, cardiac hemodynamic monitoring for the management of heart failure using implantable direct pressure monitoring of the pulmonary artery, thoracic bioimpedance, inert gas rebreathing, and arterial pressure during the Valsalva maneuver is considered **investigational**. There is insufficient evidence to support a general conclusion concerning the health outcomes or benefits associated with these procedures.

Policy Guidelines

This policy refers only to the use of stand-alone cardiac output measurement devices designed for use in ambulatory care and outpatient settings. The use of cardiac hemodynamic monitors or intrathoracic fluid monitors that are integrated into other implantable cardiac devices, including implantable cardioverter defibrillators, cardiac resynchronization therapy devices, and cardiac pacing devices, is addressed in MP 2.007.

Cross-reference:

MP 2.007 Cardiac Intervention Therapy for the Treatment of Heart Failure

II. PRODUCT VARIATIONS

[Top](#)

This policy is only applicable to certain programs and products administered by Capital Blue Cross and subject to benefit variations as discussed in Section VI. Please see additional information below.

FEP PPO - Refer to FEP Medical Policy Manual. The FEP Medical Policy manual can be found at:

<https://www.fepblue.org/benefit-plans/medical-policies-and-utilization-management-guidelines/medical-policies>.

MEDICAL POLICY

POLICY TITLE	CARDIAC HEMODYNAMIC MONITORING FOR THE MANAGEMENT OF HEART FAILURE IN THE OUTPATIENT SETTING
POLICY NUMBER	MP 2.051

III. DESCRIPTION/BACKGROUND

[Top](#)

A variety of outpatient cardiac hemodynamic monitoring devices are intended to improve quality of life and reduce morbidity for patients with heart failure by decreasing episodes of acute decompensation. Monitors can identify physiologic changes that precede clinical symptoms and thus allow preventive intervention. These devices operate through various mechanisms, including implantable pressure sensors, thoracic bioimpedance measurement, inert gas rebreathing, and estimation of left ventricular end diastolic pressure by arterial pressure during the Valsalva maneuver.

Chronic Heart Failure

Patients with chronic heart failure are at risk of developing acute decompensated heart failure, often requiring hospital admission. Patients with a history of acute decompensation have the additional risk of future episodes of decompensation, and death. Reasons for the transition from a stable, chronic state to an acute, decompensated state include disease progression, as well as acute events such as coronary ischemia and dysrhythmias. While precipitating factors are frequently not identified, the most common preventable cause is noncompliance with medication and dietary regimens.

Management

Strategies for reducing decompensation, and thus the need for hospitalization, are aimed at early identification of patients at risk for imminent decompensation. Programs for early identification of heart failure are characterized by frequent contact with patients to review signs and symptoms with a health care provider, education, and medication adjustments as appropriate. These encounters may occur face-to-face in the office or at home, or via cellular or computed technology.

Precise measurement of cardiac hemodynamics is often employed in the intensive care setting to carefully manage fluid status in acutely decompensated heart failure. Transthoracic echocardiography, transesophageal echocardiography, and Doppler ultrasound are noninvasive methods for monitoring cardiac output on an intermittent basis for the more stable patient but are not addressed herein. A variety of biomarkers and radiologic techniques may be used for dyspnea when the diagnosis of acute decompensated heart failure is uncertain.

The criterion standard for hemodynamic monitoring is pulmonary artery catheters and central venous pressure catheters. However, they are invasive, inaccurate, and inconsistent in predicting fluid responsiveness. Several studies have demonstrated that catheters fail to improve outcomes in critically ill patients and may be associated with harm. To overcome these limitations, multiple techniques and devices have been developed that use complex imaging technology and computer algorithms to estimate fluid responsiveness, volume status, cardiac output, and tissue perfusion. Many are intended for use in outpatient settings but can be used in the emergency department, intensive care unit, and operating room. Four methods are reviewed here: implantable pressure monitoring devices, thoracic bioimpedance, inert gas rebreathing, and arterial waveform during the Valsalva maneuver. Use of last three is not widespread

MEDICAL POLICY

POLICY TITLE	CARDIAC HEMODYNAMIC MONITORING FOR THE MANAGEMENT OF HEART FAILURE IN THE OUTPATIENT SETTING
POLICY NUMBER	MP 2.051

because of several limitations including use proprietary technology making it difficult to confirm their validity and lack of large randomized controlled trials to evaluate treatment decisions guided by these hemodynamic monitors.

Regulatory Status

Noninvasive Left Ventricular End-Diastolic Pressure Measurement Devices

In 2004, the VeriCor® (CVP Diagnostics), a noninvasive left ventricular end-diastolic pressure measurement device, was cleared for marketing by U.S. Food and Drug Administration (FDA) through the 510(k) process. The FDA determined that this device was substantially equivalent to existing devices for the following indication:

“The VeriCor is indicated for use in estimating non-invasively, left ventricular end-diastolic pressure (LVEDP). This estimate, when used along with clinical signs and symptoms and other patient test results, including weights on a daily basis, can aid the clinician in the selection of further diagnostic tests in the process of reaching a diagnosis and formulating a therapeutic plan when abnormalities of intravascular volume are suspected. The device has been clinically validated in males only. Use of the device in females has not been investigated.”

FDA product code: DXN.

Thoracic Bioimpedance Devices

Multiple thoracic impedance measurement devices that do not require invasive placement have been cleared for marketing by the FDA through the 510(k) process. The FDA determined that this device was substantially equivalent to existing devices used for peripheral blood flow monitoring. Table 1 presents an inexhaustive list of representative devices (FDA product code: DSB).

Table 1. Noninvasive Thoracic Impedance Plethysmography Devices

Device	Manufacturer	Clearance Date
BioZ® Thoracic Impedance Plethysmograph	SonoSite	2009
Zoe® Fluid Status Monitor	Noninvasive Medical Technologies	2004
Cheetah Starling SV	Cheetah Medical	2008
PhysioFlow® Signal Morphology-based Impedance Cardiography (SM-ICG™)	Vasocom, now NeuMeDx	2008
ReDS™ Wearable System	Sensible Medical Innovations	2015
Bodyport Cardiac Scale	Bodyport Inc.	2022

MEDICAL POLICY

POLICY TITLE	CARDIAC HEMODYNAMIC MONITORING FOR THE MANAGEMENT OF HEART FAILURE IN THE OUTPATIENT SETTING
POLICY NUMBER	MP 2.051

Also, several manufacturers market thoracic impedance measurement devices integrated into implantable cardiac pacemakers, cardioverter defibrillator devices, and cardiac resynchronization therapy devices. Thoracic bioimpedance devices integrated into implantable cardiac devices are addressed in evidence review MP 2.007.

Inert Gas Rebreathing Devices

In 2006, the Innocor® (Innovision), an inert gas rebreathing device, was cleared for marketing by the FDA through the 510(k) process. The FDA determined that this device was substantially equivalent to existing inert gas rebreathing devices for use in computing blood flow. FDA product code: BZG.

Implantable Pulmonary Artery Pressure Sensor Devices

In 2014, the CardioMEMS™ Heart Failure Monitoring System (CardioMEMS, now Abbott) was approved for marketing by the FDA through the premarket approval process. This device consists of an implantable pulmonary artery (PA) sensor, which is implanted in the distal PA, a transvenous delivery system, and an electronic sensor that processes signals from the implantable PA sensor and transmits PA pressure measurements to a secure database. The device originally underwent FDA review in 2011, at which point FDA found no reasonable assurance that the monitoring system would be effective, particularly in certain subpopulations, although the FDA agreed this monitoring system was safe for use in the indicated patient population. In 2022, the CardioMEMS™ HF Monitoring System received expanded approval for the treatment of New York Heart Association (NYHA) Class II-III patients who had been hospitalized at least 1 time in the prior year and/or had elevated natriuretic peptides.

Several other devices that monitor cardiac output by measuring pressure changes in the PA or right ventricular outflow tract have been investigated in the research setting but have not received the FDA approval. They include the Chronicle® implantable continuous hemodynamic monitoring device (Medtronic), which includes a sensor implanted in the right ventricular outflow tract, the ImPressure® device (Remon Medical Technologies), which includes a sensor implanted in the PA, and the Cordella™ PA Pressure Sensor System (Endotronix, Inc.), which includes a sensor implanted in the PA.

Note: This evidence review only addresses the use of these technologies in ambulatory care and outpatient settings.

IV. RATIONALE

[Top](#)

Summary of Evidence

For individuals with New York Heart Association (NYHA) class II-IV heart failure in outpatient settings who have had a hospitalization in the past year and/or have elevated natriuretic peptides who receive hemodynamic monitoring with an implantable pulmonary artery pressure sensor device, the evidence includes randomized controlled trials (RCTs) and nonrandomized studies. Relevant outcomes are overall survival, symptoms, functional outcomes, quality of life,

MEDICAL POLICY

POLICY TITLE	CARDIAC HEMODYNAMIC MONITORING FOR THE MANAGEMENT OF HEART FAILURE IN THE OUTPATIENT SETTING
POLICY NUMBER	MP 2.051

morbid events, hospitalizations, and treatment-related morbidity. One implantable pressure monitor, the CardioMEMS device, has U.S. Food and drug Administration (FDA) approval. The pivotal CHAMPION RCT reported a statistically significant 28% decrease in heart failure hospitalization (HFH) in patients implanted with the CardioMEMS device compared with usual care. However, trial results were potentially biased in favor of the treatment group due to the use of additional nurse communication to enhance protocol compliance with the device. The manufacturer conducted multiple analyses to address potential bias from the nurse interventions. Results were reviewed favorably by the FDA. While these analyses demonstrated the consistency of benefit of the CardioMEMS device, all such analyses have methodologic limitations. Early safety data have been suggestive of a higher rate of procedural complications, particularly related to pulmonary artery injury. While the U.S. CardioMEMS post-approval study and CardioMEMS European Monitoring Study for Heart Failure (MEMS-HF) study reported a significant decrease in HFH with few device- or system-related complications at 1 year, the impact of nursing interventions remains unclear. The subsequent GUIDE-HF RCT failed to meet its primary efficacy endpoint, the composite of HFH, urgent heart failure visits, and death at 1 year. With the approval of the FDA, the statistical analysis plan was updated to pre-specify sensitivity analyses to assess the impact of COVID-19 on the trial. For the 72% of patients who completed follow-up prior to the public health emergency declaration in March 2020, a statistically significant 19% reduction in the primary endpoint was reported, driven by a 28% reduction in HFH. However, lifestyle changes during the COVID-19 pandemic such as changes in physical activity, exposure to infections, willingness to seek medical care, and adherence to medications are unmeasured and add imprecision to treatment effect estimates, as do alterations in provider behaviors. Enrollment of NYHA Class II patients was significantly enriched in the first 500 patients, potentially impacting the pre-COVID-19 analysis. Overall, the beneficial effect of CardioMEMS, if any, appears to be on the hospitalization outcome of the composite. Both urgent heart failure visits and death outcomes had hazard ratios favoring the control group with wide confidence intervals including the null value in pre-COVID-19, during-COVID-19, and overall analyses of the GUIDE-HF trial. No significant differences were observed in secondary quality of life and functional status outcomes. While the HFH reduction of 28% found in the pre-COVID-19 analysis is consistent with findings from the CHAMPION trial, it is unclear whether physician knowledge of treatment assignment biases the decision to hospitalize and administer intravenous diuretics. Given that the intervention is invasive and intended to be used for a highly prevalent condition and, in light of the absence of a demonstrated benefit on mortality and functional outcomes, the lack of periprocedural safety data, and unclear impact of COVID-19 on remote monitoring in the GUIDE-HF trial, the net benefit of the CardioMEMS device remains uncertain. Concerns may be clarified by the ongoing open access phase of the GUIDE-HF RCT and the German non-industry-sponsored PASSPORT-HF trial. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have heart failure in outpatient settings who receive hemodynamic monitoring by thoracic bioimpedance, the evidence includes uncontrolled prospective studies and case series. Relevant outcomes are overall survival, symptoms, functional outcomes, quality of life, morbid events, hospitalizations, and treatment-related morbidity. There is a lack of RCT evidence evaluating whether the use of these technologies improves health outcomes over standard active management of heart failure patients. The case series have reported physiologic measurement-

MEDICAL POLICY

POLICY TITLE	CARDIAC HEMODYNAMIC MONITORING FOR THE MANAGEMENT OF HEART FAILURE IN THE OUTPATIENT SETTING
POLICY NUMBER	MP 2.051

related outcomes and/or associations between monitoring information and heart failure exacerbations, but do not provide definitive evidence on device efficacy. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have heart failure in outpatient settings who receive hemodynamic monitoring with inert gas rebreathing, no studies have been identified on clinical validity or clinical utility. Relevant outcomes are overall survival, symptoms, functional outcomes, quality of life, morbid events, hospitalizations, and treatment-related morbidity. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who have heart failure in outpatient settings who receive hemodynamic monitoring of arterial pressure during the Valsalva maneuver, a single study was identified. Relevant outcomes are overall survival, symptoms, functional outcomes, quality of life, morbid events, hospitalizations, and treatment-related morbidity. The study assessed the use of left ventricular end-diastolic pressure (LVEDP) monitoring and reported an 85% sensitivity and an 80% specificity to detect LVEDP greater than 15 mm Hg. The evidence is insufficient to determine that the technology results in an improvement in the net health outcome.

V. DEFINITIONS

[Top](#)

CARDIAC OUTPUT is the volume of blood expelled by the ventricles of the heart, equal to the amount of blood ejected at each beat (the stroke output) multiplied by the heart rate per minute/number of beats in the period of time used in the computation. A normal heart in a resting adult ejects from four (4) to eight (8) liters of blood per minute.

HEMODYNAMIC refers to the study of the forces involved in circulating blood through the body.

THORAX is the cage of bone and cartilage containing the principal organs of respiration and circulation and covering part of the abdominal organs.

VALSALVA'S MANEUVER is an attempt to forcibly exhale with the glottis, nose and mouth closed. This maneuver causes increased intrathoracic pressure, slowing of the pulse, decreased return of blood to the heart, and increased venous pressure. If the Eustachian tubes are not obstructed, the pressure on the tympanic membranes also will be increased. When this maneuver is done with just the glottis closed, only intrathoracic pressure will increase.

VI. BENEFIT VARIATIONS

[Top](#)

The existence of this medical policy does not mean that this service is a covered benefit under the member's health benefit plan. Benefit determinations should be based in all cases on the applicable health benefit plan language. Medical policies do not constitute a description of benefits. A member's health benefit plan governs which services are covered, which are excluded, which are subject to benefit limits, and which require preauthorization. There are different benefit plan designs in each product administered by Capital Blue Cross. Members and providers should consult the member's health benefit plan for information or contact Capital Blue Cross for benefit information.

MEDICAL POLICY

POLICY TITLE	CARDIAC HEMODYNAMIC MONITORING FOR THE MANAGEMENT OF HEART FAILURE IN THE OUTPATIENT SETTING
POLICY NUMBER	MP 2.051

VII. DISCLAIMER

[Top](#)

Capital Blue Cross's medical policies are developed to assist in administering a member's benefits, do not constitute medical advice and are subject to change. Treating providers are solely responsible for medical advice and treatment of members. Members should discuss any medical policy related to their coverage or condition with their provider and consult their benefit information to determine if the service is covered. If there is a discrepancy between this medical policy and a member's benefit information, the benefit information will govern. If a provider or a member has a question concerning the application of this medical policy to a specific member's plan of benefits, please contact Capital Blue Cross' Provider Services or Member Services. Capital Blue Cross considers the information contained in this medical policy to be proprietary and it may only be disseminated as permitted by law.

VIII. CODING INFORMATION

[Top](#)

Note: This list of codes may not be all-inclusive, and codes are subject to change at any time. The identification of a code in this section does not denote coverage as coverage is determined by the terms of member benefit information. In addition, not all covered services are eligible for separate reimbursement.

Investigational; therefore, not covered:

Procedure Codes							
0607T	0608T	33289	93050	93264	93701	C2624	

The following code is investigational when used for ambulatory care and outpatient setting. Inert Gas Rebreathing as outlined in the policy section:

Procedure Codes							
93799							

IX. REFERENCES

[Top](#)

1. *Opasich C, Rapezzi C, Lucci D, et al. Precipitating factors and decision-making processes of short-term worsening heart failure despite optimal treatment (from the IN-CHF Registry). Am J Cardiol. Aug 15, 2001;88(4):382-387. PMID 11545758*
2. *McAlister FA, Stewart S, Ferrua S, et al. Multidisciplinary strategies for the management of heart failure patients at high risk for admission: a systematic review of randomized trials. J Am Coll Cardiol. Aug 18, 2004;44(4):810- 819. PMID 15312864*
3. *Food and Drug Administration. Summary of Safety and Effectiveness Data (SSED): CardioMEMS HF System. 2014*
4. *Loh JP, Barbash IM, Waksman R. Overview of the 2011 Food and Drug Administration Circulatory System Devices Panel of the Medical Devices Advisory Committee Meeting on the CardioMEMS Champion Heart Failure Monitoring System. J Am Coll Cardiol. Apr 16, 2013;61(15):1571-1576. PMID 23352783*

MEDICAL POLICY

POLICY TITLE	CARDIAC HEMODYNAMIC MONITORING FOR THE MANAGEMENT OF HEART FAILURE IN THE OUTPATIENT SETTING
POLICY NUMBER	MP 2.051

5. Burns DJP, Arora J, Okunade O, et al. International Consortium for Health Outcomes Measurement (ICHOM): Standardized Patient-Centered Outcomes Measurement Set for Heart Failure Patients. *JACC Heart Fail.* Mar 2020; 8(3): 212-222. PMID 31838032
6. Zannad F, Garcia AA, Anker SD, et al. Clinical outcome endpoints in heart failure trials: a European Society of Cardiology Heart Failure Association consensus document. *Eur. J. Heart Fail.*, 2013 Jun 22;15(10). PMID 23787718
7. Givertz MM, Stevenson LW, Costanzo MR, et al. Pulmonary Artery Pressure-Guided Management of Patients With Heart Failure and Reduced Ejection Fraction. *J Am Coll Cardiol.* Oct 10, 2017; 70(15): 1875-1886. PMID 28982501
8. Curtain JP, Lee MMY, McMurray JJ, et al. Efficacy of implantable haemodynamic monitoring in heart failure across ranges of ejection fraction: a systematic review and meta-analysis. *Heart.* May 15 2023; 109(11): 823-831. PMID 36522146
9. Adamson PB, Abraham WT, Bourge RC, et al. Wireless pulmonary artery pressure monitoring guides management to reduce decompensation in heart failure with preserved ejection fraction. *Circ Heart Fail.* Nov 2014;7(6):935-944. PMID 25286913
10. Adamson PB, Abraham WT, Stevenson LW, et al. Pulmonary Artery Pressure-Guided Heart Failure Management Reduces 30-Day Readmissions. *Circ Heart Fail.* Jun 2016;9(6). PMID 27220593
11. Krahnke JS, Abraham WT, Adamson PB, et al. Heart failure and respiratory hospitalizations are reduced in patients with heart failure and chronic obstructive pulmonary disease with the use of an implantable pulmonary artery pressure monitoring device. *J Card Fail.* Mar 2015;21(3):240-249. PMID 25541376
12. Assmus B, Angermann CE, Alkhout B, et al. Effects of remote haemodynamic-guided heart failure management in patients with different subtypes of pulmonary hypertension: insights from the MEMS-HF study. *Eur J Heart Fail.* Dec 2022; 24(12): 2320-2330. PMID 36054647
13. Heywood JT, Jermyn R, Shavelle D, et al. Impact of Practice-Based Management of Pulmonary Artery Pressures in 2000 Patients Implanted With the CardioMEMS Sensor. *Circulation.* Apr 18, 2017; 135(16): 1509-1517. PMID 28219895
14. Abraham WT, Adamson PB, Bourge RC, et al. Wireless pulmonary artery haemodynamic monitoring in chronic heart failure: a randomised controlled trial. *Lancet.* Feb 19, 2011; 377(9766): 658-66. PMID 21315441
15. Abraham WT, Stevenson LW, Bourge RC, et al. Sustained efficacy of pulmonary artery pressure to guide adjustment of chronic heart failure therapy: complete follow-up results from the CHAMPION randomised trial. *Lancet.* Jan 30, 2016; 387(10017): 453-61. PMID 26560249
16. Adamson PB, Abraham WT, Aaron M, et al. CHAMPION trial rationale and design: the long-term safety and clinical efficacy of a wireless pulmonary artery pressure monitoring system. *J Card Fail.* Jan 2011; 17(1): 3-10. PMID 21187258
17. CardioMEMS Champion™ HF Monitoring System: FDA Review of P100045/A004FDA Presentation - CardioMEMS: Oct. 9, 2013. 2013
18. CardioMEMS Champion™ Heart Failure Monitoring System: Presentation - CardioMEMS: Oct. 9, 2013. 2013

MEDICAL POLICY

POLICY TITLE	CARDIAC HEMODYNAMIC MONITORING FOR THE MANAGEMENT OF HEART FAILURE IN THE OUTPATIENT SETTING
POLICY NUMBER	MP 2.051

19. Shavelle DM, Desai AS, Abraham WT, et al. Lower Rates of Heart Failure and All-Cause Hospitalizations During Pulmonary Artery Pressure-Guided Therapy for Ambulatory Heart Failure: One-Year Outcomes From the CardioMEMS Post-Approval Study. *Circ Heart Fail.* Aug 2020; 13(8): e006863. PMID 32757642
20. DeFilippis EM, Henderson J, Axsom KM, et al. Remote Hemodynamic Monitoring Equally Reduces Heart Failure Hospitalizations in Women and Men in Clinical Practice: A Sex-Specific Analysis of the CardioMEMS Post-Approval Study. *Circ Heart Fail.* Jun 2021; 14(6): e007892. PMID 34129363
21. Lindenfeld J, Zile MR, Desai AS, et al. Haemodynamic-guided management of heart failure (GUIDE-HF): a randomised controlled trial. *Lancet.* Sep 11, 2021; 398(10304): 991-1001. PMID 34461042
22. Zile MR, Desai AS, Costanzo MR, et al. The GUIDE-HF trial of pulmonary artery pressure monitoring in heart failure: impact of the COVID-19 pandemic. *Eur Heart J.* Mar 10, 2022. PMID 35266003
23. Kishino Y, Kuno T, Malik AH, et al. Effect of pulmonary artery pressure-guided therapy on heart failure readmission in a nationally representative cohort. *ESC Heart Fail.* Aug 2022; 9(4): 2511-2517. PMID 35560987
24. Cowie MR, Flett A, Cowburn P, et al. Real-world evidence in a national health service: results of the UK CardioMEMS HF System Post-Market Study. *ESC Heart Fail.* Feb 2022; 9(1): 48-56. PMID 34882989
25. Angermann CE, Assmus B, Anker SD, et al. Pulmonary artery pressure-guided therapy in ambulatory patients with symptomatic heart failure: the CardioMEMS European Monitoring Study for Heart Failure (MEMS-HF). *Eur J Heart Fail.* Oct 2020; 22(10): 1891-1901. PMID 32592227
26. Abraham J, Bharmi R, Jonsson O, et al. Association of Ambulatory Hemodynamic Monitoring of Heart Failure With Clinical Outcomes in a Concurrent Matched Cohort Analysis. *JAMA Cardiol.* Jun 01, 2019; 4(6): 556-563. PMID 31090869
27. Desai AS, Bhimaraj A, Bharmi R, et al. Ambulatory Hemodynamic Monitoring Reduces Heart Failure Hospitalizations in Real-World Clinical Practice. *J Am Coll Cardiol.* May 16, 2017;69(19):2357-2365. PMID 28330751
28. Lin AL, Hu G, Dhruva SS, et al. Quantification of Device-Related Event Reports Associated With the CardioMEMS Heart Failure System. *Circ Cardiovasc Qual Outcomes.* Oct 2022; 15(10): e009116. PMID 36252112
29. Vaduganathan M, DeFilippis EM, Fonarow GC, et al. Postmarketing adverse events related to the CardioMEMS HF System. *JAMA Cardiol.* Nov 1, 2017;2(11):1277-1279. PMID 28975249
30. Krzesiński P, Jankowska EA, Siebert J, et al. Effects of an outpatient intervention comprising nurse-led non-invasive assessments, telemedicine support and remote cardiologists' decisions in patients with heart failure (AMULET study): a randomized controlled trial. *Eur J Heart Fail.* Mar 2022; 24(3): 565-577. PMID 34617373
31. Kamath SA, Drazner MH, Tasissa G, et al. Correlation of impedance cardiography with invasive hemodynamic measurements in patients with advanced heart failure: the BioImpedance CardioGraphy (BIG) substudy of the Evaluation Study of Congestive Heart

MEDICAL POLICY

POLICY TITLE	CARDIAC HEMODYNAMIC MONITORING FOR THE MANAGEMENT OF HEART FAILURE IN THE OUTPATIENT SETTING
POLICY NUMBER	MP 2.051

Failure and Pulmonary Artery Catheterization Effectiveness (ESCAPE) Trial. Am Heart J. Aug 2009;158(2):217-223. PMID 19619697

32. *Anand IS, Greenberg BH, Fogoros RN, et al. Design of the Multi-Sensor Monitoring in Congestive Heart Failure (MUSIC) study: prospective trial to assess the utility of continuous wireless physiologic monitoring in heart failure. J Card Fail. Jan 2011;17(1):11-16. PMID 21187259*
33. *Anand IS, Tang WH, Greenberg BH, et al. Design and performance of a multisensor heart failure monitoring algorithm: results from the multisensor monitoring in congestive heart failure (MUSIC) study. J Card Fail. Apr 2012;18(4):289-295. PMID 22464769*
34. *Packer M, Abraham WT, Mehra MR, et al. Utility of impedance cardiography for the identification of short-term risk of clinical decompensation in stable patients with chronic heart failure. J Am Coll Cardiol. Jun 6, 2006;47(11):2245-2252. PMID 16750691*
35. *Amir O, Ben-Gal T, Weinstein JM, et al. Evaluation of remote dielectric sensing (ReDS) technology-guided therapy for decreasing heart failure re-hospitalizations. Int J Cardiol. Aug 1, 2017;240:279-284. PMID 28341372*
36. *Silber HA, Trost JC, Johnston PV, et al. Finger photoplethysmography during the Valsalva maneuver reflects left ventricular filling pressure. Am J Physiol Heart Circ Physiol. May 2012;302(10):H2043-2047. PMID 22389389*
37. *Yancy CW, Jessup M, Bozkurt B, et al. 2017 ACC/AHA/HFSA Focused Update of the 2013 ACCF/AHA Guideline for the Management of Heart Failure: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines and the Heart Failure Society of America. J Am Coll Cardiol. Aug 8, 2017;70(6):776-803. PMID 28461007*
38. *Heidenreich PA, Bozkurt B, Aguilar D, et al. 2022 AHA/ACC/HFSA Guideline for the Management of Heart Failure: A Report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines. Circulation. May 03, 2022; 145(18): e895-e1032. PMID 35363499*
39. *National Institute for Health and Care Excellence (NICE). Chronic heart failure in adults: diagnosis and management [NG106]. September 2018*
40. *Dickinson, MM, Allen, LL, Albert, NN, DiSalvo, TT, Ewald, GG, Vest, AA, Whellan, DD, Zile, MM, Givertz, MM. Remote Monitoring of Patients With Heart Failure: A White Paper From the Heart Failure Society of America Scientific Statements Committee. J. Card. Fail., 2018 Oct 12;24(10). PMID 30308242*
34. *Centers for Medicare & Medicaid Services (CMS). National coverage decision for cardiac output monitoring by thoracic electrical bioimpedance (TEB) (20.16). 2006*
35. *Blue Cross Blue Shield Association Medical Policy Reference Manual. 2.02.24, Cardiac Hemodynamic Monitoring for the Management of Heart Failure in the Outpatient Setting. July 2023*

Other: *Mosby's Medical, Nursing & Allied Health Dictionary, 6th edition. Taber's Cyclopedic Medical Dictionary, 21st edition.*

MEDICAL POLICY

POLICY TITLE	CARDIAC HEMODYNAMIC MONITORING FOR THE MANAGEMENT OF HEART FAILURE IN THE OUTPATIENT SETTING
POLICY NUMBER	MP 2.051

X. POLICY HISTORY

[Top](#)

MP 2.051	CAC 8/30/05
	CAC 6/27/06
	CAC 6/26/07
	CAC 5/27/08
	CAC 5/26/09
	CAC 5/25/10 Consensus
	CAC 7/26/11 Adopt BCBSA. CBC policy addressed thoracic bioimpedance and inert gas rebreathing only. By adopting BCBSA - arterial pressure Vasalva and implantable direct pressure monitoring of the pulmonary artery is also listed as investigational. CBC was silent on these prior to this revision. Other policy statements unchanged.
	CAC 10/30/12 Consensus Review. No change to policy statements. References updated. FEP variation revised to refer to the FEP medical policy manual. Codes reviewed 10/23/12
	CAC 11/26/13 Consensus. No change to policy statements. Changed Medicare variation to reference Novitas Solutions Local Coverage Determination (LCD) L32684 Cardiac Output Monitoring by Bioimpedance. Rationale added.
	CAC 11/25/14 Consensus review. No changes to the policy statements. References and rationale updated. Medicare variation revised to reflect NCD 20.16 Cardiac Output Monitoring by Thoracic Electrical Bioimpedance as LCD L32684 has been retired. Coding reviewed, no changes.
	03/26/2015-Administrative code review. No changes to policy stance.
	9/24/15 Administrative Change. Added 4 th Quarter New Code.
	CAC 11/24/15 Consensus review. No changes to the policy statements. References and rationale updated. Coding updated including 2016 updates.
	4/7/16 Administrative change. Added Medicare variation to reference L36419.
	5/10/2016 Administrative Change: Typographical error corrected; code is 93701, not 92701 (invalid code).
	CAC 11/29/16 Consensus review. No change to policy statements. References and rationale updated. Variation reformatting. Coding Reviewed
	Admin Update 1/1/18: Removed end dated codes 0293T and 0294T; effective 1/1/18. Medicare variations removed from Commercial Policies.
9/21/2017 Consensus review. Verbiage clarification to policy statement, intent unchanged. Description/Background, Rationale and Reference sections updated. Coding Reviewed.	
6/5/18 Consensus. No change to policy statements. References updated. Rationale condensed.	
1/1/19 Admin Update: Added new codes effective 1/1/19 and removed deleted HCPCS code.	

MEDICAL POLICY

POLICY TITLE	CARDIAC HEMODYNAMIC MONITORING FOR THE MANAGEMENT OF HEART FAILURE IN THE OUTPATIENT SETTING
POLICY NUMBER	MP 2.051

	6/10/19 Consensus review. No change to policy statements. References updated.
	01/01/20 Admin code update. New code added, C9728.
	5/20/2020 Admin code update. New codes 0607T and 0608T added.
	6/3/2020 Consensus Review. No change to policy statement. Rationale and references updated.
	11/1/2021 Consensus Review. Policy statement unchanged. FEP language updated. Background, Rationale and References updated.
	08/24/2022 Consensus review. No change in policy statement. Product variation and FEP language revised. Background, Rationale and References updated. Removed C9758 from policy.
	07/12/2023 Consensus Review. No change to policy statement. Background and Rationale updated. References added.
	1/18/2024 Administrative review. Clinical benefit added.

[Top](#)

Health care benefit programs issued or administered by Capital Blue Cross and/or its subsidiaries, Capital Advantage Insurance Company[®], Capital Advantage Assurance Company[®] and Keystone Health Plan[®] Central. Independent licensees of the Blue Cross BlueShield Association. Communications issued by Capital Blue Cross in its capacity as administrator of programs and provider relations for all companies.