

MEDICAL POLICY

POLICY TITLE	GENETIC TESTING FOR IDIOPATHIC DILATED CARDIOMYOPATHY
POLICY NUMBER	MP 2.328

CLINICAL BENEFIT	☐ MINIMIZE SAFETY RISK OR CONCERN. ☒ MINIMIZE HARMFUL OR INEFFECTIVE INTERVENTIONS. ☐ ASSURE APPROPRIATE LEVEL OF CARE. ☐ ASSURE APPROPRIATE DURATION OF SERVICE FOR INTERVENTIONS. ☐ ASSURE THAT RECOMMENDED MEDICAL PREREQUISITES HAVE BEEN MET. ☐ ASSURE APPROPRIATE SITE OF TREATMENT OR SERVICE.
Effective Date:	RETIRED 7/1/2026

[POLICY](#)
[RATIONALE](#)
[DISCLAIMER](#)
[POLICY HISTORY](#)

[PRODUCT VARIATIONS](#)
[DEFINITIONS](#)
[CODING INFORMATION](#)

[DESCRIPTION/BACKGROUND](#)
[BENEFIT VARIATIONS](#)
[REFERENCES](#)

I. POLICY

Comprehensive genetic testing for individuals with signs or symptoms of dilated cardiomyopathy (DCM) which is considered idiopathic after a negative workup for secondary causes is considered **medically necessary**.

Targeted genetic testing for asymptomatic individuals with a first-degree relative who has dilated cardiomyopathy, and a known familial variant is considered **medically necessary**.

Genetic testing for dilated cardiomyopathy is considered **investigational** in all other situations as there is insufficient evidence to support a general conclusion concerning the health outcomes or benefits associated with this procedure.

Policy Guidelines

Standard Workup for Patients with Signs or Symptoms of Dilated Cardiomyopathy

The standard workup for patients with signs or symptoms of DCM includes a clinical exam, blood pressure monitoring, electrocardiography, echocardiography, and workup for coronary artery disease as warranted by risk factors. An extensive workup including cardiac magnetic resonance imaging, exercise testing, right-sided catheterization with biopsy, and 24-hour electrocardiography monitoring will uncover only a small number of additional etiologies for DCM.

Genetics Nomenclature Update

The Human Genome Variation Society nomenclature is used to report information on variants found in DNA and serves as an international standard in DNA diagnostics. It is being implemented for genetic testing medical evidence review updates starting in 2017 (see Table PG1). The Society's nomenclature is recommended by the Human Variome Project, the Human Genome Organization, and by the Human Genome Variation Society itself.

MEDICAL POLICY

POLICY TITLE	GENETIC TESTING FOR IDIOPATHIC DILATED CARDIOMYOPATHY
POLICY NUMBER	MP 2.328

The American College of Medical Genetics and Genomics and the Association for Molecular Pathology standards and guidelines for interpretation of sequence variants represent expert opinion from both organizations, in addition to the College of American Pathologists. These recommendations primarily apply to genetic tests used in clinical laboratories, including genotyping, single genes, panels, exomes, and genomes. Table PG2 shows the recommended standard terminology - "pathogenic," "likely pathogenic," "uncertain significance," "likely benign," and "benign" - to describe variants identified that cause Mendelian disorders.

Table PG1. Nomenclature to Report on Variants Found in DNA

Previous	Updated	Definition
Mutation	Diseased-Associated Variant	Disease-associated change in the DNA sequence.
	Variant	Change in DNA sequence
	Familial Variant	Disease-associated variant identified in a proband for use in subsequent targeted genetic testing in first-degree relatives.

Table PG2. ACMG-AMP Standards and Guidelines for Variant Classification

Variant Classification	Definition
Pathogenic	Disease-causing change in the DNA sequence
Likely Pathogenic	Likely disease-causing change in the DNA sequence
Variant of uncertain significance	Change in DNA sequence with uncertain effects on disease
Likely benign	Likely benign change in the DNA sequence
Benign	Benign change in the DNA sequence

ACMG: American College of Medical Genetics and Genomics; AMP: Association of Molecular Pathology.

GENETIC COUNSELING

Genetic counseling is primarily aimed at patients who are at risk for inherited disorders, and experts recommend formal genetic counseling in most cases when genetic testing for an inherited condition is considered. The interpretation of the results of genetic tests and the understanding of risk factors can be very difficult and complex. Therefore, genetic counseling will assist individuals in understanding the possible benefits and harms of genetic testing, including the possible impact of the information on the individual's family. Genetic counseling may alter the utilization of genetic testing substantially and may reduce inappropriate testing. Genetic counseling should be performed by an individual with experience and expertise in genetic medicine and genetic testing methods.

Cross-References:

MP 2.233 Genetic Testing for Cardiac Ion Channelopathies

MP 2.248 Genetic Testing for Predisposition to Inherited Hypertrophic Cardiomyopathy

MP 2.323 General Approach to Evaluating the Utility of Genetic Panels

MEDICAL POLICY

POLICY TITLE	GENETIC TESTING FOR IDIOPATHIC DILATED CARDIOMYOPATHY
POLICY NUMBER	MP 2.328

II. PRODUCT VARIATIONS

[TOP](#)

This policy is only applicable to certain programs and products administered by Capital Blue Cross and subject to benefit variations as discussed in Section VI. Please see additional information below.

FEP PPO- Refer to FEP Medical Policy Manual. The FEP Medical Policy manual can be found at:

<https://www.fepblue.org/benefit-plans/medical-policies-and-utilization-management-guidelines/medical-policies>.

III. DESCRIPTION/BACKGROUND

[TOP](#)

Dilated cardiomyopathy is characterized by progressive left ventricular enlargement and systolic dysfunction, leading to clinical manifestations of heart failure. There are a variety of causes of DCM, including genetic and nongenetic conditions. Genetic forms of DCM are heterogeneous in their molecular basis and clinical expression. Genetic testing for DCM has potential utility for confirming a diagnosis of genetic DCM and as a prognostic test in family members when familial DCM is present.

Dilated Cardiomyopathy

Dilated cardiomyopathy is defined as the presence of left ventricular enlargement and dilatation in conjunction with significant systolic dysfunction. DCM has an estimated prevalence of 1 in 2700 in the United States. The age of onset for DCM varies, ranging from infancy to the eighth decade, with most individuals developing symptoms in the fourth through sixth decades.

Idiopathic Dilated Cardiomyopathy

When a patient presents with DCM, a workup is performed to identify underlying causes, especially those treatable. The standard workup consists of a clinical exam, blood pressure monitoring, electrocardiography, echocardiography, and workup for coronary artery disease as warranted by risk factors. An extensive workup including cardiac magnetic resonance imaging (MRI), exercise testing, right-sided catheterization with biopsy, and 24-hour electrocardiography monitoring will uncover only a small number of additional etiologies for DCM. Approximately 35% to 40% of DCM cases are thus determined to be idiopathic after a negative workup for secondary causes listed above. This has traditionally been termed idiopathic dilated cardiomyopathy (IDC).

Clustering of IDC within families has been reported, leading to the conclusion that at least some cases of DCM have a genetic basis. Familial DCM is diagnosed when two closely related family members have IDC in the absence of underlying causes. Penetrance of familial DCM is variable and age-dependent, often leading to a lack of appreciation of the familial component.

Genetic Dilated Cardiomyopathy

MEDICAL POLICY

POLICY TITLE	GENETIC TESTING FOR IDIOPATHIC DILATED CARDIOMYOPATHY
POLICY NUMBER	MP 2.328

Genetic DCM has been proposed as a newer classification that includes both familial DCM and some cases of sporadic IDC. The percentage of patients with sporadic DCM that has a genetic basis is not well characterized. Most disease-associated variants are inherited in an autosomal dominant fashion, but some autosomal recessive, X-linked, and mitochondrial patterns of inheritance also are present. Expanded numbers of genotyped individuals facilitate genotype-phenotype correlations and studies of natural disease history. Recognition of high-risk variant carriers is important as these individuals would be expected to have the most to gain from pre-emptive interventions.

In general, genotype-phenotype correlations in the inherited cardiomyopathies are either not present or not well characterized. There have been some purported correlations between certain disease-associated variants and the presence of arrhythmias. For example, patients with conduction system disease and/or a family history of sudden cardiac death may be more likely to have disease-associated variants in the lamin A/C (LM), *SCN5A*, and desmin genes. Kayvanpour et al (2017) performed a meta-analysis of genotype-phenotype associations in DCM. The analysis included 48 studies (N=8097 patients) and found a higher prevalence of sudden cardiac death, cardiac transplantation, and ventricular arrhythmias in the LM and phospholamban (PLN) disease-associated variant carriers and increasing penetrance with age of DCM phenotype in subjects with titin (TTN)-truncating variants.

There may be interactions between genetic and environmental factors that lead to the clinical manifestations of DCM. A genetic variant may not in itself be sufficient to cause DCM but may predispose to developing DCM in the presence of environmental factors such as nutritional deficiencies or viral infections. It also has been suggested that DCM genetics may be more complex than single-gene variants, with low-penetrance variants that are common in the population contributing to a cumulative risk of DCM that includes both genetic and environmental factors.

Diagnosis of Dilated Cardiomyopathy

Primary clinical manifestations of DCM are heart failure and arrhythmias. Symptoms of heart failure, such as dyspnea on exertion and peripheral edema, are the most common presentation of DCM. These symptoms are generally gradual in onset and slowly progressive over time. Progressive myocardial dysfunction also may lead to electrical instability and arrhythmias. Symptoms of arrhythmias may include light-headedness, syncope, or sudden cardiac arrest.

Many underlying conditions can cause DCM, including:

- Ischemic coronary artery disease
- Toxins
- Metabolic conditions
- Endocrine disorders
- Inflammatory and infectious diseases
- Infiltrative disorders
- Tachycardia-mediated cardiomyopathy.

Treatment of Dilated Cardiomyopathy

MEDICAL POLICY

POLICY TITLE	GENETIC TESTING FOR IDIOPATHIC DILATED CARDIOMYOPATHY
POLICY NUMBER	MP 2.328

Treatment of DCM is similar to that for other causes of heart failure. This includes medications to reduce fluid overload and relieve strain on the heart and lifestyle modifications such as salt restriction. Patients with clinically significant arrhythmias also may be treated with antiarrhythmic medications, pacemaker implantation, and/or an automatic implantable cardiac defibrillator. Automatic implantable cardiac defibrillator placement for primary prevention also may be performed if criteria for low ejection fraction and/or other clinical symptoms are present. End-stage DCM can be treated with cardiac transplantation.

Genetic Testing for Dilated Cardiomyopathy

Approximately 30% to 40% of patients with DCM referred for genetic testing will have a disease-associated variant identified. Disease-associated variants linked to DCM have been identified in more than 40 genes of various types and locations. The most common genes involved are those that code for TTN, myosin heavy chain (*MYH7*), troponin T (*TNNT2*), and alpha-tropomyosin (*TPM1*). These 4 genes account for approximately 30% of disease-associated variants identified in cohorts of patients with DCM. A high proportion of the identified disease-associated variants are rare, or novel, variants, thus creating challenges in assigning the pathogenicity of discovered variants. Some individuals with DCM will have more than one DCM-associated variant. The frequency of multiple disease-associated variants is uncertain, as is the clinical significance.

Regulatory Status

Clinical laboratories may develop and validate tests in-house and market them as a laboratory service; laboratory-developed tests must meet the general regulatory standards of the Clinical Laboratory Improvement Amendments (CLIA). Laboratories that offer laboratory-developed tests must be licensed by the CLIA for high-complexity testing. To date, the U.S. Food and Drug Administration has chosen not to require any regulatory review of this test.

IV. RATIONALE

[TOP](#)

SUMMARY OF EVIDENCE

For individuals who have signs and/or symptoms of DCM who receive comprehensive genetic testing, the evidence includes large case series reporting clinical validity and prospective observational studies reporting clinical utility. Relevant outcomes are overall survival, test validity, symptoms, change in disease status, functional outcomes, quality of life, and treatment-related morbidity. The percentage of patients with idiopathic DCM who have a genetic variant (clinical sensitivity) is relatively low, in the range of 10% to 40%. Additional studies assessed clinical outcomes of patients with DCM and at least one known variant compared with patients with DCM and no known variants. The studies reported that patients with DCM and known variants experienced lower event-free survival, earlier onset of symptoms, lower transplant-free survival, and more life-threatening arrhythmias compared with patients with DCM and no known variants. A prospective observational study has reported that patients with DCM and known variants experienced high rates of morbidity and mortality during four to eight years of follow-up. While direct evidence of clinical usefulness is lacking, confirming a diagnosis can lead to changes in clinical management, which improve net health outcomes. Changes in management may include earlier implantation of cardiac defibrillators or increased surveillance to detect

MEDICAL POLICY

POLICY TITLE	GENETIC TESTING FOR IDIOPATHIC DILATED CARDIOMYOPATHY
POLICY NUMBER	MP 2.328

worsening of symptoms, as well as cascade genetic testing of asymptomatic family members. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

For individuals who are asymptomatic with a first-degree relative who has DCM and a known familial variant who receive targeted genetic testing for a known familial variant, the evidence includes retrospective studies and case series reporting clinical value and a prospective observational study reporting clinical utility. Relevant outcomes are test validity, symptoms, morbid events, functional outcomes, quality of life, and treatment-related morbidity. For an individual at-risk due to genetic DCM in the family, genetic testing can identify whether a familial variant has been inherited. A prospective observational study with 4 to 8 years of follow-up reported the development of cardiac symptoms among patients initially asymptomatic who had DCM-related variants. While direct evidence of clinical usefulness is lacking, confirming a diagnosis can lead to changes in clinical management, which improve net health outcomes. Changes in management may include periodic clinical and cardiovascular evaluations to detect the earliest signs of disease, as well as genetic counseling. The evidence is sufficient to determine that the technology results in an improvement in the net health outcome.

V. DEFINITIONS

[TOP](#)

NA

VI. BENEFIT VARIATIONS

[TOP](#)

The existence of this medical policy does not mean that this service is a covered benefit under the member's health benefit plan. Benefit determinations are based on the applicable health benefit plan language. Medical policies do not constitute a description of benefits. Members and providers should consult the member's health benefit plan for information or contact Capital Blue Cross for benefit information.

VII. DISCLAIMER

[TOP](#)

Capital Blue Cross' medical policies are developed to assist in administering a member's benefits. These medical policies do not constitute medical advice and are subject to change. Treating providers are solely responsible for medical advice and treatment of members. Members should discuss any medical policy related to their coverage or condition with their provider and consult their benefit information to determine if the service is covered. If there is a discrepancy between this medical policy and a member's benefit information, the benefit information will govern. If a provider or a member has a question concerning the application of this medical policy to a specific member's plan of benefits, please contact Capital Blue Cross' Provider Services or Member Services. Capital Blue Cross considers the information contained in this medical policy to be proprietary and it may only be disseminated as permitted by law.

MEDICAL POLICY

POLICY TITLE	GENETIC TESTING FOR IDIOPATHIC DILATED CARDIOMYOPATHY
POLICY NUMBER	MP 2.328

VIII. CODING INFORMATION

[TOP](#)

Note: This list of codes may not be all-inclusive, and codes are subject to change at any time. The identification of a code in this section does not denote coverage as coverage is determined by the terms of member benefit information. In addition, not all covered services are eligible for separate reimbursement.

Medically Necessary when used to report Genetic Testing for Cardiomyopathy:

Procedure Codes							
81403	81405	81406	81407	81439	81479		
ICD-10-CM Diagnosis Codes	Description						
I42.0	Dilated cardiomyopathy						

IX. REFERENCES

[TOP](#)

- Hersheberger RE, Morales A. Dilated Cardiomyopathy Overview. In: Adam MP, Ardinger HH, Pagon RA, et al., eds. GeneReviews. Seattle, WA: University of Washington; 2015.
- Piran S, Liu P, Morales A, et al. Where genome meets phenome: rationale for integrating genetic and protein biomarkers in the diagnosis and management of dilated cardiomyopathy and heart failure. *J Am Coll Cardiol*. Jul 24 2012; 60(4): 283-9. PMID 22813604
- Broch K, Andreassen AK, Hopp E, et al. Results of comprehensive diagnostic work-up in 'idiopathic' dilated cardiomyopathy. *Open Heart*. 2015; 2(1): e000271. PMID 26468400
- Hersheberger RE, Morales A, Siegfried JD. Clinical and genetic issues in dilated cardiomyopathy: a review for genetics professionals. *Genet Med*. Nov 2010; 12(11): 655-67. PMID 20864896
- Lakdawala NK, Winterfield JR, Funke BH. Dilated cardiomyopathy. *Circ Arrhythm Electrophysiol*. Feb 2013; 6(1): 228-37. PMID 23022708
- Fatkin D, Huttner IG, Kovacic JC, et al. Precision Medicine in the Management of Dilated Cardiomyopathy: JACC State-of-the-Art Review. *J Am Coll Cardiol*. Dec 10 2019; 74(23): 2921-2938. PMID 31806137
- Kayvanpour E, Sedaghat-Hamedani F, Amr A, et al. Genotype-phenotype associations in dilated cardiomyopathy: meta-analysis on more than 8000 individuals. *Clin Res Cardiol*. Feb 2017; 106(2): 127-139. PMID 27576561
- National Center for Biotechnology Information. Genetic Testing Registry.
- Kadish A, Dyer A, Daubert JP, et al. Prophylactic defibrillator implantation in patients with nonischemic dilated cardiomyopathy. *N Engl J Med*. May 20 2004; 350(21): 2151-8. PMID 15152060
- Bardy GH, Lee KL, Mark DB, et al. Amiodarone or an implantable cardioverter-defibrillator for congestive heart failure. *N Engl J Med*. Jan 20 2005; 352(3): 225-37. PMID 15659722
- Brodsky GL, Muntoni F, Micić S, et al. Lamin A/C gene mutation associated with dilated cardiomyopathy with variable skeletal muscle involvement. *Circulation*. Feb 08 2000; 101(5): 473-6. PMID 10662742

MEDICAL POLICY

POLICY TITLE	GENETIC TESTING FOR IDIOPATHIC DILATED CARDIOMYOPATHY
POLICY NUMBER	MP 2.328

12. MacLeod HM, Culley MR, Huber JM, et al. Lamin A/C truncation in dilated cardiomyopathy with conduction disease. *BMC Med Genet.* Jul 10 2003; 4: 4. PMID 12854972
13. Olson TM, Michels VV, Thibodeau SN, et al. Actin mutations in dilated cardiomyopathy, a heritable form of heart failure. *Science.* May 01 1998; 280(5364): 750-2. PMID 9563954
14. Sylvius N, Duboscq-Bidot L, Bouchier C, et al. Mutational analysis of the beta- and delta-sarcoglycan genes in a large number of patients with familial and sporadic dilated cardiomyopathy. *Am J Med Genet A.* Jul 01 2003; 120A(1): 8-12. PMID 12794684
15. Taylor MR, Slavov D, Ku L, et al. Prevalence of desmin mutations in dilated cardiomyopathy. *Circulation.* Mar 13 2007; 115(10): 1244-51. PMID 17325244
16. Villard E, Duboscq-Bidot L, Charron P, et al. Mutation screening in dilated cardiomyopathy: prominent role of the beta myosin heavy chain gene. *Eur Heart J.* Apr 2005; 26(8): 794-803. PMID 15769782
17. Dhandapany PS, Razzaque MA, Muthusami U, et al. RAF1 mutations in childhood-onset dilated cardiomyopathy. *Nat Genet.* Jun 2014; 46(6): 635-639. PMID 24777450
18. McNair WP, Sinagra G, Taylor MR, et al. SCN5A mutations associate with arrhythmic dilated cardiomyopathy and commonly localize to the voltage-sensing mechanism. *J Am Coll Cardiol.* May 24 2011; 57(21): 2160-8. PMID 21596231
19. van Rijnsingen IA, Nannenberg EA, Arbustini E, et al. Gender-specific differences in major cardiac events and mortality in lamin A/C mutation carriers. *Eur J Heart Fail.* Apr 2013; 15(4): 376-84. PMID 23183350
20. Herman DS, Lam L, Taylor MR, et al. Truncations of titin causing dilated cardiomyopathy. *N Engl J Med.* Feb 16 2012; 366(7): 619-28. PMID 22335739
21. Theis JL, Sharpe KM, Matsumoto ME, et al. Homozygosity mapping and exome sequencing reveal GATAD1 mutation in autosomal recessive dilated cardiomyopathy. *Circ Cardiovasc Genet.* Dec 2011; 4(6): 585-94. PMID 21965549
22. Norton N, Li D, Rieder MJ, et al. Genome-wide studies of copy number variation and exome sequencing identify rare variants in BAG3 as a cause of dilated cardiomyopathy. *Am J Hum Genet.* Mar 11 2011; 88(3): 273-82. PMID 21353195
23. van der Meulen MH, Herkert JC, den Boer SL, et al. Genetic Evaluation of A Nation-Wide Dutch Pediatric DCM Cohort: The Use of Genetic Testing in Risk Stratification. *Circ Genom Precis Med.* Oct 2022; 15(5): e002981. PMID 36178741
24. Hofmeyer M, Haas GJ, Jordan E, et al. Rare Variant Genetics and Dilated Cardiomyopathy Severity: The DCM Precision Medicine Study. *Circulation.* Sep 12 2023; 148(11): 872-881. PMID 37641966
25. Dalin MG, Engström PG, Ivarsson EG, et al. Massive parallel sequencing questions the pathogenic role of missense variants in dilated cardiomyopathy. *Int J Cardiol.* Feb 01 2017; 228: 742-748. PMID 27886618
26. Haas J, Frese KS, Peil B, et al. Atlas of the clinical genetics of human dilated cardiomyopathy. *Eur Heart J.* May 07 2015; 36(18): 1123-35a. PMID 25163546
27. Pugh TJ, Kelly MA, Gowrisankar S, et al. The landscape of genetic variation in dilated cardiomyopathy as surveyed by clinical DNA sequencing. *Genet Med.* Aug 2014; 16(8): 601-8. PMID 24503780
28. University of Bologna. ws-SNPs&GO. n.d.
29. National Center for Biotechnology Information. Genetic Testing Registry

MEDICAL POLICY

POLICY TITLE	GENETIC TESTING FOR IDIOPATHIC DILATED CARDIOMYOPATHY
POLICY NUMBER	MP 2.328

30. Hirtle-Lewis M, Desbiens K, Ruel I, et al. The genetics of dilated cardiomyopathy: a prioritized candidate gene study of LMNA, TNNT2, TCAP, and PLN. *Clin Cardiol.* Oct 2013; 36(10): 628-33. PMID 24037902
31. van der Linde IHM, Hiemstra YL, Bökenkamp R, et al. A Dutch MYH7 founder mutation, p.(Asn1918Lys), is associated with early onset cardiomyopathy and congenital heart defects. *Neth Heart J.* Dec 2017; 25(12): 675-681. PMID 28864942
32. Myers VD, Gerhard GS, McNamara DM, et al. Association of Variants in BAG3 With Cardiomyopathy Outcomes in African American Individuals. *JAMA Cardiol.* Oct 01 2018; 3(10): 929-938. PMID 30140897
33. Verdonschot JAJ, Hazebroek MR, Derks KWJ, et al. Titin cardiomyopathy leads to altered mitochondrial energetics, increased fibrosis and long-term life-threatening arrhythmias. *Eur Heart J.* Mar 07 2018; 39(10): 864-873. PMID 29377983
34. Ebert M, Wijnmaalen AP, de Riva M, et al. Prevalence and Prognostic Impact of Pathogenic Variants in Patients With Dilated Cardiomyopathy Referred for Ventricular Tachycardia Ablation. *JACC Clin Electrophysiol.* Sep 2020; 6(9): 1103-1114. PMID 32972544
35. Millat G, Bouvagnet P, Chevalier P, et al. Clinical and mutational spectrum in a cohort of 105 unrelated patients with dilated cardiomyopathy. *Eur J Med Genet.* 2011; 54(6): e570-5. PMID 21846512
36. Lakdawala NK, Funke BH, Baxter S, et al. Genetic testing for dilated cardiomyopathy in clinical practice. *J Card Fail.* Apr 2012; 18(4): 296-303. PMID 22464770
37. Priganc M, Zigová M, Boroňová I, et al. Analysis of SCN5A Gene Variants in East Slovak Patients with Cardiomyopathy. *J Clin Lab Anal.* Mar 2017; 31(2). PMID 27554632
38. van Rijsingen IA, van der Zwaag PA, Groeneweg JA, et al. Outcome in phospholamban R14del carriers: results of a large multicentre cohort study. *Circ Cardiovasc Genet.* Aug 2014; 7(4): 455-65. PMID 24909667
39. Hasselberg NE, Edvardsen T, Petri H, et al. Risk prediction of ventricular arrhythmias and myocardial function in Lamin A/C mutation positive subjects. *Europace.* Apr 2014; 16(4): 563-71. PMID 24058181
40. Hasselberg NE, Haland TF, Saberniak J, et al. Lamin A/C cardiomyopathy: young onset, high penetrance, and frequent need for heart transplantation. *Eur Heart J.* Mar 07 2018; 39(10): 853-860. PMID 29095976
41. Reddy S, Fung A, Manhiot C, et al. Adrenergic receptor genotype influences heart failure severity and β -blocker response in children with dilated cardiomyopathy. *Pediatr Res.* Feb 2015; 77(2): 363-9. PMID 25406899
42. Wasielewski M, van Spaendonck-Zwarts KY, Westerink ND, et al. Potential genetic predisposition for anthracycline-associated cardiomyopathy in families with dilated cardiomyopathy. *Open Heart.* 2014; 1(1): e000116. PMID 25332820
43. Michels VV, Moll PP, Miller FA, et al. The frequency of familial dilated cardiomyopathy in a series of patients with idiopathic dilated cardiomyopathy. *N Engl J Med.* Jan 09 1992; 326(2): 77-82. PMID 1727235
44. Grünig E, Tasman JA, Kücherer H, et al. Frequency and phenotypes of familial dilated cardiomyopathy. *J Am Coll Cardiol.* Jan 1998; 31(1): 186-94. PMID 9426039
45. Baig MK, Goldman JH, Caforio AL, et al. Familial dilated cardiomyopathy: cardiac abnormalities are common in asymptomatic relatives and may represent early disease. *J Am Coll Cardiol.* Jan 1998; 31(1): 195-201. PMID 9426040

MEDICAL POLICY

POLICY TITLE	GENETIC TESTING FOR IDIOPATHIC DILATED CARDIOMYOPATHY
POLICY NUMBER	MP 2.328

46. Mahon NG, Murphy RT, MacRae CA, et al. Echocardiographic evaluation in asymptomatic relatives of patients with dilated cardiomyopathy reveals preclinical disease. *Ann Intern Med.* Jul 19 2005; 143(2): 108-15. PMID 16027452
47. Brodt C, Siegfried JD, Hofmeyer M, et al. Temporal relationship of conduction system disease and ventricular dysfunction in LMNA cardiomyopathy. *J Card Fail.* Apr 2013; 19(4): 233-9. PMID 23582089
48. Huggins GS, Kinnamon DD, Haas GJ, et al. Prevalence and Cumulative Risk of Familial Idiopathic Dilated Cardiomyopathy. *JAMA.* Feb 01 2022; 327(5): 454-463. PMID 35103767
49. Vissing CR, Espersen K, Mills HL, et al. Family Screening in Dilated Cardiomyopathy: Prevalence, Incidence, and Potential for Limiting Follow-Up. *JACC Heart Fail.* Nov 2022; 10(11): 792-803. PMID 36328645
50. Stava TT, Leren TP, Bogsrud MP. Molecular genetics in 4408 cardiomyopathy probands and 3008 relatives in Norway: 17 years of genetic testing in a national laboratory. *Eur J Prev Cardiol.* Oct 18 2022; 29(13): 1789-1799. PMID 35653365
51. Fernlund E, Österberg AW, Kuchinskaya E, et al. Novel Genetic Variants in BAG3 and TNNT2 in a Swedish Family with a History of Dilated Cardiomyopathy and Sudden Cardiac Death. *Pediatr Cardiol.* Aug 2017; 38(6): 1262-1268. PMID 28669108
52. Asadi M, Foo R, Salehi AR, et al. Mutation in δ -Sg Gene in Familial Dilated Cardiomyopathy. *Adv Biomed Res.* 2017; 6: 32. PMID 28401079
53. Bodian DL, Vilboux T, Hourigan SK, et al. Genomic analysis of an infant with intractable diarrhea and dilated cardiomyopathy. *Cold Spring Harb Mol Case Stud.* Nov 2017; 3(6). PMID 28701297
54. Yuan HX, Yan K, Hou DY, et al. Whole exome sequencing identifies a KCNJ12 mutation as a cause of familial dilated cardiomyopathy. *Medicine (Baltimore).* Aug 2017; 96(33): e7727. PMID 28816949
55. Petropoulou E, Soltani M, Firoozabadi AD, et al. Digenic inheritance of mutations in the cardiac troponin (TNNT2) and cardiac beta myosin heavy chain (MYH7) as the cause of severe dilated cardiomyopathy. *Eur J Med Genet.* Sep 2017; 60(9): 485-488. PMID 28642161
56. Rafiq MA, Chaudhry A, Care M, et al. Whole exome sequencing identified 1 base pair novel deletion in BCL2-associated athanogene 3 (BAG3) gene associated with severe dilated cardiomyopathy (DCM) requiring heart transplant in multiple family members. *Am J Med Genet A.* Mar 2017; 173(3): 699-705. PMID 28211974
57. Liu JS, Fan LL, Zhang H, et al. Whole-Exome Sequencing Identifies Two Novel TTN Mutations in Chinese Families with Dilated Cardiomyopathy. *Cardiology.* 2017; 136(1): 10-14. PMID 27544385
58. Posafalvi A, Herkert JC, Sinke RJ, et al. Clinical utility gene card for: dilated cardiomyopathy (CMD). *Eur J Hum Genet.* Oct 2013; 21(10). PMID 23249954
59. Bozkurt B, Colvin M, Cook J, et al. Current Diagnostic and Treatment Strategies for Specific Dilated Cardiomyopathies: A Scientific Statement From the American Heart Association. *Circulation.* Dec 06 2016; 134(23): e579-e646. PMID 27832612
60. Hersberger RE, Givertz MM, Ho CY, et al. Genetic evaluation of cardiomyopathy: a clinical practice resource of the American College of Medical Genetics and Genomics (ACMG). *Genet Med.* Sep 2018; 20(9): 899-909. PMID 29904160

MEDICAL POLICY

POLICY TITLE	GENETIC TESTING FOR IDIOPATHIC DILATED CARDIOMYOPATHY
POLICY NUMBER	MP 2.328

61. Ackerman MJ, Priori SG, Willems S, et al. HRS/EHRA expert consensus statement on the state of genetic testing for the channelopathies and cardiomyopathies this document was developed as a partnership between the Heart Rhythm Society (HRS) and the European Heart Rhythm Association (EHRA). *Heart Rhythm*. Aug 2011; 8(8): 1308-39. PMID 21787999
62. Hershberger RE, Givertz MM, Ho CY, et al. Genetic Evaluation of Cardiomyopathy-A Heart Failure Society of America Practice Guideline. *J Card Fail*. May 2018; 24(5): 281-302. PMID 29567486
63. Blue Cross Blue Shield Association Medical Policy Reference Manual Genetic Testing for Dilated Cardiomyopathy. March 2024

X. POLICY HISTORY

[TOP](#)

MP 2.328	03/13/2020 Consensus Review. No change to policy statement. References updated; coding reviewed.
	04/15/2021 Consensus Review. No change to policy statements. References updated; coding reviewed.
	03/16/2022 Consensus Review. Policy statement unchanged. FEP language updated. References added.
	03/07/2023 Consensus Review. No change to policy statement. Background, Rationale and References updated.
	03/12/2024 Consensus Review. No change to policy statement. References added.
	02/21/2025 Consensus Review. No change to policy statement. References updated.
	03/09/2026 Retirement Review. EviCore delegation.

[TOP](#)

Health care benefit programs issued or administered by Capital Blue Cross and/or its subsidiaries, Capital Advantage Insurance Company®, Capital Advantage Assurance Company®, and Keystone Health Plan® Central. Independent licensees of the Blue Cross BlueShield Association. Communications issued by Capital Blue Cross in its capacity as administrator of programs and provider relations for all companies.