

POLICY TITLE	LOW-LEVEL LASER THERAPY
POLICY NUMBER	MP- 1.097

Original Issue Date (Created):	7/26/2004
Most Recent Review Date (Revised):	5/8/2020
Effective Date:	8/1/2020

- | | | |
|----------------------------------|--|---|
| POLICY RATIONALE | PRODUCT VARIATIONS DEFINITIONS | DESCRIPTION/BACKGROUND BENEFIT VARIATIONS |
| DISCLAIMER | CODING INFORMATION | REFERENCES |
| POLICY HISTORY | | |

I. POLICY

[TOP](#)

Low-level laser therapy may be considered **medically necessary** for prevention of oral mucositis in patients undergoing cancer treatment associated with increased risk of oral mucositis, including chemotherapy and/or radiotherapy, and/or hematopoietic stem cell transplantation (see Policy Guidelines).

Low-level laser therapy is considered **investigational** for all other indications including but not limited to:

- Carpal tunnel syndrome
- Neck pain
- Subacromial impingement
- Adhesive capsulitis
- Temporomandibular joint pain
- Low back pain
- Osteoarthritis knee pain
- Heel pain (i.e., Achilles tendinopathy, plantar fasciitis)
- Rheumatoid arthritis
- Bell palsy
- Fibromyalgia
- Wound healing
- Lymphedema.

There is insufficient evidence to support a conclusion concerning the health outcomes or benefits associated with this procedure for these indications.

POLICY TITLE	LOW-LEVEL LASER THERAPY
POLICY NUMBER	MP- 1.097

Policy Guidelines

In the meta-analysis of 18 trials comparing low-level laser therapy (LLLT) to chemotherapy or chemoradiation for prevention of oral mucositis (Oberoi et al 2014), the course of LLLT was generally from day 0 through treatment. In studies of hematopoietic cell transplant (HCT), the course of LLLT began between day -7 and day 0 and continued as long as day 14 to 15. In studies that began LLLT at day -7 to day -5 before HCT, the course of laser therapy ended at day -1 to day 0.

Other protocols have used low-level laser energy applied to acupuncture points on the fingers and hand. This technique may be referred to as "laser acupuncture." Laser acupuncture is not reviewed in this herein.

Cross-reference:

MP-1.094 Skin Contact Monochromatic Infrared Energy for the Treatment of Cutaneous Ulcers, Diabetic Neuropathy, and Other Miscellaneous Musculoskeletal Conditions

II. PRODUCT VARIATIONS

[TOP](#)

This policy is only applicable to certain programs and products administered by Capital BlueCross and subject to benefit variations as discussed in Section VI. Please see additional information below.

FEP PPO: Refer to FEP Medical Policy Manual MP-2.01.56, Low-Level Laser Therapy. The FEP Medical Policy Manual can be found at: <https://www.fepblue.org/benefit-plans/medical-policies-and-utilization-management-guidelines/medical-policies>

III. DESCRIPTION/BACKGROUND

[TOP](#)

ORAL MUCOSITIS

Oral mucositis describes inflammation of the oral mucosa and typically manifests as erythema or ulcerations that appear seven to ten days after initiation of high-dose cancer therapy. Oral mucositis can cause significant pain and increased risk of systemic infection, dependency on total parenteral nutrition, and use of narcotic analgesics.

TREATMENT

Treatment planning may also need to be modified due to dose-limiting toxicity. There are a number of interventions for oral mucositis that may partially control symptoms, but none is considered a criterion standard treatment. When uncomplicated by infection, oral mucositis is

POLICY TITLE	LOW-LEVEL LASER THERAPY
POLICY NUMBER	MP- 1.097

self-limited and usually heals within two to four weeks after cessation of cytotoxic chemotherapy. Low-level laser therapy (LLLT) has been used in cancer therapy–induced oral mucositis in patients treated with radiotherapy and/or chemotherapy and hematopoietic cell transplantation.

MUSCULOSKELETAL AND NEUROLOGIC DISORDERS

Carpal tunnel syndrome (CTS) is the most common entrapment neuropathy and the most commonly performed surgery of the hand. The syndrome is related to the bony anatomy of the wrist. The carpal tunnel is bound dorsally and laterally by the carpal bones and ventrally by the transverse carpal ligament. Through this contained space run the nine flexor tendons and the median nerve. Therefore, any space occupying lesion can compress the median nerve and produce the typical symptoms of CTS - pain, numbness, and tingling in the distribution of the median nerve. Symptoms of more severe cases include hypesthesia, clumsiness, loss of dexterity, and weakness of pinch. In the most severe cases, patients experience marked sensory loss and significant functional impairment with thenar atrophy.

TREATMENT

Mild-to-moderate cases of CTS are usually first treated conservatively with splinting and cessation of aggravating activities. Other conservative therapies include oral steroids, diuretics, nonsteroidal anti-inflammatory drugs, and steroid injections into the carpal tunnel itself. Patients who do not respond to conservative therapy or who present with severe CTS with thenar atrophy may be considered candidates for surgical release of the carpal ligament, using either an open or endoscopic approach. LLLT is also used to treat CTS.

LOW-LEVEL LASER THERAPY

LLLT is the use of red-beam or near-infrared lasers with a wavelength between 600 and 1000 nm and power between 5 and 500 MW. (By comparison, lasers used in surgery typically use 300 W.) When applied to the skin, LLLT produces no sensation and does not burn the skin. Because of the low absorption by human skin, it is hypothesized that the laser light can penetrate deeply into the tissues where it has a photobiostimulative effect. The exact mechanism of its effect on tissue healing is unknown; hypotheses have included improved cellular repair and stimulation of the immune, lymphatic, and vascular systems.

LLLT is being evaluated to treat a wide variety of conditions, including soft tissue injuries, myofascial pain, tendinopathies, nerve injuries, joint pain, and lymphedema.

REGULATORY STATUS

A number of low-level lasers have been cleared for marketing by the U.S. Food and Drug Administration (FDA) through the 510(k) process for the treatment of pain. Data submitted for the MicroLight 830® Laser consisted of application of the laser over the carpal tunnel 3 times a week for 5 weeks. The labeling states that the “MicroLight 830 Laser is indicated for adjunctive

POLICY TITLE	LOW-LEVEL LASER THERAPY
POLICY NUMBER	MP- 1.097

use in the temporary relief of hand and wrist pain associated with Carpal Tunnel Syndrome.” In 2006, GRT LITE™ was cleared for marketing, listing the TUCO Erchonia PL3000, the Excalibur System, the MicroLight 830® Laser, and the Acculaser Pro as predicate devices. Indications of the GRT LITE™ for CTS are similar to the predicate devices: “adjunctive use in providing temporary relief of minor chronic pain.” In 2009, the LightStream™ LLL device was cleared for marketing by the FDA through the 510(k) process for adjunctive use in the temporary relief of pain associated with knee disorders treated in standard chiropractic practice. A number of clinical trials of LLLT are underway in the United States, including studies of wound healing. Since 2009, many more similar LLLT devices have received 510(k) clearance from the FDA; most recently, in 2018, Super Pulsed Laser technology (Multi Radiance Medical) was approved by the FDA through the premarket approval process for use in neck and shoulder pain.

IV. RATIONALE

[Top](#)

SUMMARY OF EVIDENCE

Oral Mucositis

For individuals who have increased risk of oral mucositis due to some cancer treatments (e.g., chemotherapy, radiotherapy) and/or hematopoietic cell transplantation who receive LLLT, the evidence includes randomized controlled trials (RCTs) and systematic reviews. Relevant outcomes are symptoms, morbid events, quality of life, and treatment-related morbidity. A 2014 systematic review included 18 RCTs and found better outcomes with LLLT used to prevent oral mucositis than with control treatments. RCTs published after the systematic review had similar findings. The evidence is sufficient to determine that the technology results in a meaningful improvement in the net health outcome.

Musculoskeletal and Neurologic Disorders

For individuals who have carpal tunnel syndrome who receive LLLT, the evidence includes RCTs and systematic reviews. Relevant outcomes are symptoms, functional outcomes, quality of life, and treatment related morbidity. Both a 2016 systematic review and a TEC Assessment (2010) did not find sufficient evidence from RCTs that LLLT improves outcomes. The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals who have neck pain who receive LLLT, the evidence includes RCTs and systematic reviews. The relevant outcomes are symptoms, functional outcomes, quality of life, and treatment-related morbidity. A 2013 systematic review identified 17 trials, most of which were considered low quality. Only two trials were considered moderate quality, and they found that LLLT led to better outcomes than placebo for chronic neck pain. A TEC Assessment (2010) found conflicting evidence. Additionally, laser types, application dosages, and treatment schedules vary in the available evidence and require further study. The evidence is insufficient to determine the effects of the technology on health outcomes.

POLICY TITLE	LOW-LEVEL LASER THERAPY
POLICY NUMBER	MP- 1.097

For individuals who have subacromial impingement syndrome who receive LLLT, the evidence includes RCTs. The relevant outcomes are symptoms, functional outcomes, quality of life, and treatment-related morbidity. Most trials did not show a significant benefit of LLLT compared with sham treatment or with an alternative intervention (e.g., exercise). The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals who have adhesive capsulitis who receive LLLT, the evidence includes RCTs and a systematic review. Relevant outcomes are symptoms, functional outcomes, quality of life, and treatment related morbidity. A Cochrane review evaluating treatments for adhesive capsulitis identified two RCTs assessing LLLT. Due to the small number of trials and study limitations, reviewers concluded that the evidence was insufficient to permit conclusions about the effectiveness of LLLT for adhesive capsulitis. The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals who have temporomandibular joint pain who receive LLLT, the evidence includes RCTs and several systematic reviews. Relevant outcomes are symptoms, functional outcomes, quality of life, and treatment-related morbidity. Meta-analyses of RCTs had mixed findings. A 2015 meta-analysis, which included 14 placebo-controlled randomized trials, did not find a statistically significant impact of LLLT on pain but did find that LLLT significantly improved functional outcomes (e.g., mouth opening). RCTs have not compared the impact of LLLT with physical therapy. The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals who have low back pain who receive LLLT, the evidence includes RCTs and systematic reviews. Relevant outcomes are symptoms, functional outcomes, quality of life, and treatment-related morbidity. Meta-analyses of RCTs found that LLLT resulted in a significantly greater reduction in pain scores and global assessment scores than a placebo control in the immediate posttreatment setting. Meta-analyses also found that other outcomes (e.g., disability index, range of motion) were significantly better immediately after treatment with active rather than placebo LLLT, but not at longer-term follow-up. The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals who have osteoarthritic knee pain who receive LLLT, the evidence includes RCTs and systematic reviews. Relevant outcomes are symptoms, functional outcomes, quality of life, and treatment related morbidity. A 2015 systematic review, which pooled study findings, did not find that LLLT significantly reduced pain or improved function outcomes compared with a sham intervention. The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals who have heel pain (i.e., Achilles tendinopathy, plantar fasciitis) who receive LLLT, the evidence includes RCTs. Relevant outcomes are symptoms, functional outcomes,

POLICY TITLE	LOW-LEVEL LASER THERAPY
POLICY NUMBER	MP- 1.097

quality of life, and treatment-related morbidity. Findings of two, sham-controlled randomized trials were inconsistent, and while an RCT compared LLLT with standard care lacked long-term follow-up. The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals who have rheumatoid arthritis who receive LLLT, the evidence includes RCTs and a systematic review. Relevant outcomes are symptoms, functional outcomes, quality of life, and treatment related morbidity. A systematic review of RCTs found inconsistent benefit of LLLT for a range of outcomes. A 2010 RCT, published after the systematic review, did not find that LLLT was significantly better than a placebo treatment on most outcomes. The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals who have Bell palsy who receive LLLT, the evidence includes two RCTs. Relevant outcomes are symptoms, functional outcomes, quality of life, and treatment-related morbidity. The RCT found significant short-term benefit of LLLT over exercise. Longer term outcomes (>6 weeks) were not available. Because Bell palsy often improves within weeks and may completely resolve within months, it is difficult to isolate specific improvements from laser therapy over the natural resolution of the illness. Also, no sham-controlled trials are available. The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals who have fibromyalgia who receive LLLT, the evidence includes RCTs. Relevant outcomes are symptoms, functional outcomes, quality of life, and treatment-related morbidity. The RCTs evaluating LLLT for treatment of fibromyalgia are small (i.e., <25 patients each). One RCT (N=20 patients) found significantly better outcomes with LLLT than with sham, while another (N=20 patients) did not find statistically significant between-group differences for similar outcomes. Additional RCTs with sufficient numbers of patients are needed to establish the efficacy of LLLT for fibromyalgia. The evidence is insufficient to determine the effects of the technology on health outcomes.

Wound Care and Lymphedema

For individuals who have chronic nonhealing wounds who receive LLLT, the evidence includes RCTs and systematic reviews. Relevant outcomes are symptoms, functional outcomes, quality of life, and treatment related morbidity. The few existing RCTs tend to have small sample sizes and potential risk of bias. The evidence is insufficient to determine the effects of the technology on health outcomes.

For individuals who have lymphedema who receive LLLT, the evidence includes RCTs and systematic reviews. Relevant outcomes are symptoms, functional outcomes, quality of life, and treatment-related morbidity. Two systematic reviews detected methodologic flaws in the available studies and did not consistently find better outcomes for patients receiving LLLT than

POLICY TITLE	LOW-LEVEL LASER THERAPY
POLICY NUMBER	MP- 1.097

those receiving a control condition for treatment of lymphedema. The evidence is insufficient to determine the effects of the technology on health outcomes.

V. DEFINITIONS

[TOP](#)

CARPAL TUNNEL SYNDROME is a condition of pain or numbness that affects some part of the median nerve distribution of the hand (the palmar side of the thumb, the index finger, the radial half of the ring finger, and the radial half of the palm) and may radiate into the arm.

EPICONDYLITIS is the inflammation of the epicondyle of the humerus and surrounding tissues.

FIBROMYALGIA is chronic and frequently difficult to manage pain in muscles and soft tissues surrounding joints.

RHEUMATOID ARTHRITIS is a chronic systemic disease marked by inflammation of multiple synovial joints.

TENDINITIS is an inflammation of a tendon.

TMJ SYNDROME is severe pain in and about the temporomandibular joint, made worse by chewing.

VI. BENEFIT VARIATIONS

[TOP](#)

The existence of this medical policy does not mean that this service is a covered benefit under the member's health benefit plan. Benefit determinations should be based in all cases on the applicable health benefit plan language. Medical policies do not constitute a description of benefits. A member's health benefit plan governs which services are covered, which are excluded, which are subject to benefit limits and which require preauthorization. There are different benefit plan designs in each product administered by Capital BlueCross. Members and providers should consult the member's health benefit plan for information or contact Capital BlueCross for benefit information.

VII. DISCLAIMER

[TOP](#)

Capital BlueCross's medical policies are developed to assist in administering a member's benefits, do not constitute medical advice and are subject to change. Treating providers are solely responsible for medical advice and treatment of members. Members should discuss any medical policy related to their coverage or condition with their provider and consult their benefit information to determine if the service is covered. If there is a discrepancy between this medical policy and a member's benefit information, the benefit information will govern. If a provider or a member has a question concerning the application of this medical policy to a specific member's plan of benefits, please contact Capital BlueCross' Provider Services or Member Services.

MEDICAL POLICY

POLICY TITLE	LOW-LEVEL LASER THERAPY
POLICY NUMBER	MP- 1.097

Capital BlueCross considers the information contained in this medical policy to be proprietary and it may only be disseminated as permitted by law.

VIII. CODING INFORMATION

[TOP](#)

Note: This list of codes may not be all-inclusive, and codes are subject to change at any time. The identification of a code in this section does not denote coverage as coverage is determined by the terms of member benefit information. In addition, not all covered services are eligible for separate reimbursement.

Investigational; therefore, not covered for low level laser therapy used for pain:

CPT Codes ®							
0552T							

Covered when medically necessary for low-level laser therapy for prevention of oral mucositis in patients undergoing cancer treatment:

CPT Codes ®							
97026							

Current Procedural Terminology (CPT) copyrighted by American Medical Association. All Rights Reserved.

HCPCS Code	Description
S8948	Application of a modality (requiring constant provider attendance) to one or more areas; low-level laser; each 15 minutes

ICD-10 CM Diagnosis Codes	Description
K12.31	Oral mucositis (ulcerative) due to antineoplastic therapy
K12.33	Oral mucositis (ulcerative) due to radiation
K12.39	Other oral mucositis (ulcerative)

POLICY TITLE	LOW-LEVEL LASER THERAPY
POLICY NUMBER	MP- 1.097

IX. REFERENCE

[TOP](#)

1. Lalla RV, Bowen J, Barasch A, et al. MASCC/ISOO clinical practice guidelines for the management of mucositis secondary to cancer therapy. *Cancer*. May 15 2014;120(10):1453-1461. PMID 24615748
2. Schubert MM, Eduardo FP, Guthrie KA, et al. A phase III randomized double-blind placebo-controlled clinical trial to determine the efficacy of low level laser therapy for the prevention of oral mucositis in patients undergoing hematopoietic cell transplantation. *Support Care Cancer*. Oct 2007;15(10):1145-1154. PMID 17393191
3. Figueiredo AL, Lins L, Cattony AC, et al. Laser therapy in the control of oral mucositis: a meta-analysis. *Rev Assoc Med Bras (1992)*. Sep-Oct 2013;59(5):467-474. PMID 24119379
4. Oberoi S, Zamperlini-Netto G, Beyene J, et al. Effect of prophylactic low level laser therapy on oral mucositis: a systematic review and meta-analysis. *PLoS One*. Sep 2014;9(9):e107418. PMID 25198431
5. Doeuk C, Hersant B, Bosc R, et al. Current indications for low level laser treatment in maxillofacial surgery: a review. *Br J Oral Maxillofac Surg*. Apr 2015;53(4):309-315. PMID 25740083
6. Gautam AP, Fernandes DJ, Vidyasagar MS, et al. Low level laser therapy for concurrent chemoradiotherapy induced oral mucositis in head and neck cancer patients - A triple blinded randomized controlled trial. *Radiother Oncol*. Sep 2012;104(3):349-354. PMID 22884841
7. Gautam AP, Fernandes DJ, Vidyasagar MS, et al. Low Level Helium Neon Laser therapy for chemoradiotherapy induced oral mucositis in oral cancer patients - A randomized controlled trial. *Oral Oncol*. Sep 2012;48(9):893-897. PMID 22502814
8. Gautam AP, Fernandes DJ, Vidyasagar MS, et al. Effect of low-level laser therapy on patient reported measures of oral mucositis and quality of life in head and neck cancer patients receiving chemoradiotherapy--a randomized controlled trial. *Support Care Cancer*. May 2013;21(5):1421-1428. PMID 23224689
9. Gautam AP, Fernandes DJ, Vidyasagar MS, et al. Low level laser therapy against radiation induced oral mucositis in elderly head and neck cancer patients-a randomized placebo controlled trial. *J Photochem Photobiol B*. Mar 2015;144:51-56. PMID 25704314
10. Oton-Leite AF, Silva GB, Morais MO, et al. Effect of low-level laser therapy on chemoradiotherapy-induced oral mucositis and salivary inflammatory mediators in head and neck cancer patients. *Lasers Surg Med*. Apr 2015;47(4):296-305. PMID 25824475
11. Ferreira B, da Motta Silveira FM, de Orange FA. Low-level laser therapy prevents severe oral mucositis in patients submitted to hematopoietic stem cell transplantation: a randomized clinical trial. *Support Care Cancer*. Mar 2016;24(3):1035-1042. PMID 26248655

POLICY TITLE	LOW-LEVEL LASER THERAPY
POLICY NUMBER	MP- 1.097

12. *Blue Cross and Blue Shield Technology Evaluation Center (TEC). Low-level laser therapy for carpal tunnel syndrome and chronic neck pain. TEC Assessment. Nov 2010;Volume 25:Tab 4. PMID 21638940*
13. *Li ZJ, Wang Y, Zhang HF, et al. Effectiveness of low-level laser on carpal tunnel syndrome: A meta-analysis of previously reported randomized trials. Medicine (Baltimore). Aug 2016;95(31):e4424. PMID 27495063*
14. *Fusakul Y, Aranyavalai T, Saensri P, et al. Low-level laser therapy with a wrist splint to treat carpal tunnel syndrome: a double-blinded randomized controlled trial. Lasers Med Sci. May 2014;29(3):1279-1287. PMID 24477392*
15. *Chow RT, Heller GZ, Barnsley L. The effect of 300 mW, 830 nm laser on chronic neck pain: a double-blind, randomized, placebo-controlled study. Pain. Sep 2006;124(1-2):201-210. PMID 16806710*
16. *Gross AR, Dziengo S, Boers O, et al. Low level laser therapy (LLLT) for neck pain: a systematic review and meta-regression. Open Orthop J. Oct 2013;7:396-419. PMID 24155802*
17. *Yeldan I, Cetin E, Ozdincler AR. The effectiveness of low-level laser therapy on shoulder function in subacromial impingement syndrome. Disabil Rehabil. Nov 2009;31(11):935-940. PMID 19031167*
18. *Dogan SK, Ay S, Evcik D. The effectiveness of low laser therapy in subacromial impingement syndrome: a randomized placebo controlled double-blind prospective study. Clinics (Sao Paulo). Dec 2010;65(10):1019-1022. PMID 21120304*
19. *Abrisham SM, Kermani-Alghoraishi M, Ghahramani R, et al. Additive effects of low-level laser therapy with exercise on subacromial syndrome: a randomised, double-blind, controlled trial. Clin Rheumatol. May 4 2011;30(10):1341-1346. PMID 21538218*
20. *Bal A, Eksioglu E, Gurcay E, et al. Low-level laser therapy in subacromial impingement syndrome. Photomed Laser Surg. Feb 2009;27(1):31-36. PMID 19250050*
21. *Calis HT, Berberoglu N, Calis M. Are ultrasound, laser and exercise superior to each other in the treatment of subacromial impingement syndrome? A randomized clinical trial. Eur J Phys Rehabil Med. Mar 2 2011;47(3):375-380. PMID 21364511*
22. *Page MJ, Green S, Kramer S, et al. Electrotherapy modalities for adhesive capsulitis (frozen shoulder). Cochrane Database Syst Rev. Oct 1 2014;10:CD011324. PMID 25271097*
23. *Stergioulas A. Low-power laser treatment in patients with frozen shoulder: preliminary results. Photomed Laser Surg. Apr 2008;26(2):99-105. PMID 18341417*
24. *Chen J, Huang Z, Ge M, et al. Efficacy of low-level laser therapy in the treatment of TMDs: a meta-analysis of 14 randomised controlled trials. J Oral Rehabil. Apr 2015;42(4):291-299. PMID 25491183*
25. *Chang WD, Lee CL, Lin HY, et al. A meta-analysis of clinical effects of low-level laser therapy on temporomandibular joint pain. J Phys Ther Sci. Aug 2014;26(8):1297-1300. PMID 25202201*

POLICY TITLE	LOW-LEVEL LASER THERAPY
POLICY NUMBER	MP- 1.097

26. Shobha R, Narayanan VS, Jagadish Pai BS, et al. Low-level laser therapy: A novel therapeutic approach to temporomandibular disorder - A randomized, double-blinded, placebo-controlled trial. *Indian J Dent Res.* Jul-Aug 2017;28(4):380-387. PMID 28836528
27. Glazov G, Yelland M, Emery J. Low-level laser therapy for chronic non-specific low back pain: a meta-analysis of randomised controlled trials. *Acupunct Med.* Oct 2016;34(5):328-341. PMID 27207675
28. Huang Z, Ma J, Chen J, et al. The effectiveness of low-level laser therapy for nonspecific chronic low back pain: a systematic review and meta-analysis. *Arthritis Res Ther.* Dec 15 2015;17:360. PMID 26667480
29. Koldas Dogan S, Ay S, Evcik D. The effects of two different low level laser therapies in the treatment of patients with chronic low back pain: A double-blinded randomized clinical trial. *J Back Musculoskelet Rehabil.* Jul2017;30(2):235-240. PMID 27472858
30. Huang Z, Chen J, Ma J, et al. Effectiveness of low-level laser therapy in patients with knee osteoarthritis: a systematic review and meta-analysis. *Osteoarthritis Cartilage.* Sep 2015;23(9):1437-1444. PMID 25914044
31. Bjordal JM, Johnson MI, Lopes-Martins RA, et al. Short-term efficacy of physical interventions in osteoarthritic knee pain. A systematic review and meta-analysis of randomised placebo-controlled trials. *BMC Musculoskelet Disord.* Jun 22 2007;8:51. PMID 17587446
32. Stergioulas A, Stergioula M, Aarskog R, et al. Effects of low-level laser therapy and eccentric exercises in the treatment of recreational athletes with chronic achilles tendinopathy. *Am J Sports Med.* May 2008;36(5):881-887. PMID 18272794
33. Tumilty S, McDonough S, Hurley DA, et al. Clinical effectiveness of low-level laser therapy as an adjunct to eccentric exercise for the treatment of Achilles' tendinopathy: a randomized controlled trial. *Arch Phys Med Rehabil.* May 2012;93(5):733-739. PMID 22541305
34. Macias DM, Coughlin MJ, Zang K, et al. Low-level laser therapy at 635 nm for treatment of chronic plantar fasciitis: a placebo-controlled, randomized study. *J Foot Ankle Surg.* Sep-Oct 2015;54(5):768-772. PMID 25769363
35. Kiritsi O, Tsitas K, Malliaropoulos N, et al. Ultrasonographic evaluation of plantar fasciitis after low-level laser therapy: results of a double-blind, randomized, placebo-controlled trial. *Lasers Med Sci.* Mar 2010;25(2):275-281. PMID 19841862
36. Cinar E, Saxena S, Uygur F. Low-level laser therapy in the management of plantar fasciitis: a randomized controlled trial. *Lasers Med Sci.* Dec 23 2017. PMID 29273892
37. Brosseau L, Robinson V, Wells G, et al. Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis. *Cochrane Database Syst Rev.* Oct 19 2005(4):CD002049. PMID 16235295
38. Meireles SM, Jones A, Jennings F, et al. Assessment of the effectiveness of low-level laser therapy on the hands of patients with rheumatoid arthritis: a randomized double-blind controlled trial. *Clin Rheumatol.* May 2010;29(5):501-509. PMID 20082104

POLICY TITLE	LOW-LEVEL LASER THERAPY
POLICY NUMBER	MP- 1.097

39. Alayat MS, Elsodany AM, El Fiky AA. Efficacy of high and low level laser therapy in the treatment of Bell's palsy: a randomized double blind placebo-controlled trial. *Lasers Med Sci.* Jan 2014;29(1):335-342. PMID 23709010
40. Ordahan B, Karahan AY. Role of low-level laser therapy added to facial expression exercises in patients with idiopathic facial (Bell's) palsy. *Lasers Med Sci.* May 2017;32(4):931-936. PMID 28337563
41. Ruaro JA, Frez AR, Ruaro MB, et al. Low-level laser therapy to treat fibromyalgia. *Lasers Med Sci.* Nov 2014;29(6):1815-1819. PMID 24801056
42. Matsutani LA, Marques AP, Ferreira EA, et al. Effectiveness of muscle stretching exercises with and without laser therapy at tender points for patients with fibromyalgia. *Clin Exp Rheumatol.* May-Jun 2007;25(3):410-415. PMID 17631737
43. Samson D, Lefevre F, Aronson N. Wound-healing technologies: low-level laser and vacuum-assisted closure. *Evid Rep Technol Assess (Summ).* Dec 2004(111):1-6. PMID 15663354
44. Chen C, Hou WH, Chan ES, et al. Phototherapy for treating pressure ulcers. *Cochrane Database Syst Rev.* 2014;7:CD009224. PMID 25019295
45. Machado RS, Viana S, Sbruzzi G. Low-level laser therapy in the treatment of pressure ulcers: systematic review. *Lasers Med Sci.* May 2017;32(4):937-944. PMID 28116536
46. Taradaj J, Halski T, Kucharzewski M, et al. Effect of laser irradiation at different wavelengths (940, 808, and 658 nm) on pressure ulcer healing: results from a clinical study. *Evid Based Complement Alternat Med* 2013;2013:960240. PMID 24111111
47. Lucas C, van Gemert MJ, de Haan RJ. Efficacy of low-level laser therapy in the management of stage III decubitus ulcers: a prospective, observer-blinded multicentre randomised clinical trial. *Lasers Med Sci.* 2003;18(2):72-77. PMID 12928815
48. Nussbaum EL, Biemann I, Mustard B. Comparison of ultrasound/ultraviolet-C and laser for treatment of pressure ulcers in patients with spinal cord injury. *Phys Ther.* Sep 1994;74(9):812-823; discussion 824-815. PMID 8066108
49. Taly AB, Sivaraman Nair KP, Murali T, et al. Efficacy of multiwavelength light therapy in the treatment of pressure ulcers in subjects with disorders of the spinal cord: A randomized double-blind controlled trial. *Arch Phys Med Rehabil.* Oct 2004;85(10):1657-1661. PMID 15468027
50. Smoot B, Chiavola-Larson L, Lee J, et al. Effect of low-level laser therapy on pain and swelling in women with breast cancer-related lymphedema: a systematic review and meta-analysis. *J Cancer Surviv.* Jun 2015;9(2):287-304. PMID 25432632
51. Omar MT, Shaheen AA, Zafar H. A systematic review of the effect of low-level laser therapy in the management of breast cancer-related lymphedema. *Support Care Cancer.* Aug 9 2012;20:2977-2984. PMID 22875413
52. Omar MTA, Ebid AA, El Morsy AM. Treatment of post-mastectomy lymphedema with laser therapy: double blind placebo control randomized study. *J Surg Res.* Jan 2011;165(1):82-90. PMID 20538293

POLICY TITLE	LOW-LEVEL LASER THERAPY
POLICY NUMBER	MP- 1.097

53. Sung L, Robinson P, Treister N, et al. Guideline for the prevention of oral and oropharyngeal mucositis in children receiving treatment for cancer or undergoing hematopoietic stem cell transplantation. *BMJ Support Palliat Care*. Mar 2017;7(1):7-16. PMID 25818385
54. Carcia CR, Martin RL, Houck J, et al. Achilles pain, stiffness, and muscle power deficits: Achilles tendinitis. *J Orthop Sports Phys Ther*. Sep 2010;40(9):A1-26. PMID 20805627
55. News Now Staff. Laser Therapy a Slight Winner in Analysis of Electrotherapy Modalities for Adhesive Capsulitis PT in Motion News 2014; <http://www.apta.org/PTinMotion/News/2014/10/3/LaserTherapyCapsulitis> Accessed May 8, 2020.
56. National Institute for Health and Care Excellence (NICE). Low back pain in adults: early management [CG88]. 2009; <https://www.nice.org.uk/guidance/CG88> . Accessed May 8, 2020.
57. National Institute for Health and Care Excellence (NICE). Low back pain and sciatica in over 16s: assessment and management [NG59]. 2016; <https://www.nice.org.uk/guidance/NG59/chapter/Recommendations> . Accessed May 8, 2020.
58. Chou R, Qaseem A, Snow V, et al. Diagnosis and treatment of low back pain: a joint clinical practice guideline from the American College of Physicians and the American Pain Society. *Ann Intern Med*. Oct 2 2007;147(7):478-491. PMID 17909209
59. Chou R, Loeser JD, Owens DK, et al. Interventional therapies, surgery, and interdisciplinary rehabilitation for low back pain: an evidence-based clinical practice guideline from the American Pain Society. *Spine (Phila Pa 1976)*. May 1 2009;34(10):1066-1077. PMID 19363457
60. Qaseem A, Wilt TJ, McLean RM, et al. Noninvasive treatments for acute, subacute, and chronic low back pain: a clinical practice guideline from the American College of Physicians. *Ann Intern Med*. Apr 4 2017;166(7):514-530. PMID 28192789
61. American Academy of Orthopaedic Surgeons. Management of Carpal Tunnel Syndrome: Evidence-Based Clinical Guideline.2016; <https://aaos.org/globalassets/quality-and-practice-resources/carpal-tunnel/management-of-carpal-tunnel-syndrome-7-31-19.pdf> Accessed May 8, 2020.
62. Wang W, Jiang W, Tang C, et al. Clinical efficacy of low-level laser therapy in plantar fasciitis: A systematic review and meta-analysis. *Medicine (Baltimore)*. 2019 Jan;98(3):e14088. PMID: 30653125
63. Honda Y, Sakamoto J, Hamaue Y, et al. Effects of Physical-Agent Pain Relief Modalities for Fibromyalgia Patients: A Systematic Review and Meta-Analysis of Randomized Controlled Trials. *Pain Res Manag*. 2018 Oct 1;2018:2930632. PMID: 30402199
64. Blue Cross Blue Shield Association Medical Policy Reference Manual. 2.01.56, Low-Level Laser Therapy. June 13, 2019.

MEDICAL POLICY

POLICY TITLE	LOW-LEVEL LASER THERAPY
POLICY NUMBER	MP- 1.097

X. POLICY HISTORY

[TOP](#)

MP-1.097	CAC 10/28/03
	CAC 5/31/05
	CAC 2/28/06
	CAC 2/27/07
	CAC 1/29/08
	CAC 1/27/09
	CAC 1/26/10 Consensus review.
	CAC 4/26/11 Consensus review.
	CAC 6/26/12 Consensus review. BCBSA Background/Description adopted for low-level laser therapy. No change to policy statements, references updated.
	7/26/13 Admin coding review complete
	CAC 9/24/13 Consensus review. No change to policy statements. References updated.
	CAC 9/30/14 Consensus review. No change to policy statements. Rationale added for low level laser therapy. References updated.
	CAC 9/29/15 Consensus review. No change to the policy statements. Reference and rationale update. Coding Review
	CAC 5/31/16 Minor revision. BCBSA adopted for this review. Policy title revised to “Low Level Laser Therapy”. Statement added that low-level laser therapy may be considered medically necessary for prevention of oral mucositis in selected patients. Additional bullet points to the investigational statement and the statement was changed to investigational for “all other indications”. High power Class IV therapeutic laser light lasers removed from policy. Background, rationale, and references updated. FEP variation revised. Coding reviewed.
	Administrative Update 11/22/16 Variation section reformatted.
	CAC 7/25/17 Consensus review. Policy statements unchanged. Medicare variation to NCD 270.6 added. Policy Guidelines, Description/Background, Rationale and Reference sections updated. Coding reviewed.
	1/1/18 Admin Update: Medicare variations removed from Commercial Policies.
	2/28/18 Admin coding review. No changes.
	6/13/18 Consensus review. No changes to the policy statements. Background and references updated. Rationale revised.
	05/08/2019 Consensus review. No changes to the policy statements. References reviewed and updated. New code 0552T added as investigational.
04/14/2020 Admin Update: Correct procedure code 0052T to read 0552T	
5/8/2020 Consensus review. Background, Rationale and coding reviewed. References added. No change to policy statement.	

MEDICAL POLICY



POLICY TITLE	LOW-LEVEL LASER THERAPY
POLICY NUMBER	MP- 1.097

[Top](#)

.Health care benefit programs issued or administered by Capital BlueCross and/or its subsidiaries, Capital Advantage Insurance Company®, Capital Advantage Assurance Company® and Keystone Health Plan® Central. Independent licensees of the BlueCross BlueShield Association. Communications issued by Capital BlueCross in its capacity as administrator of programs and provider relations for all companies